

A bipartite Sachdev-Ye-Kitaev model: Conformal limit and level statistics

Mikael Fremling,¹ Masudul Haque,^{2,3} and Lars Fritz¹

¹*Institute for Theoretical Physics and Center for Extreme Matter and Emergent Phenomena, Utrecht University, Princetonplein 5, 3584 CC Utrecht, The Netherlands*

²*Institut für Theoretische Physik, Technische Universität Dresden, 01062 Dresden, Germany*

³*Department of Theoretical Physics, Maynooth University, Co. Kildare, Ireland*

We study a bipartite version of the Sachdev-Ye-Kitaev (SYK) model. We show that the model remains solvable in the limit of large- N in the same sense as the original model if the ratio of both flavors is kept finite. The scaling dimensions of the two species can be tuned continuously as a function of the ratio. We also investigate the finite-size spectral properties of the model. We show how the level statistics differs from the original SYK model and infer an additional exchange symmetry in the bipartite model.

I. INTRODUCTION

The Sachdev-Ye-Kitaev (SYK) model [1–5] describes a system with many degrees of freedom with random all-to-all (q -body) interactions. The original model of Sachdev and Ye consists of pairwise coupled $SU(M)$ spins [1]. The more recent version proposed by Kitaev [3] has $N_\chi \gg 1$ Majorana sites. The $q = 4$ version has the Hamiltonian

$$H_{\text{SYK}} = \frac{1}{4!} \sum_{i,j,l,m} J_{ijkl} \gamma_i \gamma_j \gamma_l \gamma_m . \quad (1)$$

with N_χ localized Majorana fermions γ_i with $i = 1, \dots, N_\chi$. The term SYK is also used to refer to complex-fermion versions of this model and models with q -body interactions, with q taking values other than 4. In this work, we will restrict to the Majorana version (1) with four-body interactions. The Majorana degrees of freedom have no kinetic energy in this setup; in fact, since the interactions are all-to-all, the system has zero spatial dimensions. The interactions are usually taken to be Gaussian with mean $\langle J_{ijkl} \rangle = 0$ and variance

$$\langle J_{ijkl} J_{i'j'l'm'} \rangle = \frac{6J^2}{N_\chi^3} \delta_{i,i'} \delta_{j,j'} \delta_{l,l'} \delta_{m,m'} .$$

The SYK model has been studied intensely in the last few years, and has a number of fascinating properties. It is a strongly coupled quantum many-body system that is maximally chaotic, as evidenced by a maximal Lyapunov exponent extracted from out-of-time-ordered correlators, and hence acts as a fast scrambler of quantum information [4–6]. It is nearly conformally invariant, and is exactly solvable in the large N_χ limit [4, 7–10]. It has been used to describe two dimensional gravity and black holes [2–4, 9, 11, 12]. The SYK model and its extensions have also been used as a mean field model for non-Fermi liquids, and metals without quasiparticles [13–19].

The subject of this work is a variant of the SYK model, which we henceforth refer to as the bipartite SYK (b-SYK) model. The b-SYK was reported recently in Ref. [20] by two of the present authors to arise as the




FIG. 1. Graphical representation of the b-SYK model. Two sets of Majorana fermions, A and B , do not interact within the set but strongly interact between sets.

effective low-energy model in finite-size strained Kitaev honeycomb systems in the presence of the so-called Γ -term and moderate disorder. The b-SYK consists of two sets of Majorana fermions, A and B , with random 4-body interaction terms that each involve exactly two Majorana fermions from A and two from B . The difference with the standard SYK model is that there are no interactions within each set, only between the sets. This is illustrated in a sketch in Fig. 1.

The b-SYK model Hamiltonian is

$$H_{\text{b-SYK}} = \frac{1}{4} \sum_{i,j=1}^{N_A} \sum_{l,m=1}^{N_B} J_{ijkl} a_i^A a_j^A a_l^B a_m^B , \quad (2)$$

where a^A is a Majorana fermion in set A whereas a^B is one in set B . There are N_A and N_B fermions in the two sets, respectively. The distribution of the couplings follows

$$\langle J_{ijkl} J_{i'j'l'm'} \rangle = \frac{J^2}{2\sqrt{N_A N_B}} \delta_{i,i'} \delta_{j,j'} \delta_{l,l'} \delta_{m,m'} .$$

We show that the b-SYK model has an asymptotic conformal symmetry in the large- N_χ limit with tunable scaling dimensions — the scaling dimensions are

a function of the relative sizes of the A and B sets. By exploring the level statistics in finite-size realizations of the system, we infer that the b-SYK system has an additional Z_2 symmetry compared to the SYK system when the two sets contain equal numbers of Majorana fermions.

The organization of the paper is as follows. In Sec. II we study the large- N_χ limit of the theory: in Sec. II A we consider the two-point correlator and the scaling dimensions, whereas in Sec. II B we present the mean-field free energy. In Sec. III we study the level statistics of the b-SYK Hamiltonian and of the Hamiltonian interpolating between b-SYK and SYK. The findings of the paper and their context are discussed in Sec. IV

II. SOME ANALYTICAL PROPERTIES IN THE CONFORMAL LIMIT

A. The two point correlation function

One of the features of the SYK model is that it shows conformal invariance in the infrared in the large- N_χ limit. This allows for an asymptotically exact solution of its correlation functions [4]. We find that the emerging conformal symmetry of the SYK model also carries over to the b-SYK model if the large- N_χ limit is taken in a specific way: the conformal symmetry exists in the limit $N_A, N_B \rightarrow \infty$ as long as the ratio $N_A/N_B = \kappa$

is kept constant. Consequently, instead of having one scaling dimension of the Majorana fermions, like in the SYK model, the two sets of Majorana fermions, A and B , generally have different scaling dimensions, Δ_A , and Δ_B . Their scaling dimensions depend on the parameter κ , and they can assume values between 0 and 1/2 while $\Delta_A + \Delta_B = 1/2$. To demonstrate this, we first define the imaginary time-ordered correlation functions

$$\begin{aligned} G_{ij}^{AA}(\tau) &= \langle T_\tau (a_i^A(\tau) a_j^A(0)) \rangle ; \\ G_{ij}^{BB}(\tau) &= \langle T_\tau (a_i^B(\tau) a_j^B(0)) \rangle ; \\ G_{ij}^{AB}(\tau) &= \langle T_\tau (a_i^A(\tau) a_j^B(0)) \rangle ; \\ G_{ij}^{BA}(\tau) &= \langle T_\tau (a_i^B(\tau) a_j^A(0)) \rangle . \end{aligned} \quad (3)$$

The Green function of the non-interacting problem is given by

$$\begin{aligned} G_{0,ij}^{AA}(\tau) &= \frac{1}{2} \text{sgn}(\tau) \delta_{i,j} , \\ G_{0,ij}^{BB}(\tau) &= \frac{1}{2} \text{sgn}(\tau) \delta_{i,j} \\ G_{0,ij}^{AB}(\tau) &= G_{0,ij}^{BA}(\tau) = 0 . \end{aligned} \quad (4)$$

meaning it is local in the index i, j as well as the set label A, B . It constitutes the starting point for the perturbation theory to follow. The most general Dyson equation reads

$$\int d\tau' \begin{pmatrix} G_{0,ij}^{AA-1}(\tau, \tau') - \Sigma_{ij}^{AA}(\tau, \tau') & G_{0,ij}^{AB-1}(\tau, \tau') - \Sigma_{ij}^{AB}(\tau, \tau') \\ G_{0,ij}^{BA-1}(\tau, \tau') - \Sigma_{ij}^{BA}(\tau, \tau') & G_{0,ij}^{BB-1}(\tau, \tau') - \Sigma_{ij}^{BB}(\tau, \tau') \end{pmatrix} \begin{pmatrix} G_{jk}^{AA}(\tau', \tau'') & G_{jk}^{AB}(\tau', \tau'') \\ G_{jk}^{BA}(\tau', \tau'') & G_{jk}^{BB}(\tau', \tau'') \end{pmatrix} = \delta(\tau - \tau'') \delta_{i,k} \mathbb{1} \quad (5)$$

where Σ^{AA} , Σ^{AB} , Σ^{BA} , and Σ^{BB} are the self-energies whereas G^{AA} , G^{AB} , G^{BA} , and G^{BB} are the Green functions. Summation over double indices is implied. In general, this equation is non-local in both the indices i, j as well as the set labels A, B . The most transparent way to determine the self-energies is based on a diagrammatic representation of perturbation theory in terms of Feynman diagrams. To leading order in N_A and N_B , the diagrams shown in Fig. 2 constitute the entire per-

turbative series and can be resummed exactly. This implies that in this limit, the theory remains local in i, j as well as A, B . Consequently, to leading order the off-diagonal self-energies $\Sigma_{ii}^{AB}(\tau_1, \tau_2)$ as well as the off-diagonal Green functions G^{AB} and G^{BA} vanish. Furthermore, as shown explicitly below, the self-energies $\Sigma_{ii}^{AA}(\tau_1, \tau_2)$ and $\Sigma_{ii}^{BB}(\tau_1, \tau_2)$ dominate the bare propagators, Eq. (4), in the infrared. This implies we have to solve

$$\int d\tau' \begin{pmatrix} \Sigma_{ii}^{AA}(\tau, \tau') G_{ii}^{AA}(\tau', \tau'') & 0 \\ 0 & \Sigma_{ii}^{BB}(\tau, \tau') G_{ii}^{BB}(\tau', \tau'') \end{pmatrix} = -\delta(\tau - \tau'') \mathbb{1} . \quad (6)$$

Since we have translational invariance in time all quantities only contain relative coordinates, so that

$$\begin{aligned} \Sigma_{ii}^{AA}(\tau) &= \frac{J^2 N_A N_B^2}{\sqrt{N_A^3 N_B^3}} G_{ii}^{AA}(\tau) G_{i'i'}^{BB}(\tau) G_{i'i'}^{BB}(\tau) = \frac{J^2}{\sqrt{\kappa}} G_{ii}^{AA}(\tau) G_{i'i'}^{BB}(\tau) G_{i'i'}^{BB}(\tau) \\ \Sigma_{ii}^{BB}(\tau) &= \frac{J^2 N_A^2 N_B}{\sqrt{N_A^3 N_B^3}} G_{ii}^{BB}(\tau) G_{i'i'}^{AA}(\tau) G_{i'i'}^{AA}(\tau) = J^2 \sqrt{\kappa} G_{ii}^{BB}(\tau) G_{i'i'}^{AA}(\tau) G_{i'i'}^{AA}(\tau) . \end{aligned} \quad (7)$$




FIG. 2. The straight line denotes the propagator G^{AA} whereas the wiggly line is G^{BB} . The upper line shows the Dyson equation for G^{AA} and the second line the Dyson equation for G^{BB} , compare Eq. (6). The lowest line shows the approximation for the self-energies Σ^{AA} and Σ^{BB} , Eq. (7), which becomes exact in the limit of large N_A and N_B .

Due to the time reparametrization symmetry of the theory we expect conformal invariance. Since we are expecting different scaling dimensions for the Majorana fermions in the two sets we introduce the scaling dimension Δ_A for Majorana fermions in set A , whereas we introduce Δ_B for those in set B . We then have

$$\begin{aligned} G^{AA}(\tau) &= A \frac{\text{sgn}(\tau)}{|\tau|^{2\Delta_A}} \\ G^{BB}(\tau) &= B \frac{\text{sgn}(\tau)}{|\tau|^{2\Delta_B}}, \end{aligned} \quad (8)$$

for the full Green functions. To decouple the Dyson equation, Eq. (6), one has to Fourier transform it. A recurring Fourier transform is of the type

$$\int_{-\infty}^{\infty} d\tau e^{i\omega\tau} \frac{\text{sgn}(\tau)}{|\tau|^\alpha} = i 2^{1-\alpha} \sqrt{\pi} \frac{\Gamma(1-\frac{\alpha}{2})}{\Gamma(\frac{1}{2}+\frac{\alpha}{2})} \text{sgn}(\omega) |\omega|^{\alpha-1}. \quad (9)$$

In combination with Eq. (6) it leads to two conditions on the scaling dimensions Δ_A and Δ_B :

$$\begin{aligned} \frac{1}{2} &= \Delta_A + \Delta_B \\ \kappa &= \frac{2\Delta_A}{1-2\Delta_A} \left(\frac{1}{\tan(\pi\Delta_A)} \right)^2. \end{aligned} \quad (10)$$

For $\kappa = 1$, we find $\Delta_A = \Delta_B = 1/4$, as expected, just like in the standard SYK model. For other values of κ , both scaling dimensions interpolate between 0 and $1/2$ while always fulfilling $\Delta_A + \Delta_B = 1/2$. This behavior is presented in Fig. 3 on a logarithmic scale which shows

the A - B symmetry explicitly. Tunable scaling dimensions have already been seen in other variants of the SYK model e.g. Ref. [21, 22].

Due to the conformal invariance and the reparametrization invariance it is straightforward to determine the finite temperature and real time




FIG. 3. Scaling dimensions Δ_A and Δ_B as a function of κ . Both interpolate between 0 and $1/2$.

correlators. At finite temperatures we find

$$\begin{aligned} G^{AA}(\tau) &= A \text{sgn}(\tau) \left(\frac{\pi}{\beta \sinh(\frac{\pi\tau}{\beta})} \right)^{2\Delta_A}, \\ G^{BB}(\tau) &= B \text{sgn}(\tau) \left(\frac{\pi}{\beta \sinh(\frac{\pi\tau}{\beta})} \right)^{2\Delta_B}, \end{aligned} \quad (11)$$

whereas for the retarded propagator at finite temperature we obtain

$$\begin{aligned} G_{\text{ret}}^{AA}(t) &= \theta(t) A \cos(\pi\Delta_A) \left(\frac{\pi}{\beta \sinh(\frac{\pi t}{\beta})} \right)^{2\Delta_A}, \\ G_{\text{ret}}^{BB}(t) &= \theta(t) B \cos(\pi\Delta_B) \left(\frac{\pi}{\beta \sinh(\frac{\pi t}{\beta})} \right)^{2\Delta_B}. \end{aligned} \quad (12)$$

B. Free energy

There is a standard prescription to calculate the effective action of the present problem, outlined for instance in Ref. [4]. After some manipulations, the effective replica symmetric action of the b-SYK model is given by

$$\begin{aligned} \mathcal{S}_{\text{eff}} = & -\frac{N_A}{2} \log \det (\partial_\tau - \Sigma^{AA}) - \frac{N_B}{2} \log \det (\partial_\tau - \Sigma^{BB}) + \frac{1}{2} \int d\tau_1 d\tau_2 \frac{J^2 N_A^2 N_B^2}{2} (G^{AA}(\tau_1, \tau_2))^2 (G^{BB}(\tau_1, \tau_2))^2 \\ & + \frac{1}{2} \int d\tau_1 d\tau_2 (N_A \Sigma^{AA}(\tau_1, \tau_2) G^{AA}(\tau_1, \tau_2) + N_B \Sigma^{BB}(\tau_1, \tau_2) G^{BB}(\tau_1, \tau_2)) \end{aligned} \quad (13)$$

It is straightforward to verify that its saddle point reproduces the series of melon diagrams shown in Fig. 2. This expression can be used as starting point to determine thermodynamic properties of the model and higher-order correlators as well as the Lyapunov exponent.

III. LEVEL STATISTICS

In this section, we focus on the level spacing statistics of the b-SYK model. For this purpose, we consider finite N_A and N_B . We will concentrate on the case $\kappa = 1$, so that we use $N_A = N_B$ for even $N_\chi = N_A + N_B$ and $N_A = N_B \pm 1$ for odd N_χ .

Level statistics can help identify the existence of chaos (non-integrability) in quantum Hamiltonians, and also to distinguish between different symmetry classes. The interest in the SYK model is partly due to its being maximally chaotic. Therefore, eigenvalue statistics has been a widely used diagnostic for characterizing the SYK model [12, 23–27] and its various variants [26, 28–42]. A noteworthy feature of the SYK level statistics is that it depends on the number of Majorana fermions N_χ . We show below that the level statistics of the b-SYK model in the $N_A = N_B$ case is systematically shifted with respect to that of the standard SYK model, consistent with the presence of an extra Z_2 symmetry in the b-SYK system.

A. Relevant ensembles

The universality classes of random matrices that are relevant for us include the Gaussian Orthogonal Ensemble (GOE), the Gaussian Unitary Ensemble (GUE), and the Gaussian Symplectic Ensemble (GSE). In Table I we refer to these as O, U, and S respectively for conciseness. Additionally, we will encounter below the level statistics obtained by merging the spectra of two GOE matrices; we refer to this as $2 \times \text{GOE}$, or for conciseness 2O in Table I.

For characterizing the level statistics with a single number, it has become common to use the average ratio of successive level spacings [43, 44]. One starts with calculating the finite size spectrum E_n , which are ordered from lowest to highest energy. The set of level spacings are defined as $s_n = E_{n+1} - E_n$. This allows to define the ratio

$$r_n = \frac{\min(s_n, s_{n-1})}{\max(s_n, s_{n-1})}. \quad (14)$$

$N_\chi \pmod{8}$	0	1	2	3	4	5	6	7
H_{SYK}	O	O	U	S	S	S	U	O
$H_{\text{b-SYK}}$	2O	2O	O	U	U	U	O	2O

TABLE I. Level statistics of the Majorana fermion SYK model, H_{SYK} as compared to $H_{\text{b-SYK}}$ in (2). Here O=GOE, U=GUE, S=GSE and 2O = $2 \times \text{GOE}$ are the different universal random matrix ensembles. Note how the level statistics of $H_{\text{b-SYK}}$ traces those of H_{SYK} but with a reduction in the symmetry classifications of one step such that S \rightarrow U \rightarrow O \rightarrow 2O.

Analyzing the statistics of this quantity r_n has advantages over the statistics of the bare level spacings s_n themselves. It bypasses the need to account for a varying density of states through unfolding procedures. In addition, the average of this quantity has characteristic values for the different ensembles, thus enabling one to distinguish symmetry classes without analyzing complete distributions.

For the Wigner-Dyson ensembles, the probability distributions of the ratio r are well-approximated by the surmise [44] $P(r) \propto (r + r^2)^\beta / (1 + r + r^2)^{1+3\beta/2}$ up to normalization, with $\beta = 1$ for GOE, $\beta = 2$ for GUE, and $\beta = 4$ for GSE. The averages of these distributions are found to be $\langle r \rangle_{\text{GOE}} \approx 0.53$, $\langle r \rangle_{\text{GUE}} \approx 0.60$, and $\langle r \rangle_{\text{GSE}} \approx 0.67$ [44].

Integrable (non-chaotic) Hamiltonians, which do not show level repulsion, generically have Poisson statistics, for which the level spacing ratio has probability distribution $P(r) = 2/(1+r)^2$ and mean value $\langle r \rangle = 2 \ln 2 - 1 \approx 0.39$. An integrable system can be thought of as having a large number of conserved quantities or quantum numbers. Therefore, adding one or a few conservation laws to a GOE system is expected to change the distribution to a form intermediate between the GOE and Poisson cases. The $2 \times \text{GOE}$ spectrum can be interpreted as that obtained when a GOE system acquires a single quantum number with two possible values, which splits the GOE spectrum into two sectors. Thus, we expect its level spacing distribution to be intermediate between Poisson and GOE distributions. Indeed, we find numerically, by merging the spectra of two GOE matrices, that the $2 \times \text{GOE}$ distribution has $\langle r \rangle \approx 0.425$, intermediate between the Poisson and GOE values. Some analytic formulas for the $2 \times \text{GOE}$ distribution were also provided in Ref. 45.

As discrete symmetries are common in quantum Hamiltonians, spectra formed out of two or more independent GOE or GUE components are the subject of longstanding interest in the quantum-chaos and random-matrix literature [34, 45–57]. Here, we will only




FIG. 4. Average level spacing ratio $\langle r \rangle$ for H_{SYK} and $H_{\text{b-SYK}}$, plotted against system size N_χ . For each system size, the value of $\langle r \rangle$ for $H_{\text{b-SYK}}$ is lower than that for H_{SYK} . The overall $\langle r \rangle$ vs N_χ behavior for $H_{\text{b-SYK}}$ mimics that for H_{SYK} but with a shifted classification. Thus instead of the SYK sequence we get a relative shift as $\text{O} \rightarrow 2\text{O}$, $\text{U} \rightarrow \text{O}$ and $\text{S} \rightarrow \text{U}$.

be concerned with the $2 \times \text{GOE}$ case because restricting the couplings of the SYK Hamiltonian to obtain the b-SYK Hamiltonian effectively adds a single Z_2 symmetry.

B. Level statistics of b-SYK

In the case of the SYK model, the random matrix ensemble describing the level statistics changes with the number of Majorana fermions N_χ [12, 24–26] as listed in Table I. This dependence is cyclic modulo 8 in N_χ . Here we will compare the level spacing statistics properties of the SYK model to that of the b-SYK model.

For this analysis we choose $N_\chi = 2N_a = 2N_b$ or $N_\chi = 2N_a + 1 = 2N_b - 1$ depending on the parity of N_χ .

Before discussing our results, we make some technical remarks. In order to obtain the correct level statistics for the SYK-model it is important that the vacuum is such that the N_χ Majorana fermions can form $N_f = \lfloor \frac{N_\chi}{2} \rfloor$ spinless fermions, e.g., by enforcing $\gamma_j |0\rangle = i\gamma_{N_\chi-j+1} |0\rangle$ for $j \leq N_f$. For the b-SYK model, we choose this constraint such that $a_j^A |0\rangle = \alpha a_j^B |0\rangle$. Furthermore, for the symplectic cases, the full spectrum has a two-fold degeneracy, and half of the energy levels must be pruned to get rid of this residual last symmetry from the spectrum. Finally, in order to satisfactorily resolve the level statistics, one needs to average over many realizations of the two models and aggregate the level spacing ratio statistics.

We quantify the level statistics by the average ratio $\langle r \rangle$, described above. Some results are summarized in Fig. 4, for both the SYK and the b-SYK Hamiltonians. For each N_χ , the averaging of the spacing ratio is per-




FIG. 5. Distribution of level spacing ratios for the b-SYK Hamiltonian with $N_\chi = 21, 25, 26$, corresponding to the three distinct classes listed in Table I. The distributions for $N_\chi = 21$ and $N_\chi = 26$ match well the expected surmise for the GUE and GOE classes. The third example, $N_\chi = 25$, closely follows the $2 \times \text{GOE}$ distribution. The GSE and Poissonian distributions are also shown as lines. None of the b-SYK distributions follow the GSE class (in contrast to the SYK case) or the Poissonian distribution.

formed over the spectra of many coupling realizations so that the results are sufficiently converged. In Fig. 4 the horizontal lines represent the average r values for different relevant ensembles, as discussed above.

We observe that the average spacing ratio of the b-SYK model is always lower than the average spacing ratio expected from the SYK model, irrespective of the size of the system. However, it follows the same 8-fold periodicity in the total number of Majorana fermions, N_χ . Compared to the SYK sequence, we find relative shifts $\text{O} \rightarrow 2\text{O}$, $\text{U} \rightarrow \text{O}$, $\text{S} \rightarrow \text{U}$, i.e., the GSE, GUE, and GOE get converted to GUE, GOE, and $2 \times \text{GOE}$ respectively. The shift is also seen by comparing the two rows of Table I.

Going beyond the average, in Figure 5 we show the full distributions (numerical histograms) of the level spacing ratio, for the $H_{\text{b-SYK}}$ Hamiltonian with $N_\chi = 21, 25, 26$. Clearly, the three classes follow the expected distributions for $2 \times \text{GOE}$, GOE, and GUE, shown as dotted lines. The GSE distribution is not obtained in the b-SYK system for any value of N_χ .

The shift in level statistics relative to SYK is clearly due to the restriction to bipartite interactions. The results are consistent with the explanation that the bipartite structure leads to an additional Z_2 symmetry of the Hamiltonian. A system of the GUE symmetry class, if endowed with an additional Z_2 symmetry, shows GOE level statistics [58–61]. This effect was discussed early in the context of single-particle (billiard) systems with a magnetic field [58, 59]. This system would naively be expected to have GUE statistics due to broken time-reversal symmetry. However, when reflection symmetry is present, the level statistics is of the GOE class. This phenomenon has also been observed in a many-body

system [62]. In the present case, the anti-unitary symmetry involved is not time, but the effect is the same: For $N_\chi = 10, 14, 18, 22 \dots$, the SYK level statistics are GUE, but the b-SYK level statistics are of GOE type.

For values of N_χ for which the level statistics if of GSE type, a corresponding effect is seen. The additional symmetry reduces the degree of level repulsion, and one obtains GUE statistics instead, as seen in Fig. 4 and Table I. A GSE to GUE shift due to a parity symmetry is discussed in Section 2.7 of Ref. [55]. Still, we do not know of another example in the literature involving a many-body Hamiltonian. The b-SYK spectrum retains the Kramers degeneracy; the level repulsion is between pairs of degenerate states.

C. The Z_2 symmetry

The numerical data implies that the b-SYK Hamiltonian possesses a Z_2 symmetry which is not present in the SYK model. One idea might be a label-flipping symmetry where the A and B labels are switched, such that the Majorana fermions in the A partition are moved to the B partition and vice-versa. However, any individual realization of the b-SYK class of Hamiltonians does not remain invariant under such a label-flipping operation. We have not been able to explicitly identify the Z_2 operation, which keeps a b-SYK Hamiltonian (but not an SYK Hamiltonian) invariant.

The b-SYK Hamiltonian admits a number of operations that leave the Hamiltonian isospectral, although not invariant: exchanging any one of the A -fermions with any one of the B -fermions leaves the Hamiltonian isospectral, i.e., amount to unitary operations. This emerges due to the restriction from SYK to b-SYK: For the SYK Hamiltonian, such operations are not isospectral. For both SYK and b-SYK, bi-partitioning the Majorana fermions into arbitrary halves and then exchanging the two halves is an isospectral (unitary) operation. The extra feature of the b-SYK is that exchanging a single A fermion with a single B fermion is also a unitary operation. At present, it is unclear whether these spectrum-preserving ('gauge') transformations in the b-SYK are related to the yet-to-be-identified Z_2 symmetry, which must keep the Hamiltonian invariant, not just isospectral.

D. Interpolation between $H_{\text{b-SYK}}$ to H_{SYK}

Since the level statistics classification is systematically shifted from H_{SYK} to $H_{\text{b-SYK}}$ this begs the question what one obtains for a mixture of the two Hamiltonians. We therefore define an interpolation Hamiltonian

$$H_{\text{Mix}} = (1 - \lambda) H_{\text{b-SYK}} + \lambda H_{\text{SYK}}$$

and investigate its level statistics as a function of λ . In the following analysis, we choose the coupling constants

J in (1) and (2) such that the variance of J_{ijlm} in both cases is unity for all system sizes. The main results are summarized in Fig. 6. In panel a) we demonstrate how for already a small SYK-mixing, $\lambda \leq 0.1$, results in a drift of the level statistics from the $H_{\text{b-SYK}}$ to the H_{SYK} random matrix class. This makes sense because, as soon as interactions are allowed which violate the bipartite restriction, the additional symmetry of the b-SYK Hamiltonian is lost. The crossover happens faster (at even smaller values of λ) for larger system sizes, indicating that, in the large- N_χ limit, an infinitesimal influence of H_{SYK} is enough to move the system into the lower-symmetry class of the un-restricted SYK Hamiltonian.

To quantify this size dependence of the $H_{\text{b-SYK}} \rightarrow H_{\text{SYK}}$ crossover, we fit $\langle r \rangle$ to the function

$$f(\lambda) = \langle r_{\text{SYK}} \rangle + (\langle r_{\text{b-SYK}} \rangle - \langle r_{\text{SYK}} \rangle) e^{-\frac{\lambda}{\lambda_C}}$$

such that $f(0) = \langle r_{\text{b-SYK}} \rangle$, and $f(\lambda \gg \lambda_C) \rightarrow \langle r_{\text{SYK}} \rangle$. λ_C is the value for which $f(\lambda_C) = \langle r_{\text{SYK}} \rangle$ to first order, i.e. $f(0) + \lambda_C f'(0) = \langle r_{\text{SYK}} \rangle$. This function captures the transition from $\langle r_{\text{b-SYK}} \rangle$ to $\langle r_{\text{SYK}} \rangle$ as a function of λ . The best fit is shown in dashed lines, and the numerical $\langle r \rangle$ is shown with statistical errors.

In panel c) we show $\frac{\langle r \rangle - \langle r_{\text{b-SYK}} \rangle}{\langle r_{\text{SYK}} \rangle - \langle r_{\text{b-SYK}} \rangle}$. The shift from $\langle r_{\text{b-SYK}} \rangle$ to $\langle r_{\text{SYK}} \rangle$ takes place at smaller values of λ if more Majorana fermions, N_χ , are involved. In panel h), this is further illustrated: we plot λ_C as a function of inverse system size $1/N_\chi$. Clearly, λ_C tends to zero in the large N_χ limit, quantifying the intuition that the shift of behavior happens at smaller λ for larger sizes.

In panels d)-g) we show a scaling analysis for $\langle r_{\text{b-SYK}} \rangle$ and $\langle r_{\text{SYK}} \rangle$, using the values at $\lambda = 0$ and $\lambda = 1$. For $\langle r_{\text{SYK}} \rangle$ (orange plus symbols), the large-size limit is consistent with the known symmetry classes, GOE, GUE, or GSE, based on the value of $N_\chi \bmod 8$ [12, 24–26]. For $\langle r_{\text{b-SYK}} \rangle$ (blue circles), the large-size limit is consistent with the values corresponding to $2 \times \text{GOE}$, GOE, or GUE. In panel d), focusing on the GSE value $\langle r \rangle \approx 0.67$, only $\lambda = 1$ data (values of $\langle r_{\text{SYK}} \rangle$) are visible. Similarly, in panel g), focusing on the $2 \times \text{GOE}$ value $\langle r \rangle \approx 0.425$, only $\lambda = 0$ data (values of $\langle r_{\text{b-SYK}} \rangle$) are visible.

IV. DISCUSSION & CONTEXT

This paper has studied a bipartite version of the quartic ($q = 4$) SYK model, which we call b-SYK. It consists of two flavors of Majorana fermions that interact between the sets, but not within — each quartic interaction term involves two Majorana fermions from one set and two from the other set. The model was motivated in Ref. [20] as being realizable in a specific setup of a strained version of the Kitaev honeycomb model.

Variants of the SYK model with two species of fermions have appeared previously, perhaps most prominently with the motivation of modeling eternal traversable wormholes using two quartic SYK models




FIG. 6. Level statistics when tuning between H_{b-SYK} and H_{SYK} Hamiltonians, with $\lambda = 0$ ($\lambda = 1$) corresponding to b-SYK (SYK). In panel a) the average spacing ratio $\langle r \rangle$ is plotted as a function of λ . Panel b) is the same data in the full range $\lambda \in [0, 1]$, so that one can see both limits, $\langle r_{b-SYK} \rangle$ to $\langle r_{SYK} \rangle$. The crossover from $\langle r_{b-SYK} \rangle$ to $\langle r_{SYK} \rangle$ is steeper for a larger number of Majoranas N_χ . Panel c) illustrates the N_χ dependence of the transition via the normalized quantity $\frac{\langle r \rangle - \langle r_{b-SYK} \rangle}{\langle r_{SYK} \rangle - \langle r_{b-SYK} \rangle}$. Panels d) - g) zoom in on the regions around the various values of $\langle r_{b-SYK} \rangle$ and $\langle r_{SYK} \rangle$ that are obtained. The colors on the vertical axes should help to match with the corresponding regions in panels a) or b). These plots d) - g) show that the large-size limits are consistent with the GSE, GUE, GOE, and 2xGOE classes. In panel a) a fit is performed to the function $\langle r_{SYK} \rangle + (\langle r_{b-SYK} \rangle - \langle r_{SYK} \rangle)e^{-\frac{\lambda}{\lambda_c}}$ and on panel h) the transition parameter λ_c is shown. The “critical” λ_c decreases with increasing N_χ .

with only quadratic interactions between them [32, 63–71]. In Ref. [22] the coupling between the two SYK clusters is quartic like ours. Since our b-SYK model has no internal coupling within the two sets, it may be regarded as an infinite-coupling limit of the model of Ref. [22], i.e., the limit in which the intra-set couplings can be neglected. Ref. [72] treats a complex-fermion version. Ref. [73] also considers two SYK clusters and quartic couplings between them, but the sizes of the two clusters are parametrically different, so that one acts as a bath for the other. Several other two-flavor or two-species SYK variants have also appeared in the literature [74–77].

We study the b-SYK model both analytically and numerically. We find that in the large- N limit, the model remains asymptotically solvable, showing conformal invariance in the infrared. We establish that if we keep the ratio between the flavors a variable, we can continuously tune the scaling dimension of the respective species between 0 and 1/2.

For finite system sizes, we analyze the level statis-

tics of the model numerically for $\kappa = 1$ ($N_A = N_B$ or $N_A = N_B \pm 1$) and compare it to the known level statistics of the SYK model. We find that the level statistics deviates systematically, consistent with the b-SYK model possessing an additional Z_2 symmetry. The GOE, GUE, and GSE level statistics of the SYK model are reduced to 2xGOE, GOE, and GUE classes.

Studying the interpolation between the two models, we find that, for finite sizes, the statistics evolve smoothly from the b-SYK to the SYK as a function of interpolating parameter λ .

In the quantum chaos literature and in random matrix theory, the GOE-GUE crossover has been studied repeatedly in various contexts [51, 55, 78–88]. In the present case, we have a GUE to GSE crossover, a GOE to GUE crossover, and a 2xGOE to GOE crossover, all in the same Hamiltonian, depending on the number of Majorana fermions, according to the Bott periodicity [12, 24–26]. In addition, unlike typical models studied in traditional quantum chaos or random matrix theory, we have a well-defined thermodynamic (large N_χ) limit.

It turns out that, in this limit, the crossover happens extremely rapidly, i.e., the b-SYK statistics is lost already for an infinitesimal mixture of SYK.

The present work opens up a number of new questions. Thermodynamic and thermalization properties, as well as higher-order correlation functions, and Lyapunov exponents, remain to be studied. It may be interesting to see how b-SYK physics is explicitly obtained in the large-interaction limit of the model of Ref. [22], and to investigate the behavior of its complex-fermion version. For the level statistics, the Z_2 symmetry that we have inferred for $\kappa = 1$ remains to be identified. In ad-

dition, the level statistics for unequal-sized bipartitions ($\kappa \neq 1$) also deserves exploration.

ACKNOWLEDGEMENTS

We would like to thank Philippe Corboz, Maria Hermanns, Lukas Janssen, Graham Kells, Tobias Meng, Subir Sachdev, and Matthias Vojta for useful discussions. This work is part of the D-ITP consortium, a program of the Netherlands Organisation for Scientific Research (NWO) that is funded by the Dutch Ministry of Education, Culture and Science (OCW).

-
- [1] S. Sachdev and J. Ye, Gapless spin-fluid ground state in a random quantum heisenberg magnet, *Phys. Rev. Lett.* **70**, 3339 (1993).
 - [2] S. Sachdev, Bekenstein-hawking entropy and strange metals, *Phys. Rev. X* **5**, 041025 (2015).
 - [3] A. Kitaev, A simple model of quantum holography, *KITP strings seminar and Entanglement 2015 program* (Feb. 12, April 7, and May 27, 2015) (2015).
 - [4] J. Maldacena and D. Stanford, Remarks on the Sachdev-Ye-Kitaev model, *Phys. Rev. D* **94**, 106002 (2016).
 - [5] J. Maldacena, S. H. Shenker, and D. Stanford, A bound on chaos, *Journal of High Energy Physics* **2016**, 106 (2016).
 - [6] B. Kobrin, Z. Yang, G. D. Kahanamoku-Meyer, C. T. Olund, J. E. Moore, D. Stanford, and N. Y. Yao, Many-Body Chaos in the Sachdev-Ye-Kitaev model, *Phys. Rev. Lett.* **126**, 030602 (2021).
 - [7] J. Polchinski and V. Rosenhaus, The spectrum in the Sachdev-Ye-Kitaev model, *Journal of High Energy Physics* **2016**, 1 (2016).
 - [8] D. J. Gross and V. Rosenhaus, All point correlation functions in SYK, *Journal of High Energy Physics* **2017**, 148 (2017).
 - [9] A. Kitaev and S. J. Suh, The soft mode in the Sachdev-Ye-Kitaev model and its gravity dual, *Journal of High Energy Physics* **2018**, 183 (2018).
 - [10] V. Rosenhaus, An introduction to the SYK model, *Journal of Physics A: Mathematical and Theoretical* **52**, 323001 (2019).
 - [11] K. Jensen, Chaos in AdS₂ holography, *Phys. Rev. Lett.* **117**, 111601 (2016).
 - [12] J. S. Cotler, G. Gur-Ari, M. Hanada, J. Polchinski, P. Saad, S. H. Shenker, D. Stanford, A. Streicher, and M. Tezuka, Black holes and random matrices, *Journal of High Energy Physics* **2017**, 118 (2017).
 - [13] R. A. Davison, W. Fu, A. Georges, Y. Gu, K. Jensen, and S. Sachdev, Thermoelectric transport in disordered metals without quasiparticles: The Sachdev-Ye-Kitaev models and holography, *Phys. Rev. B* **95**, 155131 (2017).
 - [14] H. Wang, A. L. Chudnovskiy, A. Gorsky, and A. Kamenev, Sachdev-Ye-Kitaev superconductivity: Quantum kuramoto and generalized richardson models, *Phys. Rev. Research* **2**, 033025 (2020).
 - [15] A. Altland, D. Bagrets, and A. Kamenev, Sachdev-Ye-Kitaev non-fermi-liquid correlations in nanoscopic quantum transport, *Phys. Rev. Lett.* **123**, 226801 (2019).
 - [16] Y. Wang and A. V. Chubukov, Quantum phase transition in the Yukawa-SYK model, *Phys. Rev. Research* **2**, 033084 (2020).
 - [17] I. Esterlis, H. Guo, A. A. Patel, and S. Sachdev, Large- N theory of critical Fermi surfaces, *Phys. Rev. B* **103**, 235129 (2021).
 - [18] M. Tikhonovskaya, H. Guo, S. Sachdev, and G. Tarnopolsky, Excitation spectra of quantum matter without quasiparticles. I. Sachdev-Ye-Kitaev models, *Phys. Rev. B* **103**, 075141 (2021).
 - [19] E. Lantagne-Hurtubise, V. Pathak, S. Sahoo, and M. Franz, Superconducting instabilities in a spinful Sachdev-Ye-Kitaev model, *Phys. Rev. B* **104**, L020509 (2021).
 - [20] M. Fremling and L. Fritz, Sachdev-Ye-Kitaev type physics in the strained Kitaev honeycomb model, *arXiv:2105.06119* (2021), *arXiv:2105.06119*.
 - [21] E. Marcus and S. Vandoren, A new class of SYK-like models with maximal chaos, *Journal of High Energy Physics* **2019**, 166 (2019).
 - [22] J. Kim, I. R. Klebanov, G. Tarnopolsky, and W. Zhao, Symmetry breaking in coupled SYK or tensor models, *Phys. Rev. X* **9**, 021043 (2019).
 - [23] A. M. García-García and J. J. M. Verbaarschot, Spectral and thermodynamic properties of the Sachdev-Ye-Kitaev model, *Phys. Rev. D* **94**, 126010 (2016).
 - [24] Y.-Z. You, A. W. W. Ludwig, and C. Xu, Sachdev-Ye-Kitaev model and thermalization on the boundary of many-body localized fermionic symmetry-protected topological states, *Phys. Rev. B* **95**, 115150 (2017).
 - [25] A. M. García-García and J. J. M. Verbaarschot, Analytical spectral density of the Sachdev-Ye-Kitaev model at finite N , *Phys. Rev. D* **96**, 066012 (2017).
 - [26] M. Haque and P. A. McClarty, Eigenstate thermalization scaling in Majorana clusters: From chaotic to integrable Sachdev-Ye-Kitaev models, *Phys. Rev. B* **100**, 115122 (2019).
 - [27] J. Behrends, J. H. Bardarson, and B. Béri, Tenfold way and many-body zero modes in the sachdev-ye-kitaev model, *Phys. Rev. B* **99**, 195123 (2019).
 - [28] T. Li, J. Liu, Y. Xin, and Y. Zhou, Supersymmetric SYK model and random matrix theory, *Journal of High Energy Physics* **2017**, 111 (2017).

- [29] T. Kanazawa and T. Wettig, Complete random matrix classification of SYK models with $n=0, 1$ and 2 supersymmetry, *Journal of High Energy Physics* **2017**, 50 (2017).
- [30] A. M. García-García, B. Loureiro, A. Romero-Bermúdez, and M. Tezuka, Chaotic-integrable transition in the Sachdev-Ye-Kitaev model, *Phys. Rev. Lett.* **120**, 241603 (2018).
- [31] E. Iyoda, H. Katsura, and T. Sagawa, Effective dimension, level statistics, and integrability of Sachdev-Ye-Kitaev-like models, *Phys. Rev. D* **98**, 086020 (2018).
- [32] A. M. García-García, T. Nosaka, D. Rosa, and J. J. M. Verbaarschot, Quantum chaos transition in a two-site Sachdev-Ye-Kitaev model dual to an eternal traversable wormhole, *Phys. Rev. D* **100**, 026002 (2019).
- [33] F. Sun and J. Ye, Periodic table of the ordinary and supersymmetric Sachdev-Ye-Kitaev models, *Phys. Rev. Lett.* **124**, 244101 (2020).
- [34] F. Sun, Y. Yi-Xiang, J. Ye, and W.-M. Liu, Classification of the quantum chaos in colored Sachdev-Ye-Kitaev models, *Phys. Rev. D* **101**, 026009 (2020).
- [35] T. Nosaka and T. Numasawa, Quantum chaos, thermodynamics and black hole microstates in the mass deformed SYK model, *Journal of High Energy Physics* **2020**, 81 (2020).
- [36] J. Behrends and B. Béri, Symmetry classes, many-body zero modes, and supersymmetry in the complex sachdev-ye-kitaev model, *Phys. Rev. D* **101**, 066017 (2020).
- [37] J. Behrends and B. Béri, Supersymmetry in the standard sachdev-ye-kitaev model, *Phys. Rev. Lett.* **124**, 236804 (2020).
- [38] Y. Liao, A. Vikram, and V. Galitski, Many-body level statistics of single-particle quantum chaos, *Phys. Rev. Lett.* **125**, 250601 (2020).
- [39] P. H. C. Lau, C.-T. Ma, J. Murugan, and M. Tezuka, Correlated disorder in the SYK₂ model, *Journal of Physics A: Mathematical and Theoretical* **54**, 095401 (2021).
- [40] A. M. García-García, Y. Jia, D. Rosa, and J. J. M. Verbaarschot, Sparse Sachdev-Ye-Kitaev model, quantum chaos, and gravity duals, *Phys. Rev. D* **103**, 106002 (2021).
- [41] L. Sá and A. M. García-García, The Wishart-Sachdev-Ye-Kitaev model: Q-Laguerre spectral density and quantum chaos, *arXiv preprint arXiv:2104.07647* (2021).
- [42] A. M. García-García, L. Sá, and J. J. Verbaarschot, Symmetry classification and universality in non-Hermitian many-body quantum chaos by the Sachdev-Ye-Kitaev model, *arXiv preprint arXiv:2110.03444* (2021).
- [43] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, *Phys. Rev. B* **75**, 155111 (2007).
- [44] Y. Y. Atas, E. Bogomolny, O. Giraud, and G. Roux, Distribution of the ratio of consecutive level spacings in random matrix ensembles, *Physical review letters* **110**, 084101 (2013).
- [45] O. Giraud, N. Macé, E. Vernier, and F. Alet, Probing symmetries of quantum many-body systems through gap ratio statistics, *arXiv preprint arXiv:2008.11173* (2020).
- [46] N. Rosenzweig and C. E. Porter, "repulsion of energy levels" in complex atomic spectra, *Phys. Rev.* **120**, 1698 (1960).
- [47] T. Guhr and H. Weidenmuller, Correlations in anticrossing spectra and scattering theory. analytical aspects, *Chemical Physics* **146**, 21 (1990).
- [48] U. Hartmann, H. Weidenmüller, and T. Guhr, Correlations in anticrossing spectra and scattering theory: Numerical simulations, *Chemical Physics* **150**, 311 (1991).
- [49] J.-Z. Ma, Correlation hole of survival probability and level statistics, *Journal of the Physical Society of Japan* **64**, 4059 (1995).
- [50] H. Alt, H.-D. Gräf, T. Guhr, H. L. Harney, R. Hoferberber, H. Rehfeld, A. Richter, and P. Schardt, Correlation-hole method for the spectra of superconducting microwave billiards, *Phys. Rev. E* **55**, 6674 (1997).
- [51] T. Guhr, A. Müller-Groeling, and H. A. Weidenmüller, Random-matrix theories in quantum physics: common concepts, *Physics Reports* **299**, 189 (1998).
- [52] L. Reichl, *The Transition to Chaos: Conservative Classical Systems and Quantum Manifestations* (Springer, 2004).
- [53] R. Molina, J. Retamosa, L. M. noz, A. R. no, and E. Faleiro, Power spectrum of nuclear spectra with missing levels and mixed symmetries, *Physics Letters B* **644**, 25 (2007).
- [54] H. A. Weidenmüller and G. E. Mitchell, Random matrices and chaos in nuclear physics: Nuclear structure, *Rev. Mod. Phys.* **81**, 539 (2009).
- [55] F. Haake, *Quantum Signatures of Chaos* (Springer, 2010).
- [56] J. de la Cruz, S. Lerma-Hernández, and J. G. Hirsch, Quantum chaos in a system with high degree of symmetries, *Phys. Rev. E* **102**, 032208 (2020).
- [57] S. H. Tekur and M. S. Santhanam, Symmetry deduction from spectral fluctuations in complex quantum systems, *Phys. Rev. Research* **2**, 032063(R) (2020).
- [58] M. Robnik and M. V. Berry, False time-reversal violation and energy level statistics: the role of anti-unitary symmetry, *Journal of Physics A: Mathematical and General* **19**, 669 (1986).
- [59] M. V. Berry and M. Robnik, Statistics of energy levels without time-reversal symmetry: Aharonov-bohm chaotic billiards, *Journal of Physics A: Mathematical and General* **19**, 649 (1986).
- [60] F. M. Izraïlev, Simple models of quantum chaos: Spectrum and eigenfunctions, *Physics Reports* **196**, 299 (1990).
- [61] T. Seligman and J. Verbaarschot, Quantum spectra of classically chaotic systems without time reversal invariance, *Physics Letters A* **108**, 183 (1985).
- [62] M. Fremling, C. Repellin, J.-M. Stéphan, N. Moran, J. Slingerland, and M. Haque, Dynamics and level statistics of interacting fermions in the lowest landau level, *New Journal of Physics* **20**, 103036 (2018).
- [63] J. Maldacena and X.-L. Qi, Eternal traversable wormhole, *arXiv preprint arXiv:1804.00491* (2018).
- [64] S. Plugge, E. Lantagne-Hurtubise, and M. Franz, Revival dynamics in a traversable wormhole, *Phys. Rev. Lett.* **124**, 221601 (2020).
- [65] S. Sahoo, E. Lantagne-Hurtubise, S. Plugge, and M. Franz, Traversable wormhole and hawking-page transition in coupled complex syk models, *Phys. Rev.*

- Research **2**, 043049 (2020).
- [66] T. Nosaka and T. Numasawa, Chaos exponents of SYK traversable wormholes, *Journal of High Energy Physics* **2021**, 150 (2021).
- [67] R. Haenel, S. Sahoo, T. H. Hsieh, and M. Franz, Traversable wormhole in coupled Sachdev-Ye-Kitaev models with imbalanced interactions, *Phys. Rev. B* **104**, 035141 (2021).
- [68] F. Alet, M. Hanada, A. Jevicki, and C. Peng, Entanglement and confinement in coupled quantum systems, *Journal of High Energy Physics* **2021**, 34 (2021).
- [69] J. Maldacena and A. Milekhin, Syk wormhole formation in real time, *Journal of High Energy Physics* **2021**, 258 (2021).
- [70] A. M. García-García, J. P. Zheng, and V. Ziogas, Phase diagram of a two-site coupled complex syk model, *Phys. Rev. D* **103**, 106023 (2021).
- [71] P. Zhang, More on complex Sachdev-Ye-Kitaev eternal wormholes, *Journal of High Energy Physics* **2021**, 87 (2021).
- [72] I. R. Klebanov, A. Milekhin, G. Tarnopolsky, and W. Zhao, Spontaneous breaking of $U(1)$ symmetry in coupled complex SYK models, *Journal of High Energy Physics* **2020**, 162 (2020).
- [73] Y. Chen, H. Zhai, and P. Zhang, Tunable quantum chaos in the Sachdev-Ye-Kitaev model coupled to a thermal bath, *Journal of High Energy Physics* **2017**, 150 (2017).
- [74] S. Banerjee and E. Altman, Solvable model for a dynamical quantum phase transition from fast to slow scrambling, *Phys. Rev. B* **95**, 134302 (2017).
- [75] A. Haldar and V. B. Shenoy, Strange half-metals and mott insulators in sachdev-ye-kitaev models, *Phys. Rev. B* **98**, 165135 (2018).
- [76] A. Haldar, P. Haldar, S. Bera, I. Mandal, and S. Banerjee, Quench, thermalization, and residual entropy across a non-fermi liquid to fermi liquid transition, *Phys. Rev. Research* **2**, 013307 (2020).
- [77] A. Haldar, O. Tavakol, and T. Scaffidi, Variational wave functions for sachdev-ye-kitaev models, *Phys. Rev. Research* **3**, 023020 (2021).
- [78] A. Pandey and M. L. Mehta, Gaussian ensembles of random hermitian matrices intermediate between orthogonal and unitary ones, *Communications in Mathematical Physics* **87**, 449 (1983).
- [79] J. French, V. Kota, A. Pandey, and S. Tomsovic, Statistical properties of many-particle spectra v. Fluctuations and symmetries, *Annals of Physics* **181**, 198 (1988).
- [80] G. Lenz and F. Haake, Transitions between universality classes of random matrices, *Phys. Rev. Lett.* **65**, 2325 (1990).
- [81] G. Lenz and F. Haake, Reliability of small matrices for large spectra with nonuniversal fluctuations, *Phys. Rev. Lett.* **67**, 1 (1991).
- [82] P. Shukla and A. Pandey, The effect of symmetry-breaking in chaotic spectral correlations, *Nonlinearity* **10**, 979 (1997).
- [83] S.-H. Chung, A. Gokirmak, D.-H. Wu, J. S. A. Bridgewater, E. Ott, T. M. Antonsen, and S. M. Anlage, Measurement of wave chaotic eigenfunctions in the time-reversal symmetry-breaking crossover regime, *Phys. Rev. Lett.* **85**, 2482 (2000).
- [84] S. Schierenberg, F. Bruckmann, and T. Wettig, Wigner surmise for mixed symmetry classes in random matrix theory, *Phys. Rev. E* **85**, 061130 (2012).
- [85] F. Schweiner, J. Main, and G. Wunner, Goe-gue-poisson transitions in the nearest-neighbor spacing distribution of magnetoexcitons, *Phys. Rev. E* **95**, 062205 (2017).
- [86] F. Schweiner, J. Laturner, J. Main, and G. Wunner, Crossover between the gaussian orthogonal ensemble, the gaussian unitary ensemble, and poissonian statistics, *Phys. Rev. E* **96**, 052217 (2017).
- [87] A. Sarkar, M. Kothiyal, and S. Kumar, Distribution of the ratio of two consecutive level spacings in orthogonal to unitary crossover ensembles, *Phys. Rev. E* **101**, 012216 (2020).
- [88] A. L. Corps and A. Relaño, Distribution of the ratio of consecutive level spacings for different symmetries and degrees of chaos, *Phys. Rev. E* **101**, 022222 (2020).