



Anatilmnocola floriformis sp. nov., a novel member of the family *Pirellulaceae* from a boreal lake, and emended description of the genus *Anatilmnocola*

Irina S. Kulichevskaya · Anastasia A. Ivanova · Nataliya E. Suzina · Jaap S. Sinninghe Damsté · Svetlana N. Dedysh

Received: 23 July 2022 / Accepted: 7 August 2022 / Published online: 14 August 2022
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2022

Abstract Planctomycetes of the family *Pirellulaceae* are commonly addressed as budding aquatic bacteria with a complex lifestyle. Although this family is well represented by cultured and taxonomically characterized isolates, nearly all of them were obtained from brackish or marine habitats. The examples of described freshwater *Pirellulaceae* planctomycetes are limited to two species only, *Pirellula staley* and '*Anatilmnocola aggregata*'. In this study,

we characterized a novel freshwater planctomycete of the genus '*Anatilmnocola*', strain PX40^T, which was isolated from a boreal eutrophic lake. Strain PX40^T was represented by budding, unpigmented, ellipsoidal to pear-shaped cells, which often occurred in characteristic flower-like rosettes. Cells were covered by bundles of fimbriae; crateriform-like structures were localized on a reproductive cell pole only. These planctomycetes were obligately aerobic, heterotrophic bacteria that utilized various sugars and some polysaccharides, and were highly sensitive to NaCl. Growth occurred in the pH range 5.0–7.5 (with an optimum at pH 6.5–7.0), and at temperatures between 15 and 30 °C (with an optimum at 22–25 °C). The major fatty acids of strain PX40^T were C18:1 ω 9c, C16:0, and 16:1 ω 7c; cells also contained a wide variety of hydroxy- and dihydroxy-fatty acids and a C31:9 alkene. The major intact polar lipids were diacylglyceryl-(N,N,N)-trimethylhomoserines. The 16S rRNA gene sequence of strain PX40^T displayed 96.6% similarity to that of '*Anatilmnocola aggregata*' ETA_A8^T. The genome of strain PX40^T was 8.93 Mb in size and contained one copy of rRNA operon, 76 tRNA genes and 7092 potential protein-coding genes. The DNA G+C content was 57.8%. The ANI value between strain PX40^T and '*Anatilmnocola aggregata*' ETA_A8^T was 78.3%, suggesting that these planctomycetes represent distinct species. We, therefore, propose a novel species of the genus '*Anatilmnocola*', '*A. floriformis*' sp. nov., with strain

The annotated genome and 16S rRNA gene sequences of strain PX40^T have been deposited in NCBI GenBank under the accession numbers JAMLFW000000000 and OP020302, respectively.

I. S. Kulichevskaya · A. A. Ivanova · S. N. Dedysh (✉)
Winogradsky Institute of Microbiology, Research Center of Biotechnology of the Russian Academy of Sciences, Moscow 119071, Russia
e-mail: dedysh@mail.ru

N. E. Suzina
Institute of Biochemistry and Physiology of Microorganisms, Pushchino Scientific Center for Biological Research of the Russian Academy of Sciences, Moscow Region, 142290 Pushchino, Russia

J. S. Sinninghe Damsté
Department of Marine Microbiology and Biogeochemistry, NIOZ Royal Netherlands Institute for Sea Research, PO Box 59, 1790 AB Den Burg, The Netherlands

J. S. Sinninghe Damsté
Faculty of Geosciences, Department of Earth Sciences, Geochemistry, Utrecht University, Utrecht, The Netherlands

PX40^T (=KCTC 92369^T=VKM B-3621^T=UQM 41463^T) as the type strain.

Keywords Freshwater planctomycetes · Family *Pirellulaceae* · *Anatolimnocola* · Cultivation studies · Boreal lake

Introduction

The *Planctomycetota* (formerly, *Planctomycetes*) is one of the morphologically distinct, widespread, and most puzzling phyla of the domain *Bacteria* (Fuerst 1995; Jenkins and Staley 2013; Wiegand et al. 2018; Rivas-Marín and Devos 2018; Wiegand et al. 2020). Taxonomically characterized members of this phylum belong to the classes *Planctomycetia* and *Phycisphaerae*, which accommodate chemo-organotrophic bacteria that utilize a wide range of organic compounds and differ with regard to environmental adaptations (Ward and Dedysh 2022; Fukunaga 2022). The third class-level group within the *Planctomycetota*, “*Candidatus Brocadii*”, is represented by the chemo-lithoautotrophic “anammox” planctomycetes, which are capable of anaerobic ammonium oxidation (Jetten et al. 2009; Kartal et al. 2012). Members of the *Planctomycetia* divide by budding, while representatives of the *Phycisphaerae* and “*Candidatus Brocadii*” multiply by binary fission.

The largest diversity of cultured and described planctomycetes belongs to the class *Planctomycetia*, with the largest pool of the currently available isolates belonging to the order *Pirellulales*, the family *Pirellulaceae*. This family accommodates Gram-stain negative, budding, chemoheterotrophic planctomycetes with ovoid, ellipsoidal, pear- or teardrop-shaped cells, which frequently form eye-catching rosette-like clusters (Dedysh et al. 2020). Cells of *Pirellulaceae* planctomycetes do not possess a stalk but may form a fibrillar holdfast. Members of this family are aerobic and facultatively anaerobic, neutrophilic bacteria that grow on various sugars and polysaccharides. These microorganisms are most abundant in marine and freshwater habitats and, therefore, are commonly addressed as aquatic bacteria (Schlesner 1994; Fuerst 1995; Brümmer et al. 2004; Bondoso et al. 2014, 2015, 2017; Wiegand et al. 2020, 2021). However, *Pirellulaceae* planctomycetes also occur in peatlands, soils, activated sludge and various other environments

(Chouari et al. 2015; Dedysh and Ivanova 2019; Kallscheuer et al. 2020b; Rensink et al. 2020; Waqqas et al. 2020).

The type genus of this family is the genus *Pirellula* (Schlesner and Hirsch 1987). Other described genera in the family *Pirellulaceae* with validly published names are *Aureliella* (Rast et al. 2020), *Blastopirellula* (Schlesner et al. 2004; Kallscheuer et al. 2020b), *Bremerella* (Rensink et al. 2020), *Lignipirellula* (Peeters et al. 2020a), *Crateriforma* (Peeters et al. 2020b), *Mariniblastus* (Lage et al. 2017), *Novipirellula* (Kallscheuer et al. 2020c), *Rhodopirellula* (Schlesner et al. 2004; Kallscheuer et al. 2020d), *Roseimaritima* (Bondoso et al. 2015), *Rosistilla* (Waqqas et al. 2020), *Rubripirellula* (Bondoso et al. 2015) and *Stieleria* (Kallscheuer et al. 2020a). Two additional described genera in this family with pending validation status are ‘*Anatolimnocola*’ (Kallscheuer et al. 2021) and ‘*Roseiconus*’ (Kumar et al. 2021). The isolates representing most of the above listed genera have been obtained from marine and brackish habitats such as natural or artificial marine surfaces, macroalgae, shallow-sea hydrothermal vents and others. The only currently described freshwater *Pirellulaceae* planctomycetes belong to the genera *Pirellula* and ‘*Anatolimnocola*’. Each of these genera is represented by a single species. *Pirellula staley*, the first cultured and described organism in the family, was isolated from the Lake Lansing, Michigan, USA (Staley 1973; Schlesner and Hirsch 1984, 1987). ‘*Anatolimnocola aggregata*’ ETA_A8^T was isolated from a municipal duck pond in Wolfenbüttel, Germany, during a phytoplankton bloom (Kallscheuer et al. 2021). Notably, these planctomycetes form a common phylogenetic lineage in the 16S rRNA gene-based tree of *Pirellulaceae* planctomycetes and display significant phenotypic similarity being represented by mesophilic, neutrophilic, obligately aerobic heterotrophic bacteria. The major morphological differences noticed between the *Pirellula* and strain ETA_A8^T was the absence of a holdfast and crateriform structures in the latter bacterium, which was also slightly larger in size (Kallscheuer et al. 2021). Neither the range of growth substrates, nor the chemotaxonomic characteristics were reported for ‘*Anatolimnocola aggregata*’ ETA_A8^T. In this study, we isolated one additional representative of the genus ‘*Anatolimnocola*’ from a freshwater ecosystem, i.e., a boreal eutrophic lake in Northern Russia, determined

its genome sequence and described morphological, phenotypic and chemotaxonomic characteristics. Based on these results, we emend description of the genus '*Anatilimnocola*' and propose a novel species within this genus, *A. floriformis* sp. nov.

Materials and methods

Sampling site

Strain PX40^T was isolated from water collected from the upper oxic layer (0–10 cm) of the boreal eutrophic lake Morotskoye (Vologda region, European North Russia, 58°43'28.5"N, 37°39'07"E) in August 2017. Its surface area is 6.24 km² and its maximum depth is 2.1 m. Site-specific parameters are as follows (range given in parenthesis): water conductivity (30–50 μS cm⁻¹), dissolved organic carbon (25.0–37.5 mg L⁻¹), total nitrogen (2.0 and 3.7 mg L⁻¹), and total phosphorus (45–77 μg L⁻¹). The pH is 7.5–7.8.

Isolation and cultivation

The enrichment culture was obtained using an agar medium prepared with original lake water and containing 0.05 g L⁻¹ carbenicillin (sodium salt). An aliquot of lake water (10 ml) was spread onto this medium, the plates were placed in bags to prevent drying, and incubated at room temperature as recommended by Hirsch et al. (1977). Aliquots (20 μl) of the microbial cell masses that developed on these plates after 4 weeks of incubation were spread plated onto the MPYVG medium (modification of medium 621 DSMZ), solidified with 10 g phytigel (Sigma–Aldrich) and containing (per liter distilled water): 0.1 g peptone (Fluka), 0.25 g yeast extract, 0.1 g NH₄NO₃, 20 ml Hutner's basal salts (Staley et al. 1992). After sterilization, the medium was supplemented with 5 ml L⁻¹ 5% glucose solution, 1 ml L⁻¹ Staley's vitamin solution (Staley et al. 1992), 0.05 g carbenicillin (sodium salt) L⁻¹, and maintained at pH 7.5. The plates were subsequently incubated at 22 °C for 4 weeks. Colonies that developed on plates were screened microscopically for the presence of budding cells with a planctomycete-like morphology. The selected cell material was re-streaked onto the same medium MPYVG, supplemented with 0.05%

(w/v) glucose. This procedure resulted in isolation of two strains, PX69^T and PX40^T. The former strain was described in an earlier publication (Dedysh et al. 2020), while the characterization of the later isolate is described here. Once obtained in pure culture, strain PX40^T was maintained on MPYVC medium and was sub-cultured at 2 month intervals.

Microscopic studies

Morphological observations and cell size measurements were made with a Zeiss Axioplan 2 microscope and Axiovision 4.2 software (Zeiss, Germany). For the electron microscopy of negatively stained preparations, cells suspension was placed onto grids coated with formvar film, dried and treated with 0.3% aqueous solution of uranyl acetate (pH 4.0). The specimen samples were examined with JEM-1400 transmission electron microscope (JEOL, Japan) at the UNIQEM Collection Core Facility, Research Center of Biotechnology of the Russian Academy of Sciences, at an accelerating voltage of 80 kV.

Physiological tests

Physiological tests were performed in liquid MPYVG medium. Growth of strain PX40^T was monitored by nephelometry at 600 nm in an Eppendorf BioPhotometer for 2–3 weeks under a variety of conditions, including temperatures of 4–37 °C, pH 3.8–8.0 and NaCl concentrations of 0–3.0% (w/v). Incubations at various temperatures were made under static conditions in triplicate; OD₆₀₀ was determined after 2 weeks of incubation. Variations in the pH were achieved by using MES (pH 4.0–6.5) and MOPS (pH 6.5–7.9) buffer systems. Carbon source utilization was determined using mineral medium M1 (Kulichvskaya et al. 2017), supplemented with 0.005% yeast extract and the individual carbon sources given in the species description in a concentration 0.05% (w/v). Cultivation was done in 120 ml flasks containing 20 ml medium. Cultures were incubated at 22 °C for 2–3 weeks on a shaker. All experiments were performed in triplicate.

The ability to grow under anoxic conditions was tested in tightly closed 120 ml serum flasks containing 80 ml of liquid medium MPYVG. Before autoclaving, these flasks were flushed for 5 min with a

mixture of CO₂ and N₂ (7: 93). Growth was assessed by measuring OD₆₀₀ after 3 weeks of incubation.

Oxidative and fermentative utilization of carbohydrates was determined as described for the Hugh-Leifson test (Gerhardt 1981). Analyzes of enzymatic profiles, oxidase test, gelatin and urease hydrolysis were made with API ZYM and API 20NE kits (bio-Mérieux). Catalase test was carried out by standard method (Gerhardt 1981).

Lipid analyses

For lipid analysis, cells of strain PX40^T were grown in liquid MPVYG medium and harvested in the late exponential growth phase. Fatty acids were analyzed after acid hydrolysis of whole cells following the procedure described elsewhere (Sinninghe Damsté et al. 2011). The main intact polar lipids (IPLs) in strain PX40^T were analyzed following procedures reported previously (Moore et al. 2013).

Genome sequencing, annotation and analysis

Genomic DNA was isolated from strain PX40^T using the phenol–chloroform assay with addition of hexadecyltrimethylammonium bromide (Wilson 2001). The sequencing library for Illumina sequencing was prepared using the NEBNext Ultra II DNA Library Prep Kit (New England Biolabs, USA) following the manufacturer's instructions. The sequencing of this library was performed on the Illumina HiSeq2500 platform using HiSeq Rapid Run v2 sequencing reagents. Primer sequences were removed from the Illumina reads using Cutadapt v.1.17 (Martin 2011) with the default settings, and low quality read ends were trimmed using Sickle v.1.33 (option q=30) (<https://github.com/najoshi/sickle>). For Nanopore sequencing the library was prepared using the 1D ligation sequencing kit (SQK-LSK108, Oxford Nanopore, UK). Sequencing of this library was carried out in an R9.4 flow cell (FLO-MIN106) using MinION device. Hybrid assembly of Illumina and Nanopore reads was performed using Unicycler v. 0.4.8 (Wick et al. 2017).

Gene search and annotation were performed using MicrobeAnnotator (Ruiz-Perez et al. 2021) and GhostKoala (Kanehisa et al. 2016) packages. Search for the secondary metabolites was carried out with AntiSmash (Blin et al. 2019). The overall genome similarity (GGDC) between strain PX40^T and closely

related planctomycetes was determined using an online distance calculator available at DSMZ webpage. The average nucleotide identity (ANI) values were determined using an online calculator available on the server <http://enve-omics.ce.gatech.edu/ani/>.

The annotated genome and 16S rRNA gene sequences of strain PX40^T have been deposited in NCBI GenBank under the accession numbers JAM-LFW000000000 and OP020302, respectively.

Phylogenetic analysis

The 16S rRNA gene-based phylogenetic analysis was carried out using the MEGAX (Kumar et al. 2018). The phylogenetic tree was built applying the maximum-likelihood statistical method. Visualization of the tree was performed on iTOL platform (Letunic and Bork 2019).

In addition, the GTDB-Tk v. 2.0.0 toolkit (Parks et al. 2018) was used to identify 120 single-copy, phylogenetically informative bacterial marker genes used in the Genome Taxonomy Database (GTDB) classification system in the genome of strain PX40^T. These were used to construct a multiple alignment of concatenated single-copy gene sequences, comprising those from PX40^T and all related species from the GTDB. A selected part of the multiple alignment built in GTDB-Tk was used to construct a phylogenetic tree in MEGAX (Kumar et al. 2018) with default parameters. The significance levels of interior branch points obtained in maximum-likelihood analysis were determined by bootstrap analysis (100 data resamplings).

Results and discussion

Cell morphology and physiology

On phytigel-solidified medium MPVYG, strain PX40^T formed small (2–3 mm), circular, unpigmented colonies with an entire edge and a smooth surface. These colonies were composed of bacteria with ellipsoidal to pear-shaped cells, 0.7 ± 0.2 µm wide and 1.9 ± 0.5 µm long, which multiplied by budding and occurred singly or in flower-shaped rosettes (Fig. 1a, c). Daughter cells were highly motile by means of one polar flagellum (Fig. 1b). Examination of negatively stained cells of strain PX40^T using




Fig. 1 **a** Phase-contrast image of cells of strain PX40^T grown for 7 days on solid medium MPVGY; bar, 10 μ m. Insert: a group of budding cells of strain PX40^T; bar, 5 μ m. **b** Electron micrograph of a negatively stained daughter cell with flagellum; bar, 0.1 μ m. **c** Electron micrograph of negatively stained cells displaying crateriform pits (Cr), buds (B) and bundles of fimbriae (F); bar, 1 μ m

electron microscopy showed the presence of bundles of fimbriae and crateriform-like structures, which were distributed over a reproductive cell pole only (Fig. 1c).

Strain PX40^T was capable of growth at pH values between 5.0 and 7.5 (with an optimum at pH 6.5–7.0), and at temperatures between 15 and 30 °C (with an optimum at 22–25 °C). The specific growth rate of this planctomycete under the optimal growth conditions was 0.035 h⁻¹ (equal to a doubling time of 20.4 h). These values are nearly identical to those reported for ‘*Anatolimnocola aggregata*’ ETA_A8^T (Kallscheuer et al. 2021). NaCl inhibited growth at concentrations >0.3% (w/v). Strain PX40^T grew on various sugars, including N-acetylglucosamine, and some polysaccharides, such as aesculin,

arabinogalactan, dextrin, laminarin, locust bean gum, starch, xanthan gum, xylan and gelatin, but did not assimilate amino acids or organic acids (see Table 2 and the species description). Pullulan, chondroitin sulfate, cellulose, carboxymethyl cellulose, casein, chitin and chitosan were not hydrolyzed. Anaerobic growth was not observed.

Strain PX40^T was found to be catalase- and cytochrome oxidase-positive but urease-negative. Nitrate was reduced to nitrite. The test for the activity of arginine dihydrolase was positive (API 20NE strip). In the API ZYM strip, alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase, acid phosphatase, α -glucosidase and α -galactosidase were detected. The following enzyme activities were not detected: lipase, trypsin, chymotrypsin, phosphohydrolase, N-acetyl- β -glucosaminidase, β -galactosidase, cystine arylamidase, β -glucuronidase, α -fucosidase and α -mannosidase.

Lipid composition

The major fatty acids of strain PX40^T were C18:1 ω 9c, C16:0, and 16:1 ω 7c (Table 1). Cells also contained a wide variety of hydroxy- and dihydroxy-fatty acids and a C31:9 polyunsaturated *n*-alkene, which is often present in planctomycetes (Kulichevskaya et al. 2016; Dedysh et al. 2020). The major intact polar lipids were diacylglyceryl-(N,N,N)-trimethylhomoserine (DGTS) lipids.

Genome characteristics

The genome of strain PX40^T was sequenced using a combination of Illumina and Nanopore technique. Combined assembly yielded 17 contigs with a total length of 8,932,541 bp. The DNA G+C content was 57.8%. One copy of rRNA operon and 76 tRNA genes were identified. Annotation of the genome sequence revealed 7,092 potential protein-coding genes of which 3,037 could be functionally assigned. The latter value is comparable to that reported for other representatives of *Planctomycetia* (Wiegand et al. 2020).

The examination of the strain PX40^T genome revealed a set of genes encoding a bacterial flagellar machinery and chemotaxis functions, consistently with observed morphological characteristics. The genes encoding metabolic pathways common

Table 1 Relative abundance (%; normalized on their sum) of fatty acids and other lipids including an alkene present in the acid hydrolysate of cell material of strain PX40^T

Lipid type	Relative abundance (%)
Fatty acids	
isoC16:0	2.0
C16:1ω7	5.7
C16:0	35.0
C17:1 ω 8	1.0
C18:1ω9	34.9
C18:1 ω 7	2.3
C18:0	1.4
C20:1 ω 9	4.4
C20:1 ω 7	2.7
Hydroxy fatty acids	
(ω -1)OH-C24:0	1.1
(ω -1)OH-C26:0	0.6
ω OH-C32:1	0.6
ω OH-C32:0	0.8
β ,x-diOH-C32:0	1.6
Other lipids	
C31:0-keto-ol	1.0
C33:1-keto-ol	0.8
C31:9 <i>n</i> -alkene	4.1
IPLs	
DGTS	+++
Lyso-DGTS	+

Only components with a relative abundance $\geq 0.5\%$ of the total are listed. Major components ($>5\%$) are given in bold type face

for chemoorganotrophic bacteria, such as glycolysis, the citrate cycle, the pentose-phosphate pathway and oxidative phosphorylation were present in the genome of strain PX40^T. Also, 30 genes encoding proteins with significant similarity to sulfatases were found in the genome. High number of sulfatase-encoding genes is typical for the genomes of *Pirellulaceae* planctomycetes (Wegner et al. 2013). Thus, the genome of *Pirellula staley* ATCC 27377^T contains 34 sulfatase-encoding genes, while the number of these genes in *Rhodopirellula* genomes exceeds one hundred. This large diversity of sulfatases in the genomes of *Pirellulaceae* planctomycetes was interpreted as a response to the diversity

of sulfated compounds in nature and especially in the marine environment (Wegner et al. 2013).

Genome mining of gene clusters that encode biosynthetic pathways for secondary metabolites in strain PX40^T revealed 6 clusters. Four clusters encode terpenes, one cluster encodes polyketide synthases (PKS) of type III and the last one codes for a non-ribosomal peptide synthetase (NRPS). The closest gene homologs to all terpene clusters from strain PX40^T belong to the planctomycete '*Anatolimnocola aggregata*' ETA_A8^T. The closest homologs for PKS type III cluster and NRPS belong to *Pirellula*-like planctomycete strain SH-Sr6A and *Mycobacterium shinjukuense* JCM 14233.

Phylogenetic placement of strain PX40^T

The 16S rRNA gene sequence of strain PX40^T was affiliated with the family *Pirellulaceae* and grouped with several environmental clone sequences obtained from a freshwater lake (GenBank accession number DQ787725), an alpine wetland (FJ801184), wastewater treatment systems (LR634621, LR642682, LR645593), and a rhizosphere soil (FJ444676) (Fig. 2). Among the taxonomically characterized planctomycetes, the highest 16S rRNA gene similarity (96.6%) was observed to that of '*Anatolimnocola aggregata*' ETA_A8^T.

To determine the genome-based phylogenetic position of strain PX40^T within the *Planctomycetes*, a phylogenetic tree based on concatenated sequences of conservative marker genes was constructed. This placed strain PX40^T within the family *Pirellulaceae* (Fig. 3). Following the proposed minimal standards for the use of genome data for the taxonomy of prokaryotes (Chun et al. 2018), we calculated both the overall genome similarity and the average nucleotide identity (ANI) between PX40^T and closely related planctomycetes. The ANI value between strain PX40^T and its closest relative with determined genome sequence, '*Anatolimnocola aggregata*' ETA_A8^T, was 78.3%. According to ANI thresholds recently proposed for taxonomic delineation (i.e. 65–95% for the same genus and 95–100% for the same species Konstantinidis et al. 2017), these two bacteria represented different species in one genus. The ANI value between strain PX40^T and *Pirellula staley* ATCC27377^T was 75.6%. The overall genome similarity between PX40^T and '*A. aggregata*' ETA_A8^T





Fig. 2 16S rRNA gene-based maximum-likelihood tree showing the phylogenetic relationship of strain PX40^T to other described *Pirellulaceae* planctomycetes and some environmental 16S rRNA gene clone sequences displaying > 95% similarity to those of strain PX40^T and *Pirellula staley* DSM 6068.

Habitats are coded by colored triangles. The clade defined by the genera *Pirellula* and ‘*Anatilimnocola*’ is depicted with a greenish background. Bootstrap values are shown as circles (70–100%) with the respective legend

Fig. 3 Genome-based phylogeny of the family *Pirellulaceae* defined in the GTDB taxonomy. The tree was reconstructed using the GTDB-Tk v.2.0.0 (Parks et al. 2018). The significance levels of interior branch points obtained in maximum-likelihood analysis were determined by bootstrap analysis (100 data resamplings). The root (not shown) was composed of 22 genomes of members of the family ‘*Scalinduaceae*’. Bar, 0.1 substitutions per amino acid position



was $20.8 \pm 2.5\%$, while the genome similarity between PX40^T and *Pirellula staley* ATCC 27377^T was $19.3 \pm 2.3\%$.

Based on the phylogenetic divergence between strain PX40^T and ‘*A. aggregata*’ ETA_A8^T and

several other characteristics that differentiate these planctomycetes (Table 2), we propose placing strain PX40^T in a novel species of the genus ‘*Anatilimnocola*’, *A. floriformis* sp. nov. We also include additional characteristics to the earlier published description of this genus.

Table 2 Major characteristics that distinguish strain PX40^T and ‘*Anatilimnocola aggregata*’ ETA_A8^T (Kallscheuer et al. 2021)

Characteristic	Strain PX40 ^T	<i>Anatilimnocola aggregata</i> ETA_A8 ^T
Phenotypic features		
Habitat	Eutrophic boreal lake	Municipal duck pond
Cell size (μm)	0.7 ± 0.2 × 1.9 ± 0.5	2.0 ± 0.3 × 1.4 ± 0.2
Cell shape	Ellipsoidal to pear shaped	Round grain rice shaped
Type of rosettes	Flower-shaped	Shapeless aggregates
Temperature range (optimum) (°C)	15–30 (22–25)	15–33 (30)
pH growth range (optimum)	5.0–7.5 (6.5–7.0)	5.0–10.0 (8.0)
Flagella	Yes	n.o
Crateriform structures	Yes	n.o
Major fatty acids	C18:1ω9c, C16:0, C16:1ω7c	n.d
Genomic features		
Genome size (Mb)	8.93	9.01
DNA G+C content, mol%	57.8	57.8

n.d. not determined, n.o. not observed

Emended description of the genus *Anatilimnocola* Kallscheuer et al. (2021)

The description of the genus '*Anatilimnocola*' is as given by Kallscheuer et al. (2021) with the following modifications. Crateriform-like structures are distributed on a reproductive cell pole; holdfasts are present. Carbon substrates are sugars and some polysaccharides. Sensitive to NaCl. The major fatty acids are C18:1 ω 9c, C16:0, and 16:1 ω 7c. The major polar lipid are diacylglyceryl-(N,N,N)-trimethylhomoserines. Main habitats are eutrophic freshwater bodies.

Description of *Anatilimnocola floriformis* sp. nov.

Anatilimnocola floriformis (flo.ri.for'mis. L. masc. n. *flos*, a flower; L. masc./fem. stuff. *-formis*, in the shape of (from L. fem. n. *forma*, shape); N.L. fem. adj. *floriformis*, referring to the characteristic of the cells to form flower-shaped aggregates).

Exhibit the following properties in addition to those given in the genus description. Colonies are small (2–3 mm in diameter), circular and unpigmented. The cells are 0.7 ± 0.2 μm wide and 1.9 ± 0.5 μm long, occur as single cells or flower-shaped rosettes and reproduce by polar budding. The temperature range for growth is 15–30 °C with the optimum at 22–25 °C. The pH range for growth is 5.0–7.5 with 6.5–7.0 as the optimum. Growth does not occur at NaCl concentrations > 0.3% (w/v). Chemoheterotrophic, aerobic, catalase- and cytochrome oxidase-positive but urease-negative. N-acetylglucosamine, D-galactose, L-rhamnose, D-arabinose, D-glucose, xylose, sucrose, maltose, lactose, D-cellobiose, D-trehalose, lactulose are assimilated. Capable of hydrolyzing aesculin, arabinogalactan, dextrin, laminarin, locust bean gum, gelatin, starch, xanthan gum and xylan. Amino acids, casamino acids, fructose, L-fucose, D-ribose, L-sorbose, D-raffinose, D-mannitol, sorbitol, glycerol, succinate, malate, citrate, benzoate, fumarate, formate, acetate, pyruvate and inulin are not assimilated. Cannot hydrolyze casein, chitosan, chitin, carboxymethyl cellulose and cellulose. Nitrate is reduced to nitrite. Arginine dihydrolase is positive (API 20NE strip). In the API ZYM strip, alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase, acid phosphatase, α -glucosidase and α -galactosidase were detected. The following enzyme activities were not

detected: lipase, trypsin, chymotrypsin, phosphohydrolase, N-acetyl- β -glucosaminidase, β -galactosidase, cystine arylamidase, β -glucuronidase, α -fucosidase and α -mannosidase. The major fatty acids are C18:1 ω 9c, C16:0, and 16:1 ω 7c. The major polar lipids are diacylglyceryl-(N,N,N)-trimethylhomoserines. The DNA G+C of the type strain is 57.8 mol%. The type strain is strain PX40^T (= KCTC 92369^T = VKM B-3621^T = UQM 41463^T), which was isolated from the boreal lake Morotskoye (Vologda region, European North Russia).

Acknowledgements The authors thank W. Irene C. Rijpstra (NIOZ Royal Netherlands Institute for Sea Research) for the help with lipid analyses.

Author contributions I.S.K. obtained the isolate, described its physiology and prepared figure 1; A.A.I. determined and analysed the genome sequence and prepared figures 2 and 3; N.E.S. obtained electron microscopy images of cells; J.S.S.D. performed lipid analyses; S.N.D. wrote the main manuscript text. All authors reviewed the manuscript.

Funding This study was supported by the Ministry of Science and Higher Education of the Russian Federation (Research topic No 122041100029-2 and the Grant agreement № 075-15-2021-1051). J.S.S.D. was supported by the SIAM Gravitation Grant (024.002.002) from the Dutch Ministry of Education, Culture and Science (OCW).

Declarations

Conflict of interest The authors declare that there are no conflicts of interest.

References

- Blin K, Shaw S, Steinke K et al (2019) antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Res* 47:W81–W87
- Bondoso J, Balague V, Gasol JM, Lage OM (2014) Community composition of the *Planctomycetes* associated with different macroalgae. *FEMS Microbiol Ecol* 88:445–456
- Bondoso J, Albuquerque L, Nobre MF, Lobo-da-Cunha A, da Costa MS, Lage OM (2015) *Roseimaritima ulvae* gen. nov., sp. nov. and *Rubripirellula obstinata* gen. nov., sp. nov. two novel planctomycetes isolated from the epiphytic community of macroalgae. *Syst Appl Microbiol* 38:8–15
- Bondoso J, Godoy-Vitorino F, Balague V, Gasol JM, Harder J, Lage OM (2017) Epiphytic *Planctomycetes* communities associated with three main groups of macroalgae. *FEMS Microbiol Ecol* 93:fiw255
- Brümmer IHM, Felske ADM, Wagner-Döbler I (2004) Diversity and seasonal changes of uncultured *Planctomycetales* in river biofilms. *Appl Environ Microbiol* 70(9):5094–5101

- Chouari R, Le Paslier D, Daegelen P, Ginestet P, Weissenbach J, Sghir A (2015) Molecular evidence for novel planctomycete diversity in a municipal wastewater treatment plant. *CEUR Workshop Proc* 1542(12):33–36
- Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, da Costa MS, Rooney AP, Yi H, Xu XW, De Meyer S, Trujillo ME (2018) Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 68:461–466
- Dedysh SN, Ivanova AA (2019) Planctomycetes in boreal and subarctic wetlands: diversity patterns and potential ecological functions. *FEMS Microbiol Ecol*. <https://doi.org/10.1093/femsec/fiy227>
- Dedysh SN, Kulichevskaya IS, Beletsky AV, Ivanova AA, Rijpstra WIC, Damste JSS, Mardanov AV, Ravin NV (2020) *Lacipirellula parvula* gen. nov., sp. nov., representing a lineage of planctomycetes widespread in low oxygen habitats, description of the family *Lacipirellulaceae* fam. nov. and proposal of the orders *Pirellulales* ord. nov., *Gemmatales* ord. nov. and *Isosphaerales* ord. nov. *Syst Appl Microbiol* 43:126050
- Gerhardt P (1981) Manual of methods for general bacteriology. American Society for Microbiology
- Fuerst JA (1995) The *Planctomycetes*: emerging models for microbial ecology, evolution and cell biology. *Microbiology* 141:1493–1506
- Fukunaga Y (2022) *Phycisphaerae*. In: Trujillo ME, Dedysh S, DeVos P, Hedlund B, Kämpfer P, Rainey FA, Whitman WB (eds) *Bergey's manual of systematics of archaea and bacteria*. <https://doi.org/10.1002/9781118960608.cbm00069>
- Hirsch P, Müller M, Schlesner H (1977) New aquatic budding and prosthecate bacteria and their taxonomic position. *Soc Appl Bacteriol Symp Ser* 6:107–133
- Jenkins C, Staley JT (2013) History, classification and cultivation of the *Planctomycetes*. In: Fuerst JA (ed) *Planctomycetes: cell structure, origins and biology*. Humana Press, Totowa, pp 1–38
- Jetten MSM, Niftrik LV, Strous M, Kartal B, Keltjens JT, Op den Camp HJ (2009) Biochemistry and molecular biology of anammox bacteria. *Crit Rev Biochem Mol Biol* 44:65–84
- Kartal B, van Niftrik L, Keltjens JT, Jetten MSM, Op den Camp HJ (2012) Anammox-growth physiology, cell biology, and metabolism. *Adv Microb Physiol* 60:211–262
- Kallscheuer N, Jeske O, Sandargo B, Boedeker C, Wiegand S, Bartling P, Jogler M, Rohde M, Petersen J, Medema MH, Surup F, Jogler C (2020a) The planctomycete *Stierleria maiorica* Mal15^T employs stieleriaceins to alter the species composition in marine biofilms. *Commun Biol* 3(1):303. <https://doi.org/10.1038/s42003-020-0993-2>
- Kallscheuer N, Wiegand S, Heuer A, Rensink S, Boersma AS, Jogler M, Boedeker C, Peeters SH, Rast P, Jetten MSM, Rohde M, Jogler C (2020b) *Blastopirellula retiformator* sp. nov. isolated from the shallow-sea hydrothermal vent system close to Panarea Island. *Antonie van Leeuwenhoek* 113:1811–1822. <https://doi.org/10.1007/s10482-019-01377-2>
- Kallscheuer N, Wiegand S, Peeters SH, Jogler M, Boedeker C, Heuer A, Rast P, Jetten MSM, Rohde M, Jogler C (2020c) Description of three bacterial strains belonging to the new genus *Novipirellula* gen. nov., reclassification of *Rhodopirellula rosea* and *Rhodopirellula caenicola* and readjustment of the genus threshold of the phylogenetic marker *rpoB* for *Planctomycetaceae*. *Antonie van Leeuwenhoek* 113:1779–1795
- Kallscheuer N, Wiegand S, Jogler M, Boedeker C, Peeters SH, Rast P, Heuer A, Jetten MSM, Rohde M, Jogler C (2020d) *Rhodopirellula heiligendammensis* sp. nov., *Rhodopirellula pilleata* sp. nov., and *Rhodopirellula solitaria* sp. nov. isolated from natural or artificial marine surfaces in Northern Germany and California, USA, and emended description of the genus *Rhodopirellula*. *Antonie van Leeuwenhoek* 113(12):1737–1750. <https://doi.org/10.1007/s10482-019-01366-5>
- Kallscheuer N, Rast P, Jogler M, Kohn T, Boedeker C, Jeske O, Heuer A, Quast C, Glockner FO, Rohde M, Jogler C (2021) Analysis of bacterial communities in a municipal duck pond during a phytoplankton bloom and isolation of *Anatolimnocola aggregata* gen. nov., sp. nov., *Lacipirellula limnantheis* sp. nov. and *Urbifossiella limnaea* gen. nov., sp. nov. belonging to the phylum *Planctomycetes*. *Environ Microbiol* 23:1379–1396
- Kanehisa M, Sato Y, Morishima K (2016) BlastKOALA and GhostKOALA: KEGG tools for functional characterization of genome and metagenome sequences. *J Mol Biol* 428:726–731
- Konstantinidis KT, Rosselló-Móra R, Amann R (2017) Uncultivated microbes in need of their own taxonomy. *ISME J* 11:2399–2406
- Kulichevskaya IS, Ivanova A, Suzina NE, Rijpstra WIC, Sinnighe Damsté JS, Dedysh SN (2016) *Paludisphaera borealis* gen. nov., sp. nov., a hydrolytic planctomycete from northern wetlands, and proposal of *Isosphaeraceae* fam. nov. *Int J Syst Evol Microbiol* 66(2):837–844
- Kulichevskaya IS, Ivanova AA, Detkova EN, Rijpstra WIC, Sinnighe Damsté JS, Dedysh SN (2017) *Tundrisphaera lichenicola* gen. nov., sp. nov., a psychrotolerant representative of the family *Isosphaeraceae* from lichen-dominated tundra soils. *Int J Syst Evol Microbiol* 67(9):3583–3589
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 35:1547–1549
- Kumar D, Kumar G, Uppada J, Ahmed S, Sasikala C, Ramana CV (2021) Descriptions of *Roseiconus nitratreducens* gen. nov. sp. nov. and *Roseiconus lacunae* sp. nov. *Arch Microbiol* 203:741–754
- Lage OM, Albuquerque L, Lobo-da Cunha A, da Costa MS (2017) *Mariniblastus fucicola* gen. nov., sp. nov. a novel planctomycete associated with macroalgae. *Int J Syst Evol Microbiol* 67:1571–1576
- Letunic I, Bork P (2019) Interactive tree of life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res* 47:W256–W259. <https://doi.org/10.1093/nar/gkz239>

- Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17(1):10. <https://doi.org/10.14806/ej.17.1.200>
- Moore EK, Hopmans EC, Rijpstra WIC, Villanueva L, Dedysh SN et al (2013) Novel mono-, di-, and trimethylornithine membrane lipids in northern wetland planctomycetes. *Appl Environ Microbiol* 79(22):6874–6884
- Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA, Hugenholtz P (2018) A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nat Biotechnol* 36:996–1004
- Peeters SH, Wiegand S, Kallscheuer N, Jogler M, Heuer A, Jetten MSM, Boedeker C, Rohde M, and Jogler C (2020a) *Lignipirellula cremea* gen. nov., sp. nov., a planctomycete isolated from wood particles in a brackish river estuary. *Antonie van Leeuwenhoek* 113(12):1863–1875
- Peeters SH, Wiegand S, Kallscheuer N, Jogler M, Heuer A, Jetten MSM, Rast P, Boedeker C, Rohde M, Jogler C (2020b) Three marine strains constitute the novel genus and species *Crateriforma conspicua* in the phylum *Planctomycetes*. *Antonie van Leeuwenhoek* 10:1–13
- Rast P, Heuer A, Jetten MSM, Rohde M, Jogler C (2020) *Aureliella helgolandensis* gen. nov., sp. nov., a novel Planctomycete isolated from a jellyfish at the shore of the island Helgoland. *Antonie Van Leeuwenhoek* 113:1839–1849
- Rensink S, Wiegand S, Kallscheuer N, Rast P, Peeters SH, Heuer A, Boedeker C, Jetten MSM, Rohde M, Jogler M, Jogler C (2020) Description of the novel planctomycetal genus *Bremerella*, containing *Bremerella volcania* sp. nov., isolated from an active volcanic site, and reclassification of *Blastopirellula cremea* as *Bremerella cremea* comb. nov. *Antonie van Leeuwenhoek* 113:1823–1837
- Rivas-Marín E, Devos DP (2018) The paradigms they are a-changin’: past, present and future of PVC bacteria research. *Antonie van Leeuwenhoek* 111:785–799
- Ruiz-Perez CA, Conrad RE, Konstantinidis KT (2021) MicrobeAnnotator: a user-friendly, comprehensive functional annotation pipeline for microbial genomes. *BMC Bioinformatics* 22:11. <https://doi.org/10.1186/s12859-020-03940-5>
- Schlesner H (1994) The development of media suitable for the microorganisms morphologically resembling *Planctomyces* spp., *Pirellula* spp., and other *Planctomycetales* from various aquatic habitats using dilute media. *Syst Appl Microbiol* 17:135–145
- Schlesner H, Rensmann C, Tindall BJ, Gade D, Rabus R, Pfeiffer S, Hirsch P (2004) Taxonomic heterogeneity within the *Planctomycetales* as derived by DNA-DNA hybridization, description of *Rhodopirellula baltica* gen. nov., sp. nov., transfer of *Pirellula marina* to the genus *Blastopirellula* gen. nov. as *Blastopirellula marina* comb. nov. and emended description of the genus *Pirellula*. *Int J Syst Bacteriol* 54:1567–1580
- Schlesner H, Hirsch P (1984) Assignment of ATCC 27377 to *Pirella* gen. nov. as *Pirella staleyi* comb. nov. *Int J Syst Bacteriol* 34:492–495
- Schlesner H, Hirsch P (1987) Rejection of the genus name *Pirella* for pear-shaped budding bacteria and proposal to create the genus *Pirellula* gen. nov. *Int J Syst Bacteriol* 37:441. <https://doi.org/10.1099/00207713-37-4-441>
- Sinninghe Damsté JS, Rijpstra WIC, Hopmans EC, Weijers JWH, Foesel BU, Overmann J, Dedysh SN (2011) 13,16-Dimethyl octacosanedioic acid (iso-diabolic acid), a common membrane-spanning lipid of *Acidobacteria* subdivisions 1 and 3. *Appl Environ Microbiol* 77(12):4147–4154
- Staley JT (1973) Budding bacteria of the *Pasteuria* – *Blastobacter* group. *Can J Microbiol* 19(5):609–614
- Staley JT, Fuerst JA, Giovannoni S, Schlesner H (1992) The order *Planctomycetales* and the genera *Planctomyces*, *Pirellula*, *Gemmata*, and *Isosphaera*. In: Balows A, Trüper HG, Dworkin M, Harder W, Schleifer K-H (eds) *The prokaryotes: a handbook on the biology of bacteria: ecophysiology, isolation, identification, applications*. Springer, New York, pp 3710–3731
- Waqqas M, Salbreiter M, Kallscheuer N, Jogler M, Wiegand S, Heuer A, Rast P, Peeters SH, Boedeker C, Jetten MSM, Rohde M, Jogler C (2020) *Rosistilla oblonga* gen. nov., sp. nov. and *Rosistilla carotiformis* sp. nov., isolated from biotic or abiotic surfaces in Northern Germany, Mallorca, Spain and California, USA. *Antonie van Leeuwenhoek* 113:1939–1952
- Ward NL, Dedysh SN (2022) *Planctomycetia*. In: Trujillo ME, Dedysh S, DeVos P, Hedlund B, Kämpfer P, Rainey FA, Whitman WB (eds) *Bergey’s manual of systematics of archaea and bacteria*. <https://doi.org/10.1002/9781118960608.cbm00040.pub2>
- Wegner CE, Richter-Heitmann T, Klindworth A, Klockow C, Richter M, Achstetter T, Glöckner FO, Harder J (2013) Expression of sulfatases in *Rhodopirellula baltica* and the diversity of sulfatases in the genus *Rhodopirellula*. *Mar Genom* 9:51–61. <https://doi.org/10.1016/j.margen.2012.12.001>
- Wick RR, Judd LM, Gorrie CL, Holt KE (2017) Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>
- Wiegand S, Jogler M, Jogler C (2018) On the maverick *Planctomycetes*. *FEMS Microbiol Rev* 42:739–760. <https://doi.org/10.1093/femsre/fuy029>
- Wiegand S, Jogler M, Boedeker C, Pinto D, Vollmers J, Rivas-Marín E, Kohn T, Peeters SH, Heuer A, Rast P, Oberbeckmann S, Bunk B, Jeske O, Meyerdieks A, Storesund JE, Kallscheuer N, Lückner S, Lage OM, Pohl T, Merkel BJ, Hornburger P, Müller R-W, Brümmer F, Labrenz M, Spormann AM, Op den Camp HJM, Overmann J, Amann R, Jetten MSM, Mascher T, Medema MH, Devos DP, Kaster A-K, Øvreås L, Rohde M, Galperin MY, Jogler C (2020) Cultivation and functional characterization of 79 planctomycetes uncovers their unique biology. *Nat Microbiol* 5:126–140
- Wiegand S, Rast P, Kallscheuer N, Jogler M, Heuer A, Boedeker C, Jeske O, Kohn T, Vollmers J, Kaster A-K, Quast C, Glöckner FO, Rohde M, Jogler C (2021) Analysis of bacterial communities on North Sea macroalgae

and characterization of the isolated planctomycetes *Adhaereton mobilis* gen. nov., sp. nov., *Roseimaritima multifibrata* sp. nov., *Rosistilla ulvae* sp. nov. and *Rubripirellula lacrimiformis* sp. nov. *Microorganisms* 9(7):1494. <https://doi.org/10.3390/microorganisms9071494>

Wilson K (2001) Preparation of genomic DNA from Bacteria. *Curr Protoc Mol Biol* 2:2.4.1–2.4.5. <https://doi.org/10.1002/0471142727.mb0204s56>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.