



Long-term N-addition alters the community structure of functionally important N-cycling soil microorganisms across global grasslands

Beat Frey^{a,*}, Barbara Moser^a, Bjorn Tytgat^b, Stephan Zimmermann^a, Juan Alberti^c, Lori A. Biederman^d, Elizabeth T. Borer^e, Arthur A.D. Broadbent^f, Maria C. Caldeira^g, Kendi F. Davies^h, Nico Eisenhauer^{i,j}, Anu Eskelinen^{i,k,l}, Philip A. Fay^m, Frank Hagedorn^a, Yann Hautierⁿ, Andrew S. MacDougall^o, Rebecca L. McCulley^p, Joslin L. Moore^q, Maximilian Nepel^r, Sally A. Power^s, Eric W. Seabloom^e, Eduardo Vázquez^t, Risto Virtanen^l, Laura Yahdjian^u, Anita C. Risch^a

^a Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland

^b Department of Biology, Laboratory of Protistology and Aquatic Ecology, Ghent University, Ghent, Belgium

^c Instituto de Investigaciones Marinas y Costeras (IIMyC), CONICET – UNMDP, Mar del Plata, Argentina

^d Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, IA, USA

^e Department of Ecology, Evolution, and Behavior, University of Minnesota, Saint Paul, MN, USA

^f Department of Earth and Environmental Sciences, The University of Manchester, Manchester, UK

^g Centro de Estudos Florestais, Instituto Superior de Agronomia, Universidade de Lisboa, Lisbon, Portugal

^h Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO, USA

ⁱ German Centre of Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

^j Institute of Biology, Leipzig University, Leipzig, Germany

^k Helmholtz Centre for Environmental Research UFZ, Leipzig, Germany

^l Department of Ecology and Genetics, University of Oulu, Oulu, Finland

^m USDA-ARS Grassland, Soil, and Water Research Laboratory, Temple, TX, USA

ⁿ Ecology and Biodiversity Group, Department of Biology, Utrecht University, Utrecht, the Netherlands

^o Department of Integrative Biology, University of Guelph, Guelph, ON, Canada

^p Department of Plant and Soil Sciences, University of Kentucky, Lexington, KY, USA

^q School of Biological Sciences, Monash University, Clayton Campus, Vic., Australia

^r Department of Biodiversity Research, University of Vienna, Vienna, Austria

^s Hawkesbury Institute for the Environment, Western Sydney University, Penrith, NSW, Australia

^t Department of Soil Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Bayreuth, Germany

^u Facultad de Agronomía, IFEVA, Universidad de Buenos Aires, CONICET, Buenos Aires, Argentina

ARTICLE INFO

Keywords:

Ammonia oxidizer
Biogeography
Diazotroph
Grassland
N-cycling microbial community
N-Fertilization
N₂-fixing bacteria
nifH
Nutrient network (NutNet)
Urea

ABSTRACT

Anthropogenic nitrogen (N) input is known to alter the soil microbiome, but how N enrichment influences the abundance, alpha-diversity and community structure of N-cycling functional microbial communities in grasslands remains poorly understood. Here, we collected soils from plant communities subjected to up to 9 years of annual N-addition (10 g N m⁻² per year using urea as a N-source) and from unfertilized plots (control) in 30 grasslands worldwide spanning a large range of climatic and soil conditions. We focused on three key microbial groups responsible for two essential processes of the global N cycle: N₂ fixation (soil diazotrophs) and nitrification (AOA: ammonia-oxidizing archaea and AOB: ammonia-oxidizing bacteria). We targeted soil diazotrophs, AOA and AOB using Illumina MiSeq sequencing and measured the abundance (gene copy numbers) using quantitative PCR. N-addition shifted the structure of the diazotrophic communities, although their alpha-diversity and abundance were not affected. AOA and AOB responded differently to N-addition. The abundance and alpha-diversity of AOB increased, and their community structure shifted with N-addition. In contrast, AOA were not affected by N-addition. AOA abundance outnumbered AOB in control plots under conditions of low N availability, whereas N-addition favoured copiotrophic AOB. Overall, N-addition showed a low impact on soil diazotrophs and AOA while effects for AOB communities were considerable. These results reveal that long-

* Corresponding author.

E-mail address: beat.frey@wsl.ch (B. Frey).

<https://doi.org/10.1016/j.soilbio.2022.108887>

Received 4 July 2022; Received in revised form 7 November 2022; Accepted 14 November 2022

Available online 15 November 2022

0038-0717/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

term N-addition has important ecological implications for key microbial groups involved in two critical soil N-cycling processes. Increased AOB abundance and community shifts following N-addition may change soil N-cycling, as larger population sizes may promote higher rates of ammonia oxidation and subsequently increase N loss via gaseous and soil N-leaching. These findings bring us a step closer to predicting the responses and feedbacks of microbial-mediated N-cycling processes to long-term anthropogenic N-addition in grasslands.

1. Introduction

Grasslands are globally important ecosystems that cover up to 25% of the terrestrial land surface and help to support more than 1.3 billion people through livestock production (Lieth, 1978; Rogiers et al., 2010). The productivity of natural grasslands depends on the level of nitrogen (N) inputs (Lebauer and Treseder, 2008; Fay et al., 2015). This is largely determined by biological N₂ fixation as a natural supply of N to grasslands which can reduce the use of N fertilizers. N₂ fixation stems from soil diazotrophs (N₂-fixing bacteria and archaea) that convert dinitrogen (N₂) gas to ammonia (NH₃) using the nitrogenase protein complex (*Nif*) (Reed et al., 2011; Vitousek et al., 2013; Van Groenigen et al., 2015). Diazotrophs are common in soils but their abundance, diversity and community structure are still poorly explored in grasslands in spite of their key role in global N input (Patra et al., 2006; Regan et al., 2017; Che et al., 2018). Variations in the diversity and composition of soil diazotrophs are strongly associated with vegetation type, climatic and soil parameters, whereas soil pH and moisture were the main influencing parameters (Collavino et al., 2014; Tu et al., 2016).

The N fixed by diazotrophs is directly available to plants (Vitousek et al., 2013). Nitrification is the biological oxidation of ammonia to nitrite followed by the oxidation of the nitrite to nitrate occurring through specialized microorganisms (Van Groenigen et al., 2015). Ammonia oxidation is a particularly important N-cycle process because it transforms the relatively immobile ammonium (NH₄⁺) to mobile nitrate (NO₃⁻) (Di et al., 2009; Di and Cameron, 2016), which is then prone to loss via leaching and denitrification (N₂O emission) (Liu et al., 2018; Tao et al., 2018). Overall, grazed grasslands contribute roughly 28% of global N₂O fluxes to the atmosphere (Rafique et al., 2011; Du et al., 2021). Ammonia oxidation therefore plays a key role in N losses and environmental pollution in grassland ecosystems (Risch et al., 2019). Ammonia-oxidizing archaea (AOA) and bacteria (AOB) are the most important ammonia oxidizers central to the fate of N in the environment (Nicol et al., 2008; Alves et al., 2013; Aigle et al., 2019). Besides AOA and AOB, complete ammonia oxidation (comammox) can also contribute to the conversion of ammonia to nitrite/nitrate (Daims et al., 2015). Apart from soil pH, vegetation and soil texture, N availability has been widely reported as a key driver of AOA and AOB community structure (Nicol et al., 2008; Gubry-Rangin et al., 2011; Liu et al., 2018).

Ammonia oxidizers (AOA and AOB) are tightly linked to the diazotrophs. Diazotrophs play a critical role in global N inputs into ecosystems, and ammonia oxidizers are essential for N turnover. The latter group of microorganisms are particularly important because they are involved in a bottleneck process in the global N-cycle (Lehtovirta-Morley, 2018). However, these three important N-cycling soil functional communities are rarely studied together. In particular, the possible ecological implications of shifts in the abundance, alpha-diversity and composition of these communities in response to increased N inputs in global grasslands are still largely unknown. For example, soil diazotrophs become less important and N₂ fixation less prevalent in environments with anthropogenic N inputs, since N₂ fixation is known to be a high energy process (Fan et al., 2019; Shi et al., 2021). This may imply that N-addition could replace N₂ fixation. On the contrary, higher N availability by N-addition could increase the abundance of ammonia oxidizers, in particular increasing the AOB activity, and consequently leading to higher N losses (Di and Cameron, 2016; Cardenas et al., 2019; Du et al., 2021). From local site grassland studies responses of soil diazotrophs and ammonia oxidizers to N-addition were inconsistent.

N-addition reduced (Zhang et al., 2013; Hu et al., 2021: N added as urea) or increased the abundance of soil diazotrophs (Ning et al., 2015: ammonium nitrate). N-addition as urea increased the abundance of AOA and AOB in grassland Loess Plateau soils (Wang et al., 2021), whereas urea addition in an alpine grassland soil of the Qinghai-Tibetan Plateau only increased the AOB abundance but AOA abundance remained unchanged (Xiang et al., 2017). In addition, most of these studies have focused on high-yield grasslands with high mean annual precipitation. Hence, these results limit our ability to extrapolate how N addition affects N-cycling microorganisms and underlines the need for global studies across a large range of climatic and edaphic factors. Recent global surveys on total microbial communities in grassland soils revealed strong regional scale patterns and nutrient addition effects at short-term (Leff et al., 2015), but global studies focusing on soil diazotroph and ammonia oxidizer responses to N-addition remain limited, especially longer-term responses.

Here, we sampled soils from 30 natural and semi-natural grasslands that span a globally relevant range of mean annual temperature and mean annual precipitation. We examined the impact of repeated N-addition (added as urea) on the three functionally important soil N-cycling microbial communities (soil diazotrophs, AOA, and AOB). We added granular, non-sulfur coated urea (NH₂)₂CO because it is a slow-release N fertilizer and causes less soil acidification (Cai et al., 2014). We also explored the main biotic and abiotic factors driving the geographic distribution of the functional gene abundance, observed richness, Shannon diversity index and community structure of soil diazotrophs, AOA, and AOB. This study is part of a larger effort within the Nutrient Network Global Research Cooperative to study soil microbial communities in global grasslands (Leff et al., 2015; Prober et al., 2015; Risch et al., 2019; Lekberg et al., 2021; Nepel et al., 2022). This cross-continent experiment allows the assessment of guild-level responses across treatments and global distributions. The advantage and power of this global study lies in the standardized treatments and exact replication across sites around the world. We are not aware of any previous studies in which diazotrophic and ammonia-oxidizing communities in grasslands were concomitantly assessed across diverse grassland communities. We hypothesized (H1) that long-term N-addition would reduce the abundance of soil diazotrophs, because N-addition can replace N₂ fixation (Fan et al., 2019). Further, we expected (H2) that AOA abundance would decrease with N-addition, because AOA thrive in nutrient-poor soils with low NH₄⁺ concentrations and that AOB abundance would increase because of enhanced NH₄⁺ availability leading to a decrease in the AOA:AOB ratio (Gubry-Rangin et al., 2011). We also hypothesized (H3) that alpha-diversity (observed richness and Shannon diversity index) and community structure of AOB will respond more strongly to N-addition than the diazotrophic and AOA communities because AOB perform better in N-rich environments (Sterngrén et al., 2015; Trivedi et al., 2019). These findings will allow us to better understand the responses of microbial-mediated N-cycling processes in global grassland ecosystems to N-fertilization and atmospheric N deposition.

2. Material and methods

2.1. Study sites and experimental design

Thirty sites from the Nutrient Network Global Research Cooperative (<https://nutnet.umn.edu>: <https://nutnet.umn.edu>) were used in this

study (Fig. 1, Tables S1 and S2). Thirteen sites overlapped the ones used in previous studies conducted earlier and represent a different soil sampling campaign (Leff et al., 2015; Prober et al., 2015). Most of the 30 sites were located in North America, Europe and Australia. A total of five sites were located in South America, Asia and Africa, where below-ground diversity and processes are under-studied (Guerra et al., 2020). Sites were subjected to 1–9 years of urea fertilization (Table S2), building on the previous studies of 2–4 years of fertilization (Leff et al., 2015; Prober et al., 2015). Our study therefore sheds light on impacts of N-addition across a larger global distribution and range of climatic and edaphic conditions, including mean annual temperature from -4.1°C to 22.3°C , mean annual precipitation from 252 mm to 1592 mm, and elevation from 6 m to 4261 m above sea level (Fig. 1, Table S1). Soil organic carbon (C) varies from 0.3% to 22%, soil total N from $<0.1\%$ to 1.3% and soil C:N ratio from 9.1 to 23.6 across our 30 sites (Table S2). Soil clay content (3.0%–53.3%) and soil pH (3.4–7.7) also spanned a large range across the 30 sites (Fig. 1, Table S2). Climate data for each site were obtained from WorldClim (<http://www.worldclim.org/current>, Table S1; Hijmans et al., 2005).

The full experimental design of the NutNet global experiment has been described by Borer et al. (2014). Fertilization treatments were applied to $5\text{ m} \times 5\text{ m}$ plots arranged in a randomized complete design with at least 3 replicates. Individual plots were at least 1 m apart. This study focused on untreated control and N addition, where N was added at 10 g m^{-2} per year (equal to 100 kg N ha^{-1} per year) as time-release urea $[(\text{NH}_2)_2\text{CO}]$. This rate corresponds to a typical fertilization regime of most grasslands but is significantly higher than natural N deposition in grasslands. Time-release urea was chosen because it causes less soil acidification than that associated with ammonium nitrate (Cai et al., 2014). The slow-release form releases urea during rain events, better approximating natural wet N deposition (Tian and Niu, 2015).

Soils were sampled from a $1\text{ m} \times 1\text{ m}$ sampling plot specifically designated for soil sampling (Borer et al., 2014). Each site received an identical soil sampling kit and soils were sampled as described in detail by Risch et al. (2019). Briefly, three 5 cm diameter soil samples (0–12 cm depth) per plot were collected. Two samples were composited, and one subsample of approximately 50 g was taken to analyse total C and N content, organic C content, NH_4^+ and NO_3^- concentration and pH (Risch et al., 2020). Another subsample was sieved (4 mm) and used to

determine the N-cycling functional microbial communities (six samples per site). The third $5 \times 12\text{ cm}$ subsample was collected and sealed in a steel cylinder, remaining undisturbed. From this subsample water holding capacity, bulk density and soil texture were assessed. All soil samples were collected 6–12 weeks prior to peak aboveground biomass at each site and were immediately shipped to the Swiss Federal Institute for Forest, Snow and Landscape Research WSL in Birmensdorf, Switzerland for laboratory analyses.

2.2. Laboratory procedures

The soil samples for chemical and particle size analyses were dried at 60°C . Soil particles were fractionated into sand, silt and clay using the pipette method according to Gee & Bauder (1986). Mineral soil pH was measured potentiometrically in 0.01 M CaCl_2 (Risch et al., 2019). A subsample of the dried samples was ground, 40 mg of soil from each sample was weighed into a tin capsule, and total carbon (TC) and total nitrogen (TN) contents were measured with a CHN analyser (NC-2500; CE Instruments, Wigan, United Kingdom). The organic C (Corg) content of samples with a pH (CaCl_2) < 6.0 was assumed to be equal to the total carbon (Ctot) content, whereas samples with a pH > 6.0 were assumed to potentially contain carbonates and, thus, were fumigated with HCl vapor prior to the CN analysis (Risch et al., 2019). In addition, an equivalent of 20 g dry soil was extracted with 1 M KCl (1:5 m/w) and NO_3^- and NH_4^+ concentrations were measured. Ammonium (NH_4^+) was measured photometrically with a FIAS 300 flow injection system (PerkinElmer, Waltham, MA, United States). The NO_3^- concentration was determined using UV-absorption but with CuSO_4 -coated zinc granules as a reducing agent (Risch et al., 2019). Plant species richness and cover (in %) of each plant species were assessed each year in a $1 \times 1\text{ m}$ subplot at each site (Borer et al., 2014).

2.3. Soil DNA extraction and quantitative PCR

Soil DNA was extracted from 0.30 g of fresh soil using a PowerSoil™ DNA Isolation Kit (MO BIO Laboratories, Carlsbad, CA, USA). To remove foreign DNA and prevent microbial contaminations, working bench surfaces and non-autoclavable materials were cleaned with 5% sodiumhypochlorite and 70% ethanol solutions prior to the DNA




Fig. 1. Geographic and climatic distribution of experimental sites. (A) Location of the 30 Nutrient Network sites where the field experiment was conducted and soil samples were collected for laboratory analyses. (B) The 30 study sites represent a wide range of mean annual temperature (MAT) and mean annual precipitation (MAP) conditions (Table S1). Our sites also cover a wide range of soil edaphic conditions, as described in the main text and shown in Table S2. Numbers refer to # in Tables S1 and S2.

extractions. The extracted DNA was quantified with Quant-it Pico Green dsDNA Assay (Thermo Fisher Scientific, Reinach, Switzerland). Soil diazotrophs and ammonia oxidizers have been extensively investigated with molecular marker methods (Levy-Booth et al., 2014). The *nifH* gene family, encoding the dinitrogenase reductase subunit, is widely used to gain insight into diazotrophic communities (Hsu and Buckley, 2009; Gaby and Buckley, 2012; Collavino et al., 2014). The *amoA* gene, encoding subunit A of ammonia monooxygenase, is commonly used to investigate AOA and AOB (Tourna et al., 2008; Nicol et al., 2008). The abundance of functional marker genes encoding archaeal and bacterial ammonia monooxygenase (archaeal *amoA*, bacterial *amoA*) and the dinitrogenase gene (*nifH*) were quantified using quantitative PCR (qPCR). The primers and thermocycling conditions for bacterial *amoA* (AmoA-1F and AmoA-2R; Rothauwe et al., 1997), archaeal *amoA* (Arch-amoAF and Arch-amoAR; Francis et al., 2005) and *nifH* (PolF and PolR; Poly et al., 2001) are reported in Table S3. Negative controls for the DNA extractions (extraction buffer without soil) and PCR amplifications (high-purity water without DNA template) were included. PCR for each sample was performed in triplicate and negative controls were included in each batch of PCR.

All qPCR assays were carried out with equal amounts of template DNA (20 ng DNA) in 20 μ l reactions using QuantiTect SYBR Green PCR master mix (Qiagen, Hilden, Germany) on an ABI7500 Fast Real-Time PCR system (Applied Biosystems, Foster City, CA, USA) as previously outlined (Frey et al., 2011, 2021). Standards for qPCR were generated by serial dilution of stocks containing a known number of plasmids carrying the respective functional gene as an insert (Frey et al., 2020). The specificity of the amplification products was confirmed by melting-curve analysis, and the expected sizes of the amplified fragments were checked in a 1.5% agarose gel stained with ethidium bromide. The amplification efficiencies of qPCR assays were 95% (± 2) for archaeal *amoA*, 93% (± 3) for bacterial *amoA* and 98% (± 2) for *nifH*. R^2 values were >0.99 for all runs. The gene copy number was calculated using a regression equation that related the cycle threshold (Ct) value to the known number of copies in the standards. For each soil sample, the qPCR reactions were repeated three times. Abundances of bacterial and archaeal *amoA* and *nifH* genes refer to copy numbers per gram of dry soil.

2.4. MiSeq sequencing and bioinformatic analyses

PCR amplifications of the bacterial *amoA* and *nifH* genes were performed as described previously with primers similar to those used in qPCRs but containing adapters for MiSeq Illumina sequencing. Due to the limits of sequence lengths using the MiSeq platform (2×300 bp), primer amplification fragments of archaeal *amoA* were too long (629 bp) to be assembled; therefore, another primer set was used (Arch-amoA-104F and Arch-amoA-616R; Tourna et al., 2008; Alves et al., 2013) which generated a shorter amplicon (Table S3). Each sample consisting of 40 ng DNA was amplified in triplicate and pooled before purification with Agencourt AMPure XP beads (Beckman Colter, Brea, CA, USA) and quantified with the Qubit 2.0 fluorometric system (Life Technologies, Paisley, UK). Amplicons were sent to the Genome Quebec Innovation Centre (Montreal, Canada) for barcoding using the Fluidigm Access Array technology and paired-end V3 MiSeq sequencing (Illumina Inc., San Diego, CA, USA), enabling production of 2×300 -bp reads.

Raw *nifH* and bacterial and archaeal *amoA* reads were each merged using PEAR v.1.9.4 (Zhang et al., 2014) with a minimum overlap of 10 bp between forward and reverse paired end reads. Quality filtering was performed on the paired reads using USEARCH v.9.2.64 (Edgar, 2010) removing sequences with a Phred score below Q30, exceeding a per read maximum expected error (maxEE) of 0.5 or containing ambiguous bases. Primers were subsequently trimmed off using CUTADAPT v.1.9 (Martin, 2011) set with an error tolerance of 0.15 and 0.20 for the forward and reverse primers, respectively. Next, possible frameshifts were detected and corrected to frame 1 using an inhouse Python script. Any reads containing stop codons (TAG, TAA, TGA), indicating erroneous indels,

were removed. Reads were then sorted according to abundance after which putative chimeras were removed from the dataset using the UCHIME algorithm (Edgar et al., 2011). High quality sequences were assigned to operational taxonomic units (OTUs) using the UCLUST algorithm (Edgar, 2010) at a 5% nucleotide dissimilarity for *amoA* and *nifH* gene sequences (Gaby and Buckley, 2011; Pester et al., 2012; Penton et al., 2016; Carrell et al., 2019), with -minsize set to 2 for the *nifH* gene. Prior to diversity analyses, OTUs were rarefied to 11025 (*nifH*), 7165 (AOA *amoA*) and 9427 (AOB *amoA*) reads per sample. OTU-based alpha-diversity was reported as observed richness and Shannon diversity index, which includes community richness and evenness in the metric. Differences between community structure (beta-diversity) were estimated based on the Bray-Curtis distance. Raw sequences were uploaded in the NCBI Sequence Read Archive (SRA) with BioProject accession number PRJNA739076.

2.5. Statistical analyses

Some of the explanatory variables (Table S4) were highly skewed and were thus log or square-root transformed prior to analyses. All continuous explanatory variables were centred and scaled to have a mean of zero and variance of one. Variables were additionally filtered to avoid collinearity using correlation analysis. If variables were strongly correlated (Pearson's $|r| > 0.70$) (Dormann et al., 2013), those that allowed us to minimize the number of variables were selected (Fig. S1, Table S4). Consequently, soil total C concentration, total N concentration, sand content and bulk density were removed from the dataset. For climatic variables mean annual temperature, mean annual precipitation, temperature of the wettest quarter and temperature seasonality (standard deviation; hereafter temperature variability; Hijmans et al., 2005) were used, as these variables were not correlated under the definition given above (Fig. S1, Table S4).

To assess N-addition we fit linear mixed-effects models (LMMs). observed richness, Shannon diversity index, gene abundance (gene copy numbers) of all three groups (*nifH*, AOA, AOB), and the ratio of AOA to AOB as response variables using the 'lme' function in the *nlme* package (version 3.131.1; Pinheiro et al., 2021) in R version 3.4.4 (R Core Team, 2020). Treatment (N addition, control) was included as a fixed factor and block nested within site was included as a random factor. The number of years since the N-addition treatment began was also included as a fixed effect in all the initial models. However, with the exception of AOA Shannon diversity index and AOB richness, this factor was not significant and therefore was not retained in the models. To examine global biotic, edaphic and climatic drivers (Table S4) of *nifH*, AOA, and AOB alpha-diversity (observed richness and Shannon diversity index), gene abundance and the AOA:AOB ratio in grassland soils, multi-model inference and LMMs were used. Separate models were run for each response variable. The explanatory variables together with treatment were included as fixed effects. In addition, significant interaction terms between the treatment and explanatory variables were included. Block nested within site was again included as a random effect in these models. First, full models including all predictor variables and relevant interaction terms were fitted and then the R package *MuMin* (version 1.42.1; cran.r-project.org) was used to select the best models that explained the most variation based on Akaike's information criterion (AIC; 'model.avg' function). The corrected AIC (AICc) was used to account for the small sample size. Model averages are presented in the main text and all best models (models that fell within 2 AICc units of the best model; delta AICc < 2) are given in Tables S6 and S7. To assess if the significant explanatory variables contained in the top models had opposite trends of N additions on alpha-diversity (observed richness and Shannon diversity index) and gene abundance of *nifH*, AOA and AOB as well as the AOA to AOB ratio, e.g., if the response to N additions differed between dry and wet sites, or at sites with high versus low pH we calculated the log response ratio (LRR) for gene abundance and alpha-diversity of *nifH*, AOA and AOB as well as for the AOA:AOB ratio. The LRR was then

plotted against the explanatory variables of the control plots. To assess whether these relationships were significant, we again used LMMs.

PERMANOVA analyses (10^5 permutations) were conducted to test for significant differences in diazotrophic (*nifH*), AOA and AOB community structure between control and N-added plots, sites and their interaction. Site was included as a random factor using the statistical package of PrimerE v7 (Clarke and Gorley, 2015). Beta-diversity was visualized using canonical analysis of principal coordinates (CAP) based on Bray-Curtis dissimilarity distances. To explore environmental correlations with the beta-diversity of diazotrophic and ammonia-oxidizing communities (AOA and AOB), pairwise Bray-Curtis dissimilarities between all plots were calculated using *vegan*. Dissimilarities were averaged per treatment level within site-pairs to avoid pseudo-replication (Prober et al., 2015). Final correlations between environmental factors and beta-diversity were thus comparisons between sites at the treatment level, averaged from plot-level dissimilarities. Multi-model inference and linear models were then used to assess environmental correlates of between-site dissimilarities in diazotrophic and ammonia-oxidizing communities (AOA and AOB). The model selection procedure was identical to that described above for alpha-diversity, gene abundance and AOA:AOB ratio.

3. Results

3.1. N-addition effects, and drivers of gene abundance and alpha-diversity

AOB alpha-diversity (observed richness and Shannon diversity index) and abundance were significantly higher and the AOA:AOB ratio was significantly lower in N-added plots than in control plots (Fig. 2, Table S5, Figs. S2–S6 in the Supplementary Results). N-addition did influence *nifH* richness but not *nifH* Shannon diversity index and abundance. AOA abundance and alpha-diversity were not affected by N-addition (Fig. 2, Table S5). We detected higher *nifH* alpha-diversity in N-added plots at sites with higher MAP, while larger *nifH* gene copy numbers were correlated with higher soil organic C content, higher temperature of the wettest quarter, lower temperature variability and higher MAP across the sites (model average results are given in Table 1, whereas climatic, soil physico-chemical and vegetation parameters included in all best models [within an AICc of 2] are given in Table S6).




Fig. 2. N-addition effects on diazotrophic (*nifH*), AOA (archaeal *amoA*) and AOB (bacterial *amoA*) alpha-diversity (richness and Shannon diversity index) and abundance, as well as, AOA:AOB ratio across global grasslands. Results are based on linear mixed-effects models, with block nested in site included as a random effect. * Indicates a significant treatment effect (alpha-level = 0.05). Data points are effect sizes (N-addition) and error bars represent standard errors (SE). For statistics see Table S5.

In contrast, we were not able to explain AOA Shannon diversity index across our sites with the climatic, soil or plant variables available, although higher AOA richness was related to higher soil pH and soil organic C content. Similarly, higher abundances of AOA were correlated to higher soil pH (Table 1, Table S6). AOB Shannon diversity index and AOB abundance were both higher in N-added plots than in controls (Fig. 2). Besides of the N-addition treatment, AOB abundance was further explained by multiple factors (i.e., interaction between N-addition treatment and MAT; interaction between N-addition treatment and temperature of the wettest quarter; and NO_3^- content), whereas AOB Shannon diversity index only by the treatment (Table 1, Table S6). AOB richness was explained by soil pH and NO_3^- content. The N-addition treatment was the only significant predictor for differences in the AOA:AOB ratio across the global grasslands (Fig. 2, Table 1, Table S6). There were no clear trends in the LRR ratios of gene abundance and diversity of *nifH*, AOA and AOB as well as the AOA to AOB ratio in relationship to the significant variables included in the models (Figs. S7–S10).

3.2. N-addition effects and drivers of community structure

PERMANOVA analysis revealed a strong N-addition effect on diazotrophic (*nifH*) and AOB community structure (Table 2, Fig. 3). AOB communities additionally differed significantly among the 30 sites and also showed a significant treatment \times site interaction (Table 2). AOA communities were, in contrast, not affected by N-addition. Beta-diversity of the N-cycling communities (dissimilarity among sites) was correlated with both climatic and soil variables, while vegetation properties did not play a large role (Table 3, Table 4, Table S7). Differences between sites in terms of MAP, geographic distance, soil pH, clay, MAT, C:N ratio and temperature of the wettest quarter explained the dissimilarities in diazotrophic communities across the global grasslands. Strongest predictors in the dissimilarities in diazotrophic communities were MAP and geographic distance whereas C:N ratio and temperature were weakest (Table 3). The beta-diversity of AOA communities was related to differences among sites in MAT and in soil pH, water holding capacity, clay content and C:N ratio (Table 3, Table S7). The strongest predictors of AOB beta-diversity were differences in soil pH, MAT and geographic distance, whereas N-addition, the interaction between N-addition and the difference in C:N ratio, MAP and NO_3^- concentration had less, but still significant, explanatory power (Table 3).

Table 1

Effects of significant environmental drivers on diazotrophic (*nifH*), AOA (archaeal *amoA*) and AOB (bacterial *amoA*) alpha-diversity (observed richness and Shannon diversity index) and abundance, and the AOA:AOB ratio across global grasslands. Model averaged coefficients for the full average are shown. For all best models for each variable see Table S6.

	Estimate	SE	z-value	p-value
<i>NifH</i> richness ($R^2m = 0.47$, $R^2c = 0.95$)				
Intercept	5.11	0.17	29.56	<0.001
bioclim.P.ann	0.51	0.18	2.72	0.007
no.plspecies.nfixer	0.14	0.04	3.62	<0.001
Treatment	0.19	0.04	4.17	<0.001
<i>NifH</i> Shannon ($R^2m = 0.46$, $R^2c = 0.88$)				
Intercept	3.51	0.15	23.57	<0.001
bioclim.P.ann	0.45	0.16	2.70	0.006
<i>NifH</i> abundance ($R^2m = 0.59$, $R^2c = 0.92$)				
Intercept	16.44	0.17	97.10	<0.001
ch.C.org.perc.log	0.51	0.15	3.37	<0.001
bioclim.T.q.wet	0.94	0.30	2.99	0.003
bioclim.T.var	-0.74	0.28	2.58	0.01
bioclim.P.ann	0.52	0.19	2.55	0.01
AOA richness ($R^2m = 0.49$, $R^2c = 0.90$)				
Intercept	2.60	0.06	39.5	<0.001
ch.C.org.perc.log	-0.17	0.04	3.60	<0.001
ch.pH	0.22	0.04	4.96	<0.001
AOA Shannon ($R^2m = 0.36$, $R^2c = 0.77$)				
Intercept	1.20	0.22	5.38	<0.001
AOA abundance ($R^2m = 0.25$, $R^2c = 0.77$)				
Intercept	2.57	0.02	102.0	<0.001
ch.pH	0.06	0.03	2.25	0.02
AOB richness ($R^2m = 0.18$, $R^2c = 0.79$)				
Intercept	3.96	0.20	19.30	<0.001
ch. NO ₃ start.mgg.sqrt	0.28	0.08	3.27	0.001
ch.pH	0.43	0.15	2.78	0.005
AOB Shannon ($R^2m = 0.16$, $R^2c = 0.80$)				
Intercept	2.30	0.16	13.88	<0.001
Treatment	0.35	0.08	4.42	<0.001
AOB abundance ($R^2m = 0.38$, $R^2c = 0.78$)				
Intercept	12.49	0.38	32.04	<0.001
Treatment	1.63	0.26	6.15	<0.001
bioclim.T.ann × Treatment	-1.30	0.25	5.05	<0.001
ch. NO ₃ start.mgg.sqrt	0.89	0.20	4.34	<0.001
bioclim.T.q.wet × Treatment	1.08	0.26	4.19	<0.001
AOA:AOB ($R^2m = 0.22$, $R^2c = 0.69$)				
Intercept	0.08	0.04	2.31	0.02
Treatment	-0.15	0.03	4.68	<0.001

Predictors were.

Climate: bioclim.P.ann = MAP (mm); bioclim.Tvar = Tvar (SE monthly means * 100); bioclim.T.q.wet = T wettest quarter (°C).

Soil chemistry: ch.pH = pH (CaCl₂); ch.NO₃start.

g.sqrt = NO₃⁻ (mg kg⁻¹); ch.C.org.perc.log = organic C.

Vegetation: no.plspecies.nfixer = Number of N-fixing plant species.

Treatment = N-addition.

Table 2

PERMANOVA results for treatment, site and treatment × site interaction in diazotrophic (*nifH*), AOA (archaeal *amoA*) and AOB (bacterial *amoA*) community structure, based on 9999 permutations. Significant effects are highlighted in bold ($P < 0.05$). Site was included as a random factor.

	DF	F-value	p-value
<i>NifH</i> community structure			
Treatment	1	1.819	< 0.001
Site	29	15.749	0.616
Treatment × site	29	0.925	0.718
AOA community structure			
Treatment	1	0.602	0.865
Site	29	32.811	0.984
Treatment × site	29	0.603	0.981
AOB community structure			
Treatment	1	2.004	< 0.001
Site	29	9.367	0.002
Treatment × site	29	1.499	0.003

4. Discussion

4.1. Impact of N-addition on diazotrophs, AOA and AOB abundance

The abundance of *nifH* copy numbers was not affected by N-addition, which does not support our first hypothesis (H1). We expected that soil diazotrophs would become less abundant in environments with increased N inputs, because N₂ fixation is an energetically expensive process and the diazotrophs are outcompeted under high N availability by other non-fixing microorganisms (Fan et al., 2019; Shi et al., 2021). In contrast to our study, higher available N content in soils as a consequence of N-addition has been shown to negatively affect *nifH* gene copy numbers in other grassland soils (Zhang et al., 2013; Hu et al., 2021), as the dinitrogenase enzyme is sensitive to NH₄⁺ (Reed et al., 2011; Han et al., 2019). In a long-term study (40 years), urea addition was found to be detrimental for obligate N₂ fixers (Fan et al., 2019). The addition of N was shown to decrease symbiotic N₂-fixation in grasslands, including many of the sites in this study (Vázquez et al., 2022). The negative effect of N addition on symbiotic N₂ fixation by legumes per area was caused exclusively by the negative effect of N addition on legume biomass (Tognetti et al., 2021), and not by an effect on the N₂ fixation rate per unit biomass. In addition to symbiotic N₂ fixing microorganisms, Rui et al. (2022) reported that free-living diazotrophs may also play important roles in biological N₂ fixation of grasslands. The capacity for biological N₂ fixation relies on the nitrogenase enzyme system (Reed et al., 2011). The *nifH* gene encodes the nitrogenase reductase subunit, which is the most widely sequenced biomarker for identifying diazotrophs (Gaby and Buckley, 2012). However, *nifH* gene abundance cannot differentiate between the free-living and symbiotic diazotrophs (Rui et al., 2022). Non-fixing plants (grasses and forbs) outcompeted legumes under high N availability and a decrease of N₂ fixation could be expected. Vegetation (diversity and biomass) is a crucial factor affecting soil diazotrophs, which are highly dependent on easily available carbon sources from root exudates (Reed et al., 2011; Collavino et al., 2014). It is possible that a period of one to nine years of N-addition in our current study was not long-term enough to cause such changes.

The quantity, form and duration of N-addition play a crucial role in the N availability and acidification of the soils after N-addition (Tian and Niu, 2015). In some of the long-term grassland sites considered here (>7 years of treatment), urea fertilization acidified the soils (pH decrease >0.4 in burrawan.au, cbgb.us, cdcr.us, cdpt.us, valm.ch), but this decrease in pH ($p = 0.004$) apparently had no influence on the abundance of *nifH* gene copies. This is in accordance with Leff et al. (2015) who also included soils from some sites analyzed in our study. Overall,




Fig. 3. Canonical analysis of principal coordinates (CAP) ordinations of the three N-cycling microbial communities: (A) diazotrophic (*NifH*), (B) AOA (archaeal *amoA*) and (C) AOB (bacterial *amoA*) communities of the 30 grassland sites. Each site is represented by six samples consisting of triplicates for each treatment (open symbols = control; closed symbols = N-addition). N = 180.

Table 3

Effects of nitrogen (N) addition and differences in site characteristics (climatic, soil and vegetation variables) on dissimilarities of diazotrophic (*nifH*), AOA (archaeal *amoA*) and AOB (bacterial *amoA*) community structure (beta-diversity). Differences in site characteristics and dissimilarities in microbial communities (Bray-Curtis distance) were averaged for each treatment level within site-pairs. Model averaged coefficients (full average) are shown for those predictors that occurred in at least 50% of all models. For all best models see Table S7. Significant effects are highlighted in bold ($P < 0.05$). No. models = number of models where a variable was included. Rel. imp. = relative importance, Δ = difference between two sites. Definition of predictors see Table S4.

	Estimate	SE	z-value	p-value	No. models	Rel. imp.
<i>NifH</i> dissimilarity ($R^2 = 0.21$)						
Intercept	0.9805	0.0009	1018.6	<0.001		
Δ bioclim.P. ann.sqrt	0.0061	0.0010	6.2	<0.001	24	1.00
distGeo.km.sqrt	0.0055	0.0009	5.6	<0.001	24	1.00
Δ ch.pH	0.0053	0.0010	5.3	<0.001	24	1.00
Δ ch.clay.perc.sqrt	0.0042	0.0010	4.2	<0.001	24	1.00
Δ bioclim.T. ann.sqrt	0.0041	0.0010	4.0	<0.001	24	1.00
Δ ch.CtoN.sqrt	0.0025	0.0010	2.5	0.012	24	1.00
Δ bioclim.T.q.wet.sqrt	0.0023	0.0010	2.1	0.031	24	1.00
Δ ch.WHC.vol.sqrt	0.0017	0.0012	1.4	0.162	19	0.83
Δ no.plspecies.nfixer.sqrt	0.0014	0.0011	1.1	0.234	18	0.78
Δ ch.Corg.perc.sqrt	0.0010	0.0012	0.8	0.385	15	0.60
<i>AOA</i> dissimilarity ($R^2 = 0.14$)						
Intercept	0.9331	0.0036	256.2	<0.001		
Δ ch.pH	0.0199	0.0033	6.0	<0.001	16	1.00
Δ bioclim.T. ann.sqrt	0.0191	0.0034	5.7	<0.001	16	1.00
Δ ch.WHC.vol.sqrt	0.0146	0.0034	4.3	<0.001	16	1.00
Δ ch.clay.perc.sqrt	0.0117	0.0034	3.4	<0.001	16	1.00
Δ ch.CtoN.sqrt	0.0111	0.0034	3.2	0.001	16	1.00
Δ ch.NO ₃ start.mgg.sqrt	0.0064	0.0036	1.8	0.076	15	0.95
Δ bioclim.P. ann.sqrt	0.0053	0.0037	1.4	0.156	14	0.88
Δ bioclim.T. var.sqrt	0.0028	0.0035	0.8	0.419	9	0.57
<i>AOB</i> dissimilarity ($R^2 = 0.19$)						
Intercept	0.9659	0.0029	336.6	<0.001		
ch.pH	0.0171	0.0020	8.5	<0.001	12	1.00
Δ bioclim.T. ann.sqrt	0.0137	0.0020	6.7	<0.001	12	1.00
Δ distGeo.km.sqrt	0.0097	0.0020	4.8	<0.001	12	1.00
Treatment	-0.0090	0.0042	2.1	0.032	12	1.00
Δ ch.CtoN.sqrt × Treatment	0.0086	0.0039	2.2	0.028	12	1.00
Δ bioclim.P. ann.sqrt	0.0055	0.0020	2.7	0.007	12	1.00
ch.NO ₃ start.mgg.sqrt	0.0043	0.0021	2.0	0.042	12	1.00
Δ ch.CtoN.sqrt	-0.0016	0.0030	0.6	0.580	12	1.00
Δ ch.WHC.vol.sqrt	0.0031	0.0023	1.3	0.180	10	0.84
Δ ch.clay.perc.sqrt	0.0016	0.0021	0.8	0.447	6	0.53

Table 4

Summarizing the variables for predicting diazotrophic (*nifH*), AOA (archaeal *amoA*) and AOB (bacterial *amoA*) abundance, alpha-diversity (richness and Shannon diversity index) and community structure, and the AOA:AOB ratio. For details about individual variables see Table S4. Tvar = temperature seasonality, T = temperature, P = precipitation, log = transformed using log, sqrt = transformed using square root.

	<i>nifH</i>	AOA	AOB
Abundance	multiple factors (ch.Corg.perc.log; bioclim.T.q.wet; bioclim.Tvar; bioclim.P.ann)	ch.pH	N-addition; bioclim.T.ann x N-addition; chNO ₃ start.mgg.sqrt; bioclim.T.q.wet x N-addition
Richness	bioclim.P.ann no.plspecies.nfixer	ch.pH ch.C.org.perc.log	ch.pH chNO ₃ start.mgg.sqrt
Shannon	bioclim.P.ann		N-addition
Community structure	N-addition and multiple factors (bioclim.T.ann.sqrt; distGeo.km.sqrt; ch.pH; ch.clay.perc.sqrt; bioclim.T.ann.sqrt; ch.CtoN.sqrt; bioclim.T.q.wet.sqrt)	Multiple factors (ch.pH; bioclim.T.ann.sqrt; ch.WHC.vol.sqrt; ch.clay.perc.sqrt; ch.CtoN.sqrt)	N-addition and multiple factors (ch.pH; bioclim.T.ann.sqrt; distGeo.km.sqrt; ch.CtoN.sqrt x N-addition; bioclim.P.ann.sqrt; ch. NO ₃ start.mgg.sqrt)
AOA:AOB	N-addition		

Predictors were.

Climate: bioclim.P.ann = MAP (mm); bioclim.Tvar = Tvar (SE monthly means * 100); bioclim.T.q.wet = T wettest quarter (°C); bioclim.T.ann = MAT (°C).

Soil chemistry: ch.pH = pH (CaCl₂); ch.NO₃start.mgg.sqrt = NO₃⁻ (mg kg⁻¹); ch.C.org.perc.log = organic C; ch.clay.perc = clay content (%); ch.CtoN.sqrt = C:N ratio; ch.WHC.vol.sqrt = Water-holding capacity (vol%).

Vegetation: no.plspecies.nfixer = Number of N-fixing plant species.

Geography: distGeo.km.sqrt = Geographic distance between two sites.

Treatment = N-addition.

changes in the soil environment were surprisingly small which might explain the lack of responses in terms of the *nifH* gene abundance in our study. Although we did not observe N-addition effects on *nifH* gene abundance, soil organic C, together with climatic properties, had the strongest positive effects on *nifH* gene abundance. Edaphic variables, such as organic C, NH₄⁺, NO₃⁻, and available P content have been widely recognized as crucial factors affecting the abundance of soil diazotrophs (Reed et al., 2011; Reardon et al., 2014; Han et al., 2019).

AOA and AOB responded differently to N-addition. The gene abundance of AOB increased whereas AOA were not affected by N-addition. We expected that AOA abundance would decrease with N-addition (H2) because AOA thrive in nutrient-poor soils with low NH₄⁺ concentrations and that AOB abundance would increase because of enhanced NH₄⁺ availability. Hence, we only found partial support for our hypothesis. Yet, our findings are in agreement with other research where N-addition has been shown to positively affect (Verhamme et al., 2011) or have no influence on AOA abundance (Chen et al., 2013), whereas AOB abundance flourished with N enrichment (Xiang et al., 2017; Zhang et al., 2019). However, these earlier findings are mostly from local-scale studies.

Dominance of AOA relative to AOB in the *amoA* gene pool, as we found here, has also been reported in grasslands and agricultural environments (He et al., 2007; Gubry-Rangin et al., 2011; Zhou et al., 2014; Sterngren et al., 2015). This could be attributed to a difference in the affinity for ammonia caused by the unique physiological characteristics and preferred habitat conditions of AOA and AOB (Prosser and Nicol, 2012). AOB are typically copiotrophic (Hatzenpichler, 2012) and are most abundant in soils that have a relatively high N content (Sterngren et al., 2015; Trivedi et al., 2019). AOA perform better in habitats with more acidic, organic matter-depleted soil conditions (Banning et al., 2015; Trivedi et al., 2019) which is strongly supported by our findings in some of the grassland soils (e.g. spin.us; sgs.us; kibber.in). AOA also appear to be less constrained by the availability of N substrates and prosper in oligotrophic environments (Martens-Habbena et al., 2009; Trivedi et al., 2019), and our results confirm this pattern. In addition, AOA have lower membrane permeability, smaller cell and genome size, and higher energy efficiency than AOB, which enables AOA to better adapt to and perform their physiological activities in nutrient-limited environments (Prosser and Nicol, 2012; Banning et al., 2015). Burton and Prosser (2001) suggested that AOB in contrast to AOA can produce urease and thus respond directly to urea addition which might be

responsible for the stronger response of AOB compared to AOA to N-addition in our study. These and potentially other physiological or metabolic differences between AOA and AOB likely lead to niche differentiation between these two groups (Prosser and Nicol, 2012; Carey et al., 2016; Aigle et al., 2019), and therefore different responses to N-addition. Overall, our results suggest that AOB perform better in nutrient-rich environments than AOA when faced with an increased N input (H2).

As observed in our study, N-addition may change the dynamics of microbial-mediated N-cycling processes. Diazotrophs and both AOA and AOB are tightly linked (Van Groenigen et al., 2015). Therefore, shifts in the abundance of these functionally important microbial groups with the high N dose applied in this study are likely to have ecological implications in global grasslands. N-addition may disrupt the soil N-cycling by increasing the abundance of AOB. Larger population sizes of AOB may promote higher rates of ammonia oxidation (Banning et al., 2015; Carey et al., 2016), and subsequently change the availability of oxidized forms of N and thus their mobility in soils (Yang et al., 2018). In addition, higher AOB abundance may also increase N₂O emissions, which is a potent greenhouse gas (Hink et al., 2017; Cardenas et al., 2019; Du et al., 2021). Taken together, our findings suggest that N-fertilization has important ecological implications for the abundance of functionally important microbial groups. Specially, an increased abundance of ammonia oxidizers with N-addition may increase nitrification activity, which likely leads to N losses and environmental pollution.

4.2. Impact of N-addition on diazotrophs, AOA and AOB alpha-diversity and community structure

We also hypothesized (H3) that the alpha-diversity of AOB respond more strongly to N-addition than the diazotrophic and AOA communities because AOB have a more copiotrophic lifestyle than the other two groups. Our findings partially confirmed our expectations. AOB alpha-diversity (observed richness and Shannon diversity index) increased with N-addition, but N input had no significant effect on the AOA alpha-diversity and 'Treatment' was not included in any of the best models we developed for AOA (Table 4). The AOB response that we observed contradicted previous work that showed that N input had no significant effect on the AOB (or AOA) alpha-diversity in a semi-arid grassland on China's Loess Plateau (Wang et al., 2021). Furthermore, we did find a treatment effect on *nifH* richness, which contrasts Zhu et al. (2018) who

reported no effect on diazotroph richness with urea applied at the same rate as our study. However, at higher rates (150 and 300 kg urea ha⁻¹ per year), diazotroph richness did decrease (Zhou et al., 2021) suggesting a certain threshold of N addition levels for the *nifH* richness. The best predictor of *nifH* alpha-diversity in our study was MAP, with higher *nifH* diversity at sites with higher MAP, whereas soil pH and NO₃⁻ concentration were the best variables in explaining *nifH* diversity in agricultural land (Zhou et al., 2021), but we are not aware of a study that investigated these interactions in global grassland ecosystems.

The AOB community structure responded strongly to N-addition whereas the AOA community structure did not (Table 4). This finding is in agreement with our third hypothesis (H3) and provides new insight into the responses of AOA and AOB community structure to N-addition in global grassland ecosystems. Earlier findings are mostly from single site studies. Xiang et al. (2017) showed a significant shift in AOB but not AOA community structure within short-term N urea amendment in an alpine grassland soil of the Qinghai-Tibetan Plateau. Similarly, Zhang et al. (2019) reported that N-fertilization (as NH₄NO₃ at 10 g N m⁻² year⁻¹ for 11 years) significantly affected AOB community structure over the AOA counterparts in a semi-arid grassland system in Northern China. The most likely explanation for these findings is that most of the added N enters the soil in the form of NH₄⁺, which AOB and AOA transform to NO₃⁻ thereby producing H⁺ (nitrification) (Patra et al., 2006; Van Groenigen et al., 2015). Fertilization-induced changes in pH may partly explain the N-fertilization effects on AOB community structure. However, changes are more likely due to direct effects of increased N availability and interactions between AOA and AOB (Zhang et al., 2019). Overall, these findings demonstrate that AOB were more sensitive than AOA in response to urea suggesting that N fertilizer may be an important controller of ammonia oxidizers. Besides that, AOA communities were not affected by N-addition. We could also not detect an obvious site effect on AOA community structure in our global grassland study as found for AOB. Overall, the AOA richness was low (193 OTUs; Supplementary Results) compared to the higher AOB richness (1418 OTUs) and some of the dominant AOA taxa are known to have a cosmopolitan distribution (Alves et al., 2018).

Furthermore, we found that N-addition changed the diazotrophic community structure, even though the Shannon diversity index and abundance of *nifH* gene copy numbers were not affected (Table 4). This agrees with previous findings that diazotrophic communities are sensitive to N-addition in arable soils (Wang et al., 2017a; Zhou et al., 2021). Shifts in the *nifH* community structure under longer-term N-addition may lower the capacity for biological N₂ fixation, particularly in nitrogen-poor soils (Feng et al., 2018; Fan et al., 2019; Tognetti et al., 2021). This may have long-term ecological implications for soil diazotrophs and microbial-mediated N-cycling processes in grassland ecosystems.

Soil pH was one of the main predictors of the community structure of our three key microbial groups. All models included soil pH as the main influencing soil factor (Table 4). The importance of pH is consistent with previous work demonstrating that soil pH is the primary determinant for shaping total microbial community structure at the continental scale (Lauber et al., 2009; Fierer, 2017). In our study, both AOA and AOB communities were strongly affected by soil pH, which is in line with findings from single site studies (Ying et al., 2017; Zhang et al., 2019; Vázquez et al., 2020; Wang et al., 2021). Recent work by Wang et al. (2017b) and Zhou et al. (2021) also highlighted the importance of soil pH as a fundamental driver shaping the diazotrophic community structure. We reached the same conclusion here with the global grassland soils.

5. Conclusions

Diazotrophic and ammonia-oxidizing microorganisms play crucial roles in global N cycle. Our study provides new insight into the responses of three functionally important N-cycling microbial groups to N-addition

in grassland ecosystems. Annual addition of 10 g N m⁻² year⁻¹ for up to 9 years did not affect *nifH* gene abundance or Shannon diversity but changed the *nifH* community structure across the global grassland sites. AOB responded more dramatically to N-addition than diazotrophs or AOA, with increases in gene copy numbers and alpha-diversity and a shift in community structure. N-addition favoured copiotrophic AOB over AOA, which were unaffected by the treatment. The generality of these patterns in a globally coordinated nutrient addition experiment allows, as far as we know for the first time, to draw conclusions that consistent, guild-level responses occur worldwide despite substantial differences in plant communities, climatic conditions, and soil properties across grasslands. Long-term N-fertilization at the dose applied in this study may have considerable ecological implications for the abundance of functionally important microbial groups and effects on nutrient cycling in general. More abundant AOB under N-addition may accelerate nitrification activity in grasslands, which has the potential to increase NO₃⁻ in soils, thereby leading to higher environmental pollution in the form of NO₃⁻ leaching or N₂O emission. Long-term N-addition is also likely to change above- and belowground grassland biodiversity (Leff et al., 2015), due to soil acidification and preferential uptake of NO₃⁻, and could therefore threaten the sustainability of grassland functioning.

Author contributions

AF and BF designed the study, all authors contributed to the soil sampling and collection of data. AR, BF, BM and BT analyzed the data, AR and BF drafted the manuscript to which all authors contributed.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

This work was conducted within the Nutrient Network (<http://www.nutnet.org>) experiment, which is funded at the site scale by individual researchers. The N-cycling microbial communities add-on study was funded by an internal competitive WSL grant (PSP 5233.00176.001.01) to B.F. and A.C.R. Coordination and data management have been supported by funding from the National Science Foundation Research Coordination Network (NSF-DEB-1042132) to E.T.B. and E.W.S., the Long-Term Ecological Research (LTER) programme (NSF-DEB-1234162), and the Institute on the Environment at the University of Minnesota (DG-0001-13). We also thank the Minnesota Supercomputer Institute for hosting project data, and the Institute on the Environment for hosting NutNet meetings. We are grateful to B. Stierli, S. Baumgartner and A. Dharmarajah for their help with sample processing and analyses. We also thank M. Dawes for her valuable contribution to the editing of this article.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.soilbio.2022.108887>.

References

Aigle, A., Prosser, I.J., Gubry-Rangin, C., 2019. The application of high-throughput sequencing technology to analysis of amoA phylogeny and environmental niche

- specialisation of terrestrial bacterial ammonia-oxidisers. *Environmental Microbiome* 14, 3.
- Alves, R.J., Wanek, W., Zappe, A., Richter, A., Svenning, M.M., Schlexer, C., Ulrich, T., 2013. Nitrification rates in Arctic soils are associated with functionally distinct populations of ammonia-oxidizing archaea. *The ISME Journal* 7, 1620–1631.
- Alves, R.J., Minh, B.Q., Ulrich, T., von Haeseler, A., Schlexer, C., 2018. Unifying the global phylogeny and environmental distribution of ammonia-oxidising archaea based on *amoA* genes. *Nature Communications* 9, 1–17.
- Banning, N.C., Maccarone, L.D., Fisk, L.M., Murphy, D.V., 2015. Ammonia-oxidising bacteria not archaea dominate nitrification activity in semi-arid agricultural soil. *Scientific Reports* 5, 11146.
- Borer, E.T., Harpole, W.S., Adler, P.B., Lind, E.M., Orrock, J.L., Seabloom, E.W., Smith, M.D., 2014. Finding generality in ecology: a model for globally distributed experiments. *Methods of Ecological Evolution* 5, 65–73.
- Burton, S.A.Q., Prosser, J.I., 2001. Autotrophic ammonia oxidation at low pH through urea hydrolysis. *Applied and Environmental Microbiology* 67, 2952–2957.
- Cai, Z., Wang, B., Xu, M., Zhang, H., Zhang, L., Gao, S., 2014. Nitrification and acidification from urea application in red soil (Ferralic Cambisol) after different long-term fertilization treatments. *Journal of Soils and Sediments* 14, 1526–1536.
- Cardenas, L.M., Bhogal, A., Chadwick, D.R., McGeough, K., Misselbrook, T., Rees, R.M., Thorman, R.E., Watson, C.J., Williams, J.R., Smith, K.A., Calvet, S., 2019. Nitrogen use efficiency and nitrous oxide emissions from five UK fertilised grasslands. *Science of the Total Environment* 661, 696–710.
- Carey, C.J., Dove, N.C., Beman, J.M., Hart, S.C., Aronson, E.L., 2016. Meta-analysis reveals ammonia-oxidizing bacteria respond more strongly to nitrogen addition than ammonia-oxidizing archaea. *Soil Biology and Biochemistry* 99, 158–166.
- Carrell, A.A., Kolton, M., Glass, J.B., Pelletier, D.A., Warren, M.J., Kostka, J.E., Iversen, C.M., Hanson, P.J., Weston, D.J., 2019. Experimental warming alters the community composition, diversity, and N₂ fixation activity of peat moss (*Sphagnum fallax*) microbiomes. *Global Change Biology* 25, 2993–3004.
- Che, R., Deng, Y., Wang, F., Wang, W., Xu, Z., Hao, Y., Xue, K., Zhang, B., Tang, L., Zhou, H., Cui, X., 2018. Autotrophic and symbiotic diazotrophs dominate nitrogen-fixing communities in Tibetan grassland soils. *Science of the Total Environment* 639, 997–1006.
- Chen, Y., Xu, Z., Hu, H., Hu, Y., Hao, Z., Jiang, Y., Chen, B., 2013. Responses of ammonia-oxidizing bacteria and archaea to nitrogen fertilization and precipitation increment in a typical temperate steppe in Inner Mongolia. *Applied Soil Ecology* 68, 36–45.
- Clarke, K.R., Gorley, R.N., 2015. PRIMER v7: user manual/tutorial. PRIMER-E Plymouth.
- Collavino, M.M., Tripp, H.J., Frank, I.E., Vidoz, M.L., Calderoli, P.A., Donato, M., Zehr, J.P., Aguilari, O.M., 2014. *nifH* pyrosequencing reveals the potential for location-specific soil chemistry to influence N₂-fixing community dynamics. *Environmental Microbiology* 16, 3211–3223.
- Daims, H., Lebedeva, E.V., Pjevac, P., Han, P., Herbold, C., Albertsen, M., Jehmlich, N., Palatinszky, M., Vierheilig, J., Bulaev, A., Kirkegaard, R.H., von Bergen, M., Rattai, T., Bendinger, B., Nielsen, P.H., Wagner, M., 2015. Complete nitrification by *Nitrospira* bacteria. *Nature* 528, 504–509.
- Di, H.J., Cameron, K.C., Shen, J.P., Winefield, C.S., O'Callaghan, M., Bowatte, S., He, J.Z., 2009. Nitrification driven by bacteria and not archaea in nitrogen-rich grassland soils. *Nature Geoscience* 2, 621–624.
- Di, H.J., Cameron, K.C., 2016. Inhibition of nitrification to mitigate nitrate leaching and nitrous oxide emissions in grazed grassland: a review. *Journal of Soils and Sediments* 16, 1401–1420.
- Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., García Marquez, J.R., Gruber, B., Lafourcade, B., Leitao, P.J., et al., 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36, 27–46.
- Du, Y., Ke, X., Li, J., Wang, Y., Cao, G., Guo, X., Chen, K., 2021. Nitrogen deposition increases global grassland N₂O emission rates steeply: a meta-analysis. *Catena* 199, 105105.
- Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26, 2460–2461.
- Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., Knight, R., 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27, 2194–2200.
- Fan, K., Delgado-Baquerizo, M., Guo, X., Wang, D., Chu, H., 2019. Suppressed N fixation and diazotrophs after four decades of fertilization. *Microbiome* 7, 143.
- Fay, P., Prober, S., Harpole, W., Knops, J.M.H., Bakker, J.D., Borer, E.T., Lind, E.M., MacDougall, A.S., Seabloom, E.W., Wrapp, P.D., et al., 2015. Grassland productivity limited by multiple nutrients. *Nature Plants* 1, 15080.
- Feng, M., Adams, J.M., Fan, K., Shi, Y., Sun, R., Wang, W., Guo, X., Chu, H., 2018. Long-term fertilization influences community assembly processes of soil diazotrophs. *Soil Biology and Biochemistry* 126, 151–158.
- Fierer, N., 2017. Embracing the unknown: disentangling the complexities of the soil microbiome. *Nature Reviews Microbiology* 15, 579–590.
- Francis, C.A., Roberts, K.J., Beman, J.M., Santoro, A.E., Oakley, B.B., 2005. Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proceedings of the National Academy of Sciences of the United States of America* 102, 14683–14688.
- Frey, B., Niklaus, P.A., Kremer, J., Lüscher, P., Zimmermann, S., 2011. Heavy-machinery traffic impacts methane emissions as well as methanogen abundance and community structure in oxic forest soils. *Applied and Environmental Microbiology* 77, 6060–6068.
- Frey, B., Carnol, M., Dharmarajah, A., Brunner, I., Schleppl, P., 2020. Only minor changes in the soil microbiome of a sub-alpine forest after 20 years of moderately increased nitrogen loads. *Frontiers in Forests and Global Change* 3, 77.
- Frey, B., Walthert, L., Perez-Mon, C., Stierli, B., Koechli, R., Dharmarajah, A., Brunner, I., 2021. Deep soil layers of drought-exposed forests harbor poorly known bacterial and fungal communities. *Frontiers in Microbiology* 12, 674160.
- Gaby, J.C., Buckley, D.H., 2011. A global census of nitrogenase diversity. *Environmental Microbiology* 13, 1790–1799.
- Gaby, J.C., Buckley, D.H., 2012. A comprehensive evaluation of PCR primers to amplify the *nifH* gene of nitrogenase. *PLoS One* 7, e42149.
- Gee, G., Bauder, J., 1986. *Texture, Hydrometer Method*. Soil Science Society of America, Madison, WI, USA.
- Gubry-Rangin, C., Hai, B., Quince, C., Engel, M., Thomson, B.C., James, P., Schloter, M., Griffiths, R.I., Prosser, J.I., Nicol, G.W., 2011. Niche specialization of terrestrial archaeal ammonia oxidizers. *Proceedings of the National Academy of Sciences of the United States of America* 108, 21206–21211.
- Guerra, C.A., Heintz-Buschart, A., Sikorski, J., Chatzinotas, A., Guerrero-Ramirez, N., Cesarz, S., Beaumelle, L., Rillig, M.C., Maestre, F.T., Delgado-Baquerizo, M., et al., 2020. Blind spots in global soil biodiversity and ecosystem function research. *Nature Communications* 11, 3870.
- Han, L.L., Wang, Q., Shen, J.P., Di, H.J., Wang, J.T., Wei, W.X., Fang, Y.T., Zhang, L.M., He, J.Z., 2019. Multiple factors drive the abundance and diversity of the diazotrophic community in typical farmland soils of China. *FEMS Microbiology Ecology* 95, f12113.
- Hatzenpichler, R., 2012. Diversity, physiology, and niche differentiation of ammonia-oxidizing archaea. *Applied and Environmental Microbiology* 78, 7501–7510.
- He, J.Z., Shen, J.P., Zhang, L.M., Zhu, Y.G., Zheng, Y.M., Xu, M.G., Di, H.J., 2007. Quantitative analyses of the abundance and composition of ammonia-oxidizing bacteria and ammonia-oxidizing archaea of a Chinese upland red soil under long-term fertilization practices. *Environmental Microbiology* 9, 2364–2374.
- Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, G., Jarvis, A., 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25, 1965–1978.
- Hink, L., Nicol, G.W., Prosser, J.I., 2017. Archaea produce lower yields of N₂O than bacteria during aerobic ammonia oxidation in soil. *Environmental Microbiology* 19, 4829–4837.
- Hsu, S.-F., Buckley, D.H., 2009. Evidence for the functional significance of diazotroph community structure in soil. *The ISME Journal* 3, 124–136.
- Hu, J., Richwine, J.D., Keyser, P.D., Li, L., Yao, F., Jagadamma, S., DeBruyn, M., 2021. Nitrogen fertilization and native C4 grass species alter abundance, activity, and diversity of soil diazotrophic communities. *Frontiers in Microbiology* 12, 675693.
- Lauber, C.L., Hamady, M., Knight, R., Fierer, N., 2009. Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Applied and Environmental Microbiology* 75, 5111–5120.
- Lebauer, D.S., Treseder, K.K., 2008. Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed. *Ecology* 89, 371–379.
- Lekberg, Y., Arnillas, C.A., Borer, E.T., Bullington, L.S., Fierer, N., Kennedy, P.G., Leff, J.W., Luis, A.D., Seabloom, E.W., Henning, J.A., 2021. Nitrogen and phosphorus fertilization consistently favor pathogenic over mutualistic fungi in grasslands soils. *Nature Communications* 12, 3484.
- Leff, J.W., Jones, S.E., Prober, S.M., Barberán, A., Borer, E.T., Firn, J.L., Harpole, W.S., Hobbie, S.E., Hofmockel, K.S., Knops, J.M.H., et al., 2015. Consistent responses of soil microbial communities to elevated nutrient inputs in grasslands across the globe. *Proceedings of the National Academy of Sciences of the United States of America* 112, 10967–10972.
- Lehtovirta-Morley, L.E., 2018. Ammonia oxidation: ecology, physiology, biochemistry and why they must all come together. *FEMS Microbiology Letters* 365, fny058.
- Levy-Booth, D.J., Prescott, C.E., Grayston, S.J., 2014. Microbial functional genes involved in nitrogen fixation, nitrification and denitrification in forest ecosystems. *Soil Biology and Biochemistry* 75, 11–25.
- Lieth, H., 1978. *Patterns of Primary Productivity in the Biosphere*. Hutchinson & Ross.
- Liu, J., Yu, Z., Yao, Q., Sui, Y., Shi, Y., Chu, H., Tang, C., Franks, A.E., Jin, J., Liu, X., Wang, G., 2018. Ammonia-oxidizing archaea show more distinct biogeographic distribution patterns than ammonia-oxidizing bacteria across the black soil zone of Northeast China. *Frontiers in Microbiology* 9, 171.
- Martens-Habbena, W., Berube, P.M., Urakawa, H., José, R., Stahl, D.A., 2009. Ammonia oxidation kinetics determine niche separation of nitrifying archaea and bacteria. *Nature* 461, 976–979.
- Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17, 1.
- Nepel, M., Angel, R., Borer, E., Frey, B., MacDougall, A., McCulley, R., Risch, A., Schütz, M., Seabloom, E., Woebken, D., 2022. Global grassland diazotrophic communities are structured by combined abiotic, biotic, and spatial distance factors but resilient to fertilization. *Frontiers in Microbiology* 13, 821030.
- Nicol, G.W., Leininger, S., Schlexer, C., Prosser, J.I., 2008. The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. *Environmental Microbiology* 10, 2966–2978.
- Ning, Q., Gu, Q., Shen, J., Lv, X., Yang, J., Zhang, X., He, J., Huang, J., Wang, H., Xu, Z., Han, X., 2015. Effects of nitrogen deposition rates and frequencies on the abundance of soil nitrogen-related functional genes in temperate grassland of northern China. *Journal of Soils and Sediments* 15, 694–704.
- Patra, A.K., Abbadi, L., Clays-Josserand, A., Degrange, V., Grayston, S.J., Guillaumaud, N., Loiseau, P., Louault, F., Mahmood, S., Nazaret, S., et al., 2006. Effects of management regime and plant species on the enzyme activity and genetic structure of N-fixing, denitrifying and nitrifying bacterial communities in grassland soils. *Environmental Microbiology* 8, 1005–1016.
- Penton, C.R., Yang, C., Wu, L., Wang, Q., Zhang, J., Liu, F., Qin, Y., Deng, Y., Hemme, C.L., Zheng, T., Schuur, E.A., Tiedje, J., Zhou, J., 2016. *NifH*-harboring bacterial

- community composition across an Alaskan permafrost thaw gradient. *Frontiers in Microbiology* 7, 1894.
- Pester, M., Rattei, T., Flechl, S., Gröngroft, A., Richter, A., Overmann, J., Reinhold-Hurek, B., Loy, A., Wagner, M., 2012. *amoA*-based consensus phylogeny of ammonia-oxidizing archaea and deep sequencing of *amoA* genes from soils of four different geographic regions. *Environmental Microbiology* 14, 525–539.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team, 2021. *Nlme: linear and nonlinear mixed effects models*. R package version 3, 1–152. <https://CRAN.R-project.org/package=nlme>.
- Poly, F., Monrozier, L.J., Bally, R., 2001. Improvement in the RFLP procedure for studying the diversity of *nifH* genes in communities of nitrogen fixers in soil. *Research in Microbiology* 152, 95–103.
- Prober, S.M., Leff, J.W., Bates, S.T., Borer, E.T., Firn, J., Harpole, W.S., Lind, E.M., Seabloom, E.W., Adler, P.B., Bakker, J.D., et al., 2015. Plant diversity predicts beta but not alpha diversity of soil microbes across grasslands worldwide. *Ecological Letters* 18, 85–95.
- Prosser, J.I., Nicol, G.W., 2012. Archaeal and bacterial ammonia-oxidisers in soil: the quest for niche specialisation and differentiation. *Trends in Microbiology* 20, 523–531.
- Rafique, G., Hennessy, D., Kiely, G., 2011. Nitrous oxide emission from grazed grassland under different management systems. *Ecosystems* 14, 563–582.
- R Core Team, 2020. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- Reardon, C.L., Gollany, H.T., Wuest, S.B., 2014. Diazotroph community structure and abundance in wheat-fallow and wheat-pea crop rotations. *Soil Biology and Biochemistry* 69, 406–412.
- Reed, S.C., Cleveland, C.C., Townsend, A.R., 2011. Functional ecology of free living nitrogen fixation: a contemporary perspective. *Annual Review of Ecology, Evolution and Systematics* 42, 489–512.
- Regan, K., Stempfhuber, B., Schloter, M., Rasche, F., Prati, D., Philippot, L., Boeddinghaus, R.S., Kandeler, E., Marhan, S., 2017. Spatial and temporal dynamics of nitrogen fixing, nitrifying and denitrifying microbes in an unfertilized grassland soil. *Soil Biology and Biochemistry* 109, 214–226.
- Risch, A.C., Zimmermann, S., Ochoa-Hueso, R., Schütz, M., Frey, B., Firn, J.L., Fay, P.A., Hagedorn, F., Borer, E.T., Seabloom, E.W., et al., 2019. Soil net nitrogen mineralisation across grasslands. *Nature Communications* 10, 1–10.
- Risch, A.C., Zimmermann, S., Moser, B., Schütz, M., Hagedorn, F., Firn, J., Fay, P.A., Adler, P.B., Biederman, L.A., Blair, J.M., et al., 2020. Global impacts of fertilization and herbivore removal on soil net nitrogen mineralization are modulated by local climate and soil properties. *Global Change Biology* 26, 7173–7185.
- Rogiers, N., Conen, F., Furger, M., Stöckli, R., Eugster, W., 2010. Impact of past and present land-management on the C-balance of a grassland in Swiss Alps. *Global Change Biology* 14, 2613–2625.
- Rotthauwe, J.-H., Witzel, K.-P., Liesack, W., 1997. The ammonia monoxygenase structural gene *amoA* as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing populations. *Applied and Environmental Microbiology* 63, 4704–4712.
- Rui, J., Hu, J., Wang, F., Zhao, Y., Li, C., 2022. Altitudinal niches of symbiotic, associative and free-living diazotrophs driven by soil moisture and temperature in the alpine meadow on the Tibetan Plateau. *Environmental Research* 211, 113033.
- Shi, W., Zhao, H.Y., Chen, Y., Wang, J.S., Han, B., Li, C.P., Lu, J.Y., Zhang, L.M., 2021. Organic manure rather than phosphorus fertilization primarily determined symbiotic nitrogen fixation rate and the stability of diazotrophic community in an upland red soil. *Agriculture, Ecosystems & Environment* 319, 107535.
- Sterngren, A.E., Hallin, S., Bengtson, P., 2015. Archaeal ammonia oxidizers dominate in numbers, but bacteria drive gross nitrification in N-amended grassland soil. *Frontiers in Microbiology* 6, 1350.
- Tao, J., Bai, T., Xiao, R., Wang, P., Wang, F., Duryee, A.M., Wang, Y., Zhang, Y., Hu, S., 2018. Vertical distribution of ammonia-oxidizing microorganisms across a soil profile of the Chinese Loess Plateau and their responses to nitrogen inputs. *Science of the Total Environment* 635, 240–248.
- Tian, D., Niu, S., 2015. A global analysis of soil acidification caused by nitrogen addition. *Environmental Research Letters* 10, 024019.
- Tognetti, P.M., Prober, S., Baez, S., Chaneton, E.J., Firn, J., Risch, A.C., Schütz, M., Simonsen, A.K., Yahdjian, L., Borer, E.T., et al., 2021. Negative effects of nitrogen override positive effects of phosphorus on grassland legumes worldwide. *Proceedings of the National Academy of Sciences of the United States of America* 118, e2023718118.
- Tourna, M., Freitag, T.E., Nicol, G.W., Prosser, J.I., 2008. Growth, activity and temperature responses of ammonia-oxidizing archaea and bacteria in soil microcosms. *Environmental Microbiology* 10, 1357–1364.
- Trivedi, C., Reich, P.B., Maestre, F.T., Hu, H.W., Singh, B.K., Delgado-Baquerizo, M., 2019. Plant-driven niche differentiation of ammonia-oxidizing bacteria and archaea in global drylands. *The ISME Journal* 13, 2727–2736.
- Tu, Q., Deng, Y., Yan, Q., Shen, L., He, Z., Wu, L., Van Nostrand, J.D., Buzzard, V., Michaletz, S.T., Enquist, B.J., 2016. Biogeographic patterns of soil diazotrophic communities across six forests in the North America. *Molecular Ecology* 25, 2937–2948.
- Van Groenigen, J.W., Huygens, D., Boeckx, P., Kuyper, T.W., Lubbers, I.M., Rütting, T., Groffman, P.M., 2015. The soil N cycle: new insights and key challenges. *Soils* 1, 235–256.
- Vázquez, E., Teutschero, N., Pastorelli, R., Lagomarsino, A., Giagnoni, L., Renella, G., 2020. Liming reduces N₂O emissions from Mediterranean soil after-rewetting and affects the size, structure and transcription of microbial communities. *Soil Biology and Biochemistry* 147.
- Vázquez, E., Schleuss, P.M., Borer, E.T., Bugalho, M.N., Caldeira, M.C., Eisenhauer, N., Eskelinen, A., Fay, P.A., Haider, S., et al., 2022. Nitrogen but not phosphorus addition affects symbiotic N₂ fixation by legumes in natural and seminatural grasslands located on four continents. *Plant and Soil*. <https://doi.org/10.1007/s11104-022-05498-y> (in press).
- Verhamme, D.T., Prosser, J.I., Nicol, G.W., 2011. Ammonia concentration determines differential growth of ammonia-oxidising archaea and bacteria in soil microcosms. *The ISME Journal* 5, 1067–1071.
- Vitousek, P.M., Menge, D.N.L., Reed, S.C., Cleveland, C.C., 2013. Biological nitrogen fixation: rates, patterns and ecological controls in terrestrial ecosystems. *Philosophical Transactions of the Royal Society Serie B Biological Sciences* 368, 20130119.
- Wang, C., Zheng, M., Song, W., Wen, S., Wang, B., Zhu, C., Shen, R., 2017a. Impact of 25 years of inorganic fertilization on diazotrophic abundance and community structure in an acidic soil in southern China. *Soil Biology and Biochemistry* 113, 240–249.
- Wang, Y., Li, C., Kou, Y., Wang, J., Tu, B., Li, H., Li, X., Wang, C., Yao, M., 2017b. Soil pH is a major driver of soil diazotrophic community assembly in Qinghai-Tibet alpine meadows. *Soil Biology and Biochemistry* 115, 547–555.
- Wang, F., Li, Z., Wei, Y., Su, F., Guo, H., Guo, J., Wang, Y., Zhang, Y., Hu, S., 2021. Responses of soil ammonia-oxidizing bacteria and archaea to short-term warming and nitrogen input in a semi-arid grassland on the Loess Plateau. *European Journal of Soil Biology* 102, 1164–5563.
- Xiang, X., He, D., He, J.S., Myrold, D., Chu, H., 2017. Ammonia-oxidizing bacteria rather than archaea respond to short-term urea amendment in an alpine grassland. *Soil Biology and Biochemistry* 107, 218–225.
- Yang, Y.D., Ren, Y.F., Wang, X.Q., Hu, Y.G., Wang, Z.M., Zeng, Z.H., 2018. Ammonia-oxidizing archaea and bacteria responding differently to fertilizer type and irrigation frequency as revealed by Illumina Miseq sequencing. *Journal of Soils and Sediments* 18, 1029–1040.
- Ying, J., Li, X., Wang, N., Lan, Z., He, J., Bai, Y., 2017. Contrasting effects of nitrogen forms and soil pH on ammonia oxidizing microorganisms and their responses to long-term nitrogen fertilization in a typical steppe ecosystem. *Soil Biology and Biochemistry* 107, 10–18.
- Zhang, X., Liu, W., Schloter, M., Zhang, G., Chen, Q., Huang, J., Li, L., Elser, J.J., Han, X., 2013. Response of the abundance of key soil microbial nitrogen-cycling genes to multi-factorial global changes. *PLoS One* 8 (10), e76500.
- Zhang, J., Kobert, K., Flouri, T., Stamatakis, A., 2014. PEAR: a fast and accurate Illumina paired-end reAd mergeR. *Bioinformatics* 30, 614–620.
- Zhang, G., Huang, J., Jia, M., Liu, F., Yang, Y., Wang, Z., Han, G., 2019. Ammonia-oxidizing bacteria and archaea: response to simulated climate warming and nitrogen supplementation. *Soil Science Society of America Journal* 83, 1683–1695.
- Zhou, Z.F., Shi, X.J., Zheng, Y., Qin, Z.X., Xie, D.T., Li, Z.L., Guo, T., 2014. Abundance and community composition of ammonia-oxidizing bacteria and archaea in purple soil under long-term fertilization. *European Journal of Soil Biology* 60, 24–33.
- Zhou, J., Ma, M., Guan, D., Jiang, X., Zhang, N., Shu, F., Kong, Y., Li, J., 2021. Nitrogen has a greater influence than phosphorus on the diazotrophic community in two successive crop seasons in Northeast China. *Scientific Reports* 11, 6303.
- Zhu, C., Tian, G., Luo, G., Kong, Y., Guo, J., Wang, M., Guo, S., Ling, N., Shen, Q., 2018. N-fertilizer-driven association between the arbuscular mycorrhizal fungal community and diazotrophic community impacts wheat yield. *Agriculture, Ecosystems & Environment* 254, 191–201.