




## Gender differences in respiratory health outcomes among farming cohorts around the globe: findings from the AGRICOH consortium

Jonathan Fix, Isabella Annesi-Maesano, Isabelle Baldi, Mathilde Boulanger, Soo Cheng, Sandra Cortes, Jean- Charles Dalphin, Mohamed Aqiel Dalvie, Bruno Degano, Jeroen Douwes, Wijnand Eduard, Grethe Elholm, Catterina Ferreccio, Anne-Helen Harding, Mohamed Jeebhay, Kevin M Kelly, Hans Kromhout, Ewan MacFarlane, Cara Nichole Maesano, Diane Catherine Mitchell, Hussein Mwanga, Saloshni Naidoo, Beyene Negatu, Dorothy Ngajilo, Karl- Christian Nordby, Christine G Parks, Marc B Schenker, Aesun Shin, Torben Sigsgaard, Malcolm Sim, Thibaud Soumagne, Peter Thorne, Keun-Young Yoo & Jane A Hoppin


**To cite this article:** Jonathan Fix, Isabella Annesi-Maesano, Isabelle Baldi, Mathilde Boulanger, Soo Cheng, Sandra Cortes, Jean- Charles Dalphin, Mohamed Aqiel Dalvie, Bruno Degano, Jeroen Douwes, Wijnand Eduard, Grethe Elholm, Catterina Ferreccio, Anne-Helen Harding, Mohamed Jeebhay, Kevin M Kelly, Hans Kromhout, Ewan MacFarlane, Cara Nichole Maesano, Diane Catherine Mitchell, Hussein Mwanga, Saloshni Naidoo, Beyene Negatu, Dorothy Ngajilo, Karl- Christian Nordby, Christine G Parks, Marc B Schenker, Aesun Shin, Torben Sigsgaard, Malcolm Sim, Thibaud Soumagne, Peter Thorne, Keun-Young Yoo & Jane A Hoppin (2021) Gender differences in respiratory health outcomes among farming cohorts around the globe: findings from the AGRICOH consortium, *Journal of Agromedicine*, 26:2, 97-108, DOI: [10.1080/1059924X.2020.1713274](https://doi.org/10.1080/1059924X.2020.1713274)

**To link to this article:** <https://doi.org/10.1080/1059924X.2020.1713274>

 [View supplementary material](#) 

 Published online: 17 Mar 2020.

 [Submit your article to this journal](#) 

 Article views: 622

 [View related articles](#) 



View Crossmark data [↗](#)

---



Citing articles: 9 View citing articles [↗](#)

---



## Gender differences in respiratory health outcomes among farming cohorts around the globe: findings from the AGRICOH consortium

Jonathan Fix<sup>a</sup>, Isabella Annesi-Maesano <sup>b</sup>, Isabelle Baldi<sup>c,d</sup>, Mathilde Boulanger<sup>c,d</sup>, Soo Cheng<sup>e</sup>, Sandra Cortes<sup>f,g</sup>, Jean-Charles Dalphin<sup>\*h,i</sup>, Mohamed Aqiel Dalvie<sup>j</sup>, Bruno Degano<sup>k,l</sup>, Jeroen Douwes<sup>e</sup>, Wijnand Eduard<sup>m</sup>, Grethe Elholm<sup>n</sup>, Catterina Ferreccio<sup>f,g</sup>, Anne-Helen Harding<sup>o</sup>, Mohamed Jeebhay <sup>j</sup>, Kevin M Kelly<sup>p</sup>, Hans Kromhout<sup>q</sup>, Ewan MacFarlane<sup>r</sup>, Cara Nichole Maesano<sup>b</sup>, Diane Catherine Mitchell<sup>s</sup>, Hussein Mwangi<sup>j</sup>, Saloshni Naidoo <sup>t</sup>, Beyene Negatu<sup>q,u</sup>, Dorothy Ngajilo <sup>j</sup>, Karl-Christian Nordby<sup>v</sup>, Christine G Parks<sup>w</sup>, Marc B Schenker <sup>s</sup>, Aesun Shin<sup>x</sup>, Torben Sigsgaard <sup>n</sup>, Malcolm Sim<sup>f</sup>, Thibaud Soumagne<sup>h,i</sup>, Peter Thorne<sup>p</sup>, Keun-Young Yoo<sup>x</sup>, and Jane A Hoppin <sup>y</sup>

<sup>a</sup>Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, NC, USA; <sup>b</sup>Epidemiology of Allergic and Respiratory Diseases Department (EPAR), Pierre Louis Institute of Epidemiology and Public Health (IPLESP UMR1136), UPMC Univ Paris 06, INSERM, Saint-Antoine Medical School, Sorbonne Universités, Paris; <sup>c</sup>INSERM U1086, Caen, France; <sup>d</sup>Centre de Lutte contre le Cancer François Baclesse, Caen, France; <sup>e</sup>Centre for Public Health Research, Massey University, PO Box 756, Wellington 6021, New Zealand; <sup>f</sup>Advanced Center for Chronic Diseases (ACCDiS), Facultad Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile; <sup>g</sup>Escuela de Medicina, Pontificia Universidad Católica de Chile, Santiago, 8330077, Chile; <sup>h</sup>Service de Pneumologie, CHU de Besançon, Besançon, France; <sup>i</sup>UMR 6249 Chrono-environnement, Université de Franche-Comté, Besançon, France; <sup>j</sup>Division of Occupational Medicine and Centre for Environmental and Occupational Health Research, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa; <sup>k</sup>Research, School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa Universitaire (CHRU), Besançon, France; <sup>l</sup>EA 3920, Université de Franche-Comté, Besançon, France.; <sup>m</sup>Department of Chemical and Biological Work Environment, STAMI National Institute of Occupational Health, PO Box 5330 Majorstuen, NO-0304 Oslo, Norway; <sup>n</sup>Department of Public Health, Danish Ramazzini Centre, Aarhus University, Aarhus, Denmark; <sup>o</sup>Health and Safety Executive, Harpur Hill, Buxton SK17 9JN, UK; <sup>p</sup>Department of Occupational and Environmental Health, University of Iowa, Iowa City, IA 52242, United States; <sup>q</sup>Division of Environmental Epidemiology, Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands; <sup>r</sup>Monash Centre for Occupational and Environmental Health (MonCOEH), Department of Epidemiology and Preventive Medicine, Faculty of Medicine, Nursing and Health Sciences, Monash University, Melbourne, VIC, Australia; <sup>s</sup>Department of Public Health Sciences, University of California, Davis, Davis, California, USA; <sup>t</sup>Discipline of Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal, South Africa; <sup>u</sup>Department of Pesticide Registration and control, Ministry of Agriculture and Livestock development, Addis Ababa, Ethiopia; <sup>v</sup>National Institute of Occupational Health, Oslo, Norway; <sup>w</sup>Epidemiology Branch, National Institute of Environmental Health Sciences, National Institutes of Health (NIH), Department of Health and Human Services (DHHS), Research Triangle Park, North Carolina, USA; <sup>x</sup>Department of Preventive Medicine, Seoul National University College of Medicine, 103 Daehakro, Chongno-gu, Seoul 110-799, Korea.; <sup>y</sup>Department of Biological Sciences, Center for Human Health and the Environment, North Carolina State University, NC, USA

### ABSTRACT

**Purpose:** Respiratory hazards of farming have been identified for centuries, with little focus on gender differences. We used data from the AGRICOH consortium, a collective of prospective cohorts of agricultural workers, to assess respiratory disease prevalence among adults in 18 cohorts representing over 200,000 farmers, farm workers, and their spouses from six continents.



**Methods:** Cohorts collected data between 1992 and 2016 and ranged in size from 200 to >128,000 individuals; 44% of participants were female. Farming practices varied from subsistence farming to large-scale industrial agriculture. All cohorts provided respiratory outcome information for their cohort based on their study definitions. The majority of outcomes were based on self-report using standard respiratory questionnaires; the greatest variability in assessment methods was associated with chronic obstructive pulmonary disease (COPD).

**Results:** For all three respiratory symptoms (cough, phlegm, and wheeze), the median prevalence in men was higher than in women, with the greatest difference for phlegm (17% vs. 10%). For asthma, women had a higher prevalence (7.8% vs 6.5%), with the difference associated with allergic asthma. The relative proportion of allergic asthma varied among cohorts. In two of eight cohorts for women and two of seven cohorts for men, allergic asthma was more common than non-allergic asthma.


**Conclusions:** These findings indicate that respiratory outcomes are common among farmers around the world despite differences in agricultural production. As women in the general population are at higher risk of asthma, exploring gender differences in occupational studies is critical for a deeper understanding of respiratory disease among agricultural workers.

### KEYWORDS

Farmers; Respiratory Health; Gender; Farmworkers; Occupational Exposure

**CONTACT** Jane A Hoppin  [jahoppin@ncsu.edu](mailto:jahoppin@ncsu.edu)  Biological Sciences, Center for Human Health and the Environment Raleigh, North Carolina State University, Raleigh, NC 27695, USA

\*deceased

 Supplemental data for this article can be accessed [here](#).

© 2020 Informa UK Limited, trading as Taylor & Francis Group

## Introduction

Agricultural work is an occupation represented by both men and women, though little formal comparison of respiratory disease prevalence by gender has been conducted. Work in agriculture has been associated with respiratory diseases as early as the mid-1500s.<sup>1</sup> Even as farm work becomes more industrialized, there continues to be evidence of increased respiratory risk for agricultural workers. Factors contributing to increased risk of respiratory diseases and symptoms include frequent exposure to dusts, microorganisms, toxic gases (e.g. diesel motor exhaust, welding fumes, ammonia, hydrogen sulfide, and carbon dioxide), and pesticides.<sup>2</sup> Agricultural activities vary by region and thus, exposures will also vary. While respiratory diseases have been a concern among farming populations for many centuries, recent studies have focused on differences between allergic vs. non-allergic phenotypes<sup>3-7</sup> due to the reduced risk of allergy associated with growing up on a farm.<sup>8</sup>

Agricultural workers include both farmers who are owners and operators and farm workers (local and migrant). Women often have an active role in farm production activities, but in some populations are not regarded as farmers, and as a result, the impact of their occupational exposures may be under-represented. Agricultural work remains an occupation that people often begin at young ages.

While many papers have been published related to the respiratory health of agricultural workers, few, if any, have attempted to integrate data from across continents and farming practices to characterize the global respiratory health implications of agricultural work. In 1998, the American Thoracic Society published a research statement on Respiratory Hazards of Agriculture.<sup>1</sup> In this seminal work, extensive detail was provided on the specific respiratory outcomes, as well as the range of exposures experienced by agricultural workers primarily in developed countries. However, respiratory hazards of women and agricultural work in low- and middle-income countries were not well represented. A limited amount of literature has been published assessing gender differences in the incidence and/or prevalence of respiratory symptoms and diseases among farming populations, and nearly all of these studies focus on populations in either the United States or Europe.<sup>9-11</sup> Although there has not been

a multi-country study to estimate the global burden of respiratory disease among farmers, the numerous cohort studies that have been conducted around the world can help start to address this gap in knowledge.

The AGRICOH consortium was created in 2006 as a collaborative effort to assess relationships between farming exposures and health outcomes. Currently, 29 cohorts are participating and they include a diverse range of types of farming from all continents.<sup>12</sup> To assess the prevalence of respiratory disease and symptoms among diverse farming populations and to explore differences between men and women, we focused on common respiratory endpoints from AGRICOH cohorts with respiratory outcome information.

## Methods

### Study population

The AGRICOH consortium consists of 29 cohorts from around the world (<http://agricoh.iarc.fr/>). We focused on adults for this analysis because we were interested in occupational exposures. Fourteen cohorts were excluded from this analysis: four did not include adults, eight did not collect respiratory outcome information, and two were unable to provide data. All eligible AGRICOH cohorts agreed to participate. Additionally, three farming cohorts outside of AGRICOH, but which included AGRICOH investigators, were also identified for inclusion, resulting in 18 cohorts for this analysis.<sup>13-29</sup> These cohorts cumulatively represent 211,232 people. The data presented include farming populations around the world, including those of low- and middle-income countries with different socioeconomic settings, spanning from 1992 to 2016.

### Data collection

For this project, we developed a structured reporting form to collect standardized demographic and outcome information (see supplement 1). All cohorts were asked to provide summary statistics on their cohorts; raw data were not collected. All cohorts except the Agricultural Health Study (AHS) provided prevalence estimates for data collected at enrollment. For the AHS, data from the most recent AHS

interview were included because this information was more complete than the enrollment information.

### **Cohort demographic information**

The data collection form requested information on cohort: sample size and calendar year of data collection, as well as summary statistics regarding age distribution, gender, smoking status (current/past/never), and body mass index (BMI, <20, 20 - <25, 25 - <30, and ≥30). We also collected data on the participant types (farmer, farm worker, subsistence farmer), role of women in cohort (full-time farmers, help on farm, spouses), type of livestock raised (poultry, beef cattle, dairy cattle, pork, other), type of farming (row crops, vegetables, orchards, other), and grain handling. We applied the World Bank Atlas method to determine country economy classifications, categorized by gross national income (GNI) per capita: low-income (≤\$1,025), lower-middle-income (\$1,026-\$4,035), upper-middle-income (\$4,036–12,475), and high income (≥\$12,476).<sup>30</sup>

### **Respiratory outcome information**

Cohorts were asked to provide prevalence and 95% confidence interval information for three respiratory symptoms (cough, phlegm, and wheeze) and two respiratory diseases (asthma and chronic obstructive pulmonary disease (COPD)). Asthma was further classified as either allergic or non-allergic, based on each cohort's definition.

Outcome definitions varied by cohort with most using standard respiratory questionnaires (e.g., American Thoracic Society Questionnaire<sup>31</sup> and European Community Respiratory Health Survey<sup>32</sup>). Some studies collected only self-reported outcome information while others incorporated clinical measurements as well. All symptom information was self-reported. Asthma was reported as either ever asthma or asthma in the past 12 months. All but one study relied on self-reported doctor's diagnosis of asthma, while the remaining study included a broader definition of having had a history in the past year of an attack of shortness of breath, an asthma attack, using asthma medication or having a positive bronchodilator test conducted by study staff.<sup>16</sup>

COPD definition was most variable among the cohorts (supplement 1). Fifteen cohorts provided information on COPD prevalence. This information was based on self-report (67%, 10/15 cohorts), spirometry (27%, 4/15 cohorts), or an inclusive definition of either spirometry or self-report (7%, 1/15 cohorts). Among the 11 studies that used self-report to ascertain COPD status, definitions for COPD included: doctor diagnosis of COPD alone (27%); doctor diagnosis of chronic bronchitis alone (45%); the combination of doctor diagnosis of COPD, chronic bronchitis, or emphysema (18%); or self-reported history of COPD alone (9%).

Among seven cohorts reporting both allergic and non-allergic asthma, definitions for allergic asthma were asthma with hay fever (29%), asthma with hay fever and/or eczema (14%), asthma with rhinitis and other allergic diseases (14%), asthma with positive Phadiatop test (14%), asthma with atopy or fractional exhaled nitric oxide (FeNO) >50 ppb (14%), or asthma with three or more positive responses to a skin-prick test (14%).

### **Statistical analysis**

Our statistical analysis focused on descriptive statistics of health outcomes. We integrated the summary statistics from all cohorts to create summary tables. To describe the distribution of respiratory outcomes among the cohorts, we calculated the median, interquartile range (IQR), and total range for prevalence of respiratory symptoms and diseases, for cohorts as a whole and then by gender. Reported medians were not weighted by cohort size, and reflect the 50<sup>th</sup> percentile of individual cohort prevalence estimates. To assess the relative prevalence of allergic to non-allergic asthma, gender-stratified prevalence ratios and corresponding 95% confidence intervals were calculated for each cohort. Analysis was completed in SAS 9.4 and figures developed in R.

## **Results**

### **Cohort characteristics**

A total of 18 cohorts from 11 countries participated, including 118,520 men and 92,712 women (N = 211,232) from both crop and livestock farming

populations. Cohorts had a wide geographic spread, including populations from six continents (Figure 1). Participants include farmers, farm workers, subsistence farmers, and rural residents. Agricultural activities range from large-scale industrial agriculture to rural fruit and farmworkers in Africa (supplement 2). The majority of the cohorts were based in high-income countries (14/18, 78%), while three (17%) were based in upper-middle-income, and one (6%) in a low-income country.

The cohorts ranged in size from 206 to 128,388 participants: six (33%) included fewer than 1,000 participants; 10 studies (56%) included 1,000–10,000 participants; and two (11%) included greater than 10,000 participants (Table 1). The majority of cohorts featured a greater proportion of males; however, two cohorts (KwaZulu-Natal Crop Farmers<sup>14</sup> and Western Cape Fruit Farm Workers<sup>15</sup>) included only female farm workers. The median proportion of males among cohorts was 60% (Interquartile Range (IQR): 44–78%). Among all cohorts, ages ranged from 15 to 105 years, with a median cohort-specific age of 51 years (IQR: 40.7–55.5).

### Covariates

We collected data on two common risk factors for respiratory outcomes: smoking and body mass

index (BMI). The prevalence of smoking varied among the cohorts. The prevalence of current smoking ranging between 5% and 50%, with a median of 15% (IQR: 9%–30%) (Table 1 and supplement 3). There were no apparent differences in smoking prevalence between industrialized and non-industrialized countries or by geographic region. Smoking varied by gender, with current smoking prevalence consistently higher among males (median = 17% [IQR: 9%–30%]) compared to females (median = 11% [IQR: 7%–22%]). For BMI, the proportion of obese individuals ranged from 1% to 40% (median = 17% [IQR: 13%–33%]), and overweight individuals ranged from 11% to 44% (median = 31% [IQR: 21%–44%]). Higher proportions of underweight participants were more common among KMCC<sup>17</sup> and African cohorts<sup>13–16</sup> (Table 1 and supplement 4).

### Respiratory symptoms

We collected data on cough, phlegm, and wheeze. Most cohorts collected data on all three of these outcomes (10 of 18); 72% of cohorts provided data on cough, 61% on phlegm, and 61% on wheeze. Respiratory symptoms were common in all cohorts (Supplement 5a). The median




Figure 1. World map of included AGRICOH countries.

**Table 1.** Characteristics of the AGRICOH cohorts that included respiratory outcomes.

Cohort	Country	Sample size	Year <sup>c</sup>	Male (%)	Mean age (std)	Age range	Smoking (%) <sup>b</sup>			BMI (%) <sup>b</sup>				
							Never	Past	Current	<20	20 - <25	25 - <30	≥30	
Africa														
Ethiopia <sup>13</sup>	Ethiopia	206	2014	69	27 (7)	15–57	95	0	5	32	56	11	1	
KwaZulu-Natal Crop Farmers <sup>14</sup>	South Africa	911	2006	0	42 (13)	18–82	94	0	6	NR	NR	NR	NR	
Western Cape Fruit Farm Workers <sup>15</sup>	South Africa	211	2009	0	37 (12)	17–73	50	0	50	17	28	22	33	
North West Poultry Workers <sup>16</sup>	South Africa	230	2012	68	37 (9)	21–68	56	1	43	17	48	17	17	
Asia														
KMCC <sup>17</sup>	Korea	8,431	2004	44	58 (10)	19–91	60	11	29	14	65	18	3	
Europe														
AGRICAN <sup>18</sup>	France	128,388	2007	56	65 (15)	20–105	65	27	9	4	37	44	15	
BM3R <sup>19</sup>	France	5,095	2013	59	57 (9)	40–75	65	22	14	4	46	31	19	
FERMA <sup>20</sup>	France	473	2012	71	47 (12)	18–78	55	24	21	8	45	29	17	
Franche-Comte Farmers <sup>21</sup>	France	915	2006	74	55 (11)	30–80	62	24	15	2	49	32	17	
Norwegian Farmers <sup>22</sup>	Norway	4,735	2009	60	49 (11)	21–69	50	22	28	2	43	44	11	
PIPAH <sup>23</sup>	United Kingdom	4,536	2014	98	54 (12)	19–88	65	26	9	2	41	44	13	
SUS Study <sup>24</sup>	Denmark	1,964	1992	88	19 (3)	17–49	70	0	30	NR	NR	NR	NR	
North America														
AHS <sup>a</sup>	USA	39,464	2016	53	65 (11)	32–104	68	27	5	3	22	40	35	
Farmer Health Study <sup>25</sup>	USA	1,947	1993	90	54 (13)	21–90	55	32	12	NR	NR	NR	NR	
KCRHS <sup>26</sup>	USA	1,256	1997	44	52 (16)	18–92	62	24	15	3	31	31	36	
Oceania														
New Zealand <sup>27</sup>	New Zealand	4,288	2005	52	49 (11)	20–93	58	30	12	4	36	44	17	
Victorian Grain Farmers <sup>28</sup>	Australia	1,102	1995	100	51 (13)	19–90	57	13	30	NR	NR	NR	NR	
South America														
MAUCO <sup>29</sup>	Chile	7,080	2015	39	54 (10)	36–77	45	23	31	1	16	44	40	

<sup>a</sup>Data not yet published.

<sup>b</sup>Percentages may not add up to 100% due to rounding.

<sup>c</sup>If multiple years included in cohort, the last year of data collection is provided.

BMI, body mass index; NR, not reported.

KMCC, Korean Multi-center Cancer Cohort; AHS, Agricultural Health Study; GDMSP, Grain Dust Medical Surveillance Program; KCRHS, Keokuk County Rural Health Study.


prevalence was 18.6% (IQR: 14.4–30.0%) for cough, 13.3% (IQR: 7.8–18.8%) for phlegm, and 15.0% (IQR: 8.8–16.1%) for wheeze. Farmworkers from South Africa (Western Cape Fruit Farm Workers<sup>15</sup> and North West Poultry Workers<sup>16</sup>) and France (FERMA<sup>20</sup>) had the highest prevalence of cough and phlegm among all cohorts. Western Cape Fruit Farm workers also had the highest prevalence of wheeze (31%), while the farmers in the AHS in the United States had the second highest prevalence of wheeze (22%). When the data were stratified by gender (Figure 2 (a–c), Supplement 5b), males were more likely to report respiratory symptoms than females, though

the IQR for the median values overlapped between males and females.

### Respiratory diseases

We collected data on asthma, allergic asthma, non-allergic asthma, and COPD based on each cohort's definitions (Supplement 6a). All cohorts provided data on asthma, and 39% provided information on allergic phenotypes. 83% of cohorts provided information on COPD.

The median prevalence of asthma was 7.2% (IQR 5.7%–10.1%) with individual study values ranging from 0.5% in the Ethiopian<sup>13</sup> cohort to 16.0%



**Figure 2.** Overall and gender-stratified (a) cough prevalence, phlegm prevalence (b), and wheeze prevalence (c) among AGRICOH cohorts.

among Victorian Grain Farmers<sup>28</sup> in Australia. Differences in asthma prevalence by gender did not follow the same trend as observed for symptoms (Figure 3(a,b)). The median prevalence was slightly

higher among females (7.8% [IQR: 6.5%-10.5%]) compared with males (6.5% [IQR: 4.0%-11.3%]).

In the seven cohorts with data on allergic and non-allergic asthma, allergic asthma (5.0% [IQR:



3.0%-5.1%]) was more common than non-allergic asthma (2.0% [IQR: 1.9%-7.0%]). This trend persisted after stratifying by gender; the median prevalence of allergic and non-allergic asthma among females was 5.5% and 3.5%, respectively, while among men, the median prevalence was 3.6% for both allergic and non-allergic asthma. Among the gender-stratified results, we also observed that women had higher prevalence ratios (PRs) than men, comparing allergic to non-allergic asthma (Figure 4). Upon aggregating data across all

cohorts, women (PR: 1.01 [95% CI: 0.95–1.06]) had a significantly higher prevalence ratio of allergic to non-allergic asthma compared to men (PR: 0.76 [95% CI: 0.72–0.82]), indicating that allergic asthma and non-allergic asthma prevalence were similar in women, but that men were less likely to have allergic asthma.

Information on COPD was provided by 14 of 18 cohorts. One third of the cohorts used spirometry to classify COPD; the remaining two thirds used self-reported doctor diagnosis of at least one of these





Figure 3. Overall and gender-stratified asthma prevalence (a) and COPD prevalence (b) among AGRICOH cohorts.



**Figure 4.** Gender-stratified ratio of allergic vs. non-allergic asthma within AGRICOH cohorts.

three outcomes (chronic bronchitis, emphysema, or COPD). The median prevalence of COPD among all cohorts was 4.5% [IQR: 2.8%-10.0%] with COPD prevalence ranging from 0.5% in the MAUCO<sup>29</sup> cohort to 14.6% among Norwegian<sup>22</sup> farmers. The studies that reported using spirometry to estimate COPD prevalence had values ranging from 1.9% to 14.6% (median = 10.0% [IQR: 4.5%-11.1%]); studies that used self-reported prevalence had values ranging from 0.5% to 11.9% for COPD (median = 3.7% [IQR: 2.8%-10.0%]). When we stratified the COPD data by gender, the median prevalence for males was 5.5% (IQR: 1.9%-12.2%) and for females was 4.0% (IQR: 2.0%-7.9%) (Supplement 6b).

## Discussion

Using the data from 18 agricultural cohorts from around the world, we sought to characterize the prevalence of respiratory symptoms and diseases among the AGRICOH consortium as indicators of respiratory health among farmers worldwide with particular focus on differences by gender. In this analysis, respiratory symptoms were common and higher among men than women; while respiratory diseases were less common and women had

a higher prevalence of asthma and men had higher prevalence of COPD. Although farming practices differ around the world, we found that the prevalence of these respiratory outcomes did not vary substantially between cohorts with any notable trends by region.

The 18 cohorts included here represent a small subset of agricultural workers worldwide. No comprehensive study of agricultural workers using a common protocol around the world has been conducted. While large, our sample is by no means comprehensive and the representation between developed and developing countries is uneven. Some cohorts included a relatively small number of farmers and focused only on specific types of farming, such as the all-female Western Cape Fruit Farmers<sup>15</sup> cohort in South Africa, while others featured much larger populations that included a wide variety of types of farming activities as well as spouses of farmers (e.g., AHS, AGRICAN, and Norwegian Farmers). Overall, the average age among our cohort members (56.8 years) is similar to the average age of farmers in both developed countries and across Africa (60 years), as reported by the Food and Agriculture Organization of the United Nations in 2014.<sup>33</sup>

Overall, males reported more respiratory symptoms than women, but this was not true for every cohort. In general population samples, such as the National Health and Nutrition Examination Survey (NHANES) in the United States, the prevalence of wheeze and cough is similar in men and women while the prevalence of phlegm is much more common in men.<sup>34</sup> Here we observe greater differences between men and women in cohorts limited to specific farm activities, such as poultry work, with a high potential for exposure to respiratory irritants (e.g., Northwest Poultry Workers). We were unable to evaluate specific farming practices due to both the small number of cohorts and the complex exposure experience of individuals within the larger cohorts (e.g., AHS, AGRICAN). Occupational exposures and the types of tasks that men are more likely to engage in may contribute to this higher outcome prevalence.<sup>3</sup> Smoking is more common in men, and that may also contribute to some of the observed differences. However, in our study sample, there is more variability in the prevalence of symptoms by cohort rather than between men and women, suggesting different baseline rates of symptoms in different populations as well as differences in the type of farming and related exposures.

Asthma prevalence was higher in women agricultural workers, likely due to the higher prevalence of allergic asthma in women. Similarly, women around the world have a higher prevalence of asthma than men. In an earlier analysis of AHS data from 2005 to 2010, women in the AHS had more asthma than men in the AHS, but lower prevalence of asthma compared to the US population.<sup>35</sup> This may be due to factors associated with who becomes a farmer (for example, people with a history of asthma may choose a different profession) as well as agricultural exposures that appear to reduce allergic outcomes in farmers and their families.<sup>3,8,36</sup> The prevalence of asthma among our cohorts ranged from <1% to 16% with a median prevalence of 7.2%. Globally, asthma prevalence in adults aged 18–45 from 70 countries in 2002–2003 ranged from 0.2 (China) to 21% (Australia) with an overall prevalence of 4.3%.<sup>37</sup> While that study did not include all the same countries included here, it suggests that farmers may have a higher prevalence of asthma than the general population. Surprisingly, we observed a higher prevalence

of allergic asthma than non-allergic asthma in the cohorts with data on this outcome (5.0% vs. 2.0%); this difference may have been due to higher prevalence of asthma in women and that women had a higher prevalence of allergic asthma. There is an extensive literature suggesting that farmers have a lower prevalence of allergic asthma.<sup>3,4,38</sup> We cannot evaluate whether our findings are due to the diversity of our cohorts over continents and agricultural practices, or due to the primarily cross-sectional data used, or the fact that allergy may have been assigned based on questionnaire information. Future studies should focus on better characterization of allergic asthma.

The COPD results highlight the challenge of combining data across cohorts when disease definitions vary greatly, as has been reported by others.<sup>39</sup> Among the 15 cohorts that evaluated COPD prevalence, six different definitions of COPD were employed. COPD is classified based both on symptoms (chronic cough and phlegm) as well as airway obstruction as assessed through spirometry.<sup>40</sup> COPD encompasses the spectrum of obstructive diseases and includes chronic bronchitis and emphysema. Not all cohorts used spirometry to diagnose COPD, but rather relied on doctor diagnosis of COPD. This will capture individuals who have been diagnosed accurately by their physician, but will miss those who have not had this detailed clinical work up; suggesting a sensitive, but not specific outcome. Additionally, it will capture those who received a diagnosis but did not meet the clinical criteria, creating false positives.<sup>41</sup> Within an individual cohort, these biases may behave in the same manner, but between cohorts, the factors that influence diagnosis and over reporting may differ making it almost impossible to combine across these studies.

Although this study importantly characterizes farming cohorts around the world, it does feature some limitations. Comparability between cohorts may be limited due to differences in study year, ranging from 1992 to 2016. Cohorts ranged from small, focused cohorts with detailed information on a specific type of agriculture to large studies that include farmers engaged in a wide variety of activities. In choosing to include as many cohorts as possible, we forfeit the ability to focus on specific types of agriculture or agricultural practices.

Participating cohorts provided demographic and symptom and disease metadata, but did not share participant-level data. As a result, the prevalence estimates reported are not adjusted or standardized to account for important confounding variables such as smoking, age, status, income, education, and farming practices. While the AGRICOH consortium includes cohorts from around the world, the analysis was limited to those 15 cohorts, and three additional ones that evaluated respiratory symptoms and diseases around the world. Consequently, our descriptive analysis only features study populations from 11 countries, and more work is needed to truly characterize the global burden of respiratory outcomes in agricultural workers. Given that this study included only farmers, we were unable to make direct comparisons to the general public and assess the impact of agricultural production on respiratory health. The lack of national data on asthma prevalence makes it difficult to compare both within countries by occupation and between countries with regard to national prevalence. Global data on asthma prevalence will allow more rigorous analysis of variability by occupation and country.

International collaborative research provides valuable opportunities to investigate symptom and disease burden across many regions and geographic settings and diverse study populations. Future research would greatly benefit from sharing of participant-level data, allowing estimates to be standardized or adjusted for important confounders such as what is being done by the OMEGA NET consortium,<sup>42</sup> and improving comparability between cohorts by applying common protocols like the ISAAC study for children<sup>43</sup> and the new DEGREE study for epidemiologic studies in low- and middle-income populations.<sup>44</sup> Data harmonization and standardization are not trivial tasks and we were unable to perform these at this time.

Asthma and COPD are important contributors to morbidity and mortality worldwide.<sup>39</sup> Our analysis shows that farmers and agricultural workers are impacted by these health outcomes. Agriculture represents a complex exposure environment and respiratory diseases are multi-factorial in nature. As a result, it is difficult to identify specific etiologic factors that contribute to or

protect against respiratory disease. However, in these diverse cohorts, adult respiratory disease and symptoms were common and, it is likely that occupation contributed to these outcomes. The United Nation's International Labor Organization estimated in 2009 that there were about 1.07 billion people employed in agriculture around the world, accounting for nearly 35% of the global workforce.<sup>45</sup> Therefore, understanding the global impacts of agricultural production on respiratory health is critical.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by Dr. Hoppin's startup funds at NC State University. The Keokuk County Rural Health Study was supported by Grants # (U07/CCU706145) (Round 1) and # (U50 OH07548) (Round 2 and Round 3) from the National Institute for Occupational Safety and Health to the Great Plains Center for Agricultural Health, The University of Iowa. This research was supported in part by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences (Z01-ES049030). KwaZulu-Natal was supported by the South African Netherlands Research Program on Alternatives in Development (SANPAD) and the South African Medical Research Council (SA MRC) – grant number 04/15. MAUCO was supported by Conicyt – FONDAP No (15130011).

## ORCID

Isabella Annesi-Maesano  <http://orcid.org/0000-0002-6340-9300>

Mohamed Jeebhay  <http://orcid.org/0000-0001-6656-9193>

Saloshni Naidoo  <http://orcid.org/0000-0002-8844-0160>

Dorothy Ngajilo  <http://orcid.org/0000-0001-5267-8081>

Marc B Schenker  <http://orcid.org/0000-0002-8846-4357>

Torben Sigsgaard  <http://orcid.org/0000-0002-2043-7571>

Jane A Hoppin  <http://orcid.org/0000-0001-8456-0969>

## References

1. Schenker MB, Christiani D, Cormier Y, et al. Respiratory health hazards in agriculture. *Am J Respir Crit Care Med.* 1998;158(5):S1–S76.

2. Kirkhorn SR, Garry VF. Agricultural lung diseases. *Environ Health Perspect.* 2000 Aug;108(Suppl 4):705–712. doi:10.1289/ehp.00108s4705.
3. Eduard W, et al. Do farming exposures cause or prevent asthma? Results from a study of adult Norwegian farmers. *Thorax.* 2004 May;59(5):381–386. doi:10.1136/thx.2004.013326.
4. Eduard W, Omenaas E, Bakke PS, et al. Atopic and non-atopic asthma in a farming and a general population. *Am J Ind Med.* 2004 Oct;46(4):396–399. doi:10.1002/(ISSN)1097-0274.
5. Hoppin JA, Umbach DM, London SJ, et al. Pesticides and atopic and nonatopic asthma among farm women in the agricultural health study. *Am J Respir Crit Care Med.* 2008 Jan 1;177(1):11–18. doi:10.1164/rccm.200706-821OC.
6. Hoppin JA, Umbach DM, London SJ, et al. Pesticide use and adult-onset asthma among male farmers in the agricultural health study. *Eur Respir J.* 2009 Dec;34(6):1296–1303. doi:10.1183/09031936.00005509.
7. Hoppin JA, Umbach DM, Long S, et al. Pesticides are associated with allergic and non-allergic wheeze among male farmers. *Environ Health Perspect.* 2016 Jul 06.
8. von Mutius E. The environmental predictors of allergic disease. *J Allergy Clin Immunol.* 2000;105(1 Pt 1):9–19. doi:10.1016/S0091-6749(00)90171-4.
9. Schachter EN, Zuskin E, Moshier EL, et al. Gender and respiratory findings in workers occupationally exposed to organic aerosols: a meta analysis of 12 cross-sectional studies. *Environ Health.* 2009 Jan 12;8:1. doi:10.1186/1476-069X-8-1.
10. Senthilselvan A, Chenard L, Ulmer K, et al. Excess respiratory symptoms in full-time male and female workers in large-scale swine operations. *Chest.* 2007 Apr;131(4):1197–1204. doi:10.1378/chest.06-2323.
11. Motika CA, Papachristou C, Abney M, et al. Rising prevalence of asthma is sex-specific in a US farming population. *J Allergy Clin Immunol.* 2011 Oct;128(4):774–779. doi:10.1016/j.jaci.2011.06.046.
12. Leon ME, Beane Freeman LEB, Douwes J, et al. AGRICOH: a consortium of agricultural cohorts. *Int J Environ Res Public Health.* 2011 May;8(5):1341–1357. doi:10.3390/ijerph8051341.
13. Negatu B, Kromhout H, Mekonnen Y, et al. Occupational pesticide exposure and respiratory health: a large-scale cross-sectional study in three commercial farming systems in Ethiopia. *Thorax.* 2017 Jun;72(6):498–499. doi:10.1136/thoraxjnl-2016-208924.
14. Naidoo S, London L, Burdorf A, et al. Agricultural activities, pesticide use and occupational hazards among women working in small scale farming in Northern KwaZulu-Natal, South Africa. *Int J Occup Environ Health.* 2008 Jul-Sep;14(3):218–224. doi:10.1179/oeht.2008.14.3.218.
15. Mwanga HH, Dalvie MA, Singh TS, et al. Relationship between pesticide metabolites, cytokine patterns, and asthma-related outcomes in rural women workers. *Int J Environ Res Public Health.* 2016 Sep 27;13(10). doi:10.3390/ijerph13100957
16. Ngajilo D, Singh T, Ratshikhopha E, et al. Risk factors associated with allergic sensitization and asthma phenotypes among poultry farm workers. *Am J Ind Med.* 2018 Mar 25. doi:10.1002/ajim.22841.
17. Yoo KY, Shin HR, Chang SH, et al. Korean multi-center cancer cohort study including a biological materials bank (KMCC-I). *Asian Pac J Cancer Prev.* 2002;3(1):85–92.
18. Baldi I, Robert C, Piantoni F, et al. Agricultural exposure and asthma risk in the AGRICAN French cohort. *Int J Hyg Environ Health.* 2014 Apr-May;217(4–5):435–442. doi:10.1016/j.ijheh.2013.08.006.
19. Guillien A, Puyraveau M, Soumagne T, et al. Prevalence and risk factors for COPD in farmers: a cross-sectional controlled study. *Eur Respir J.* 2016 Jan;47(1):95–103. doi:10.1183/13993003.00153-2015.
20. Audi C, Baiz N, Maesano CN, et al. Serum cytokine levels related to exposure to volatile organic compounds and PM2.5 in dwellings and workplaces in French farmers – a mechanism to explain nonsmoking COPD. *Int J Chron Obstruct Pulmon Dis.* 2017;12:1363–1374. doi:10.2147/COPD.S117866.
21. Gainet M, Thaon I, Westeel V, et al. Twelve-year longitudinal study of respiratory status in dairy farmers. *Eur Respir J.* 2007 Jul;30(1):97–103. doi:10.1183/09031936.00150405.
22. Eduard W, Pearce N, Douwes J. Chronic bronchitis, COPD, and lung function in farmers: the role of biological agents. *Chest.* 2009 Sep;136(3):716–725. doi:10.1378/chest.08-2192.
23. Harding A-H, Fox D, Chen Y, et al. Prospective investigation of pesticide applicators' health (PIPAH) study: a cohort study of professional pesticide users in Great Britain. *BMJ Open.* 2017 Oct 10;7(10):e018212. doi:10.1136/bmjopen-2017-018212.
24. Sigsgaard T, Hjort C, Omland Ø, et al. Respiratory health and allergy among young farmers and non-farming rural males in Denmark: the SUS study. *J Agromedicine.* 2004;9(2):223–238.
25. Schenker MB, Farrar JA, Mitchell DC, et al. Agricultural dust exposure and respiratory symptoms among California farm operators. *J Occup Environ Med.* 2005 Nov;47(11):1157–1166. doi:10.1097/01.jom.0000181174.02282.0c.
26. Merchant JA, Stromquist AM, Kelly KM, et al. Chronic disease and injury in an agricultural county: the Keokuk county rural health cohort study. *J Rural Health.* 2002 Fall;18(4):521–535. doi:10.1111/j.1748-0361.2002.tb00919.x
27. Douwes J, Travier N, Huang K, et al. Lifelong farm exposure may strongly reduce the risk of asthma in adults. *Allergy.* 2007 Oct;62(10):1158–1165. doi:10.1111/all.2007.62.issue-10.
28. Macfarlane E, Chapman A, Benke G, et al. Training and other predictors of personal protective equipment

- use in Australian grain farmers using pesticides. *Occup Environ Med.* 2008 Feb;65(2):141–146. doi:10.1136/oem.2007.034843.
29. Ferreccio C, Roa JC, Bambs C, et al. Study protocol for the Maule Cohort (MAUCO) of chronic diseases, Chile 2014–2024. *BMC Public Health.* 2016 Feb 04;16:122. doi:10.1186/s12889-015-2454-2.
  30. World Bank. 2017. *World Development Indicators 2017*. Washington, D.C.: World Bank; 2017. <https://openknowledge.worldbank.org/handle/10986/26447> License: CC BY 3.0 IGO.
  31. Ferris BG. Epidemiology standardization project (American Thoracic Society). *Am Rev Respir Dis.* 1978 Dec;118(6 Pt 2):1–120.
  32. Heinrich J, Richter K, Frye C, et al. European community respiratory health survey in adults (ECRHS). *Pneumologie.* 2002 May;56(5):297–303. doi:10.1055/s-2002-30699.
  33. FAO. *Food Security for Sustainable Development and Urbanization.* 2014. <https://www.un.org/en/ecosoc/integration/pdf/foodandagricultureorganization.pdf>
  34. (CDC) Centers for Disease Control and Prevention NCHS. NHANES 1999–2000 Data Files 2005 [cited 2005 Oct 19]. Available from: [http://www.cdc.gov/nchs/about/major/nhanes/nhanes99\\_00.htm](http://www.cdc.gov/nchs/about/major/nhanes/nhanes99_00.htm)
  35. Hoppin JA, Umbach DM, Long S, et al. Respiratory disease in United States farmers. *Occup Environ Med.* 2014;71(7):484–489. doi:10.1136/oemed-2013-101983.
  36. Vogelzang PF, van der Gulden JW, Tielen MJ, et al. Health-based selection for asthma, but not for chronic bronchitis, in pig farmers: an evidence-based hypothesis. *Eur Respir J.* 1999;13(1):187–189. doi:10.1034/j.1399-3003.1999.13a34.x.
  37. To T, Stanojevic S, Moores G, et al. Global asthma prevalence in adults: findings from the cross-sectional world health survey. *BMC Public Health.* 2012 Mar;19(12):204. doi:10.1186/1471-2458-12-204.
  38. Elholm G, Schlunssen V, Doekes G, et al. Become a farmer and avoid new allergic sensitization: adult farming exposures protect against new-onset atopic sensitization. *J Allergy Clin Immunol.* 2013 Nov;132(5):1239–1241. doi:10.1016/j.jaci.2013.07.003.
  39. Collaborators GBDCRD. Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the global burden of disease study 2015. *Lancet Respir Med.* 2017 Sep;5(9):691–706. doi:10.1016/S2213-2600(17)30293-X.
  40. Luize AP, Menezes AM, Perez-Padilla R, et al. Assessment of five different guideline indication criteria for spirometry, including modified GOLD criteria, in order to detect COPD: data from 5,315 subjects in the PLATINO study. *NPJ Prim Care Respir Med.* 2014 Oct 30;24:14075. doi:10.1038/npjpcrm.2014.75.
  41. Bobadilla A, Guerra S, Sherrill D, et al. How accurate is the self-reported diagnosis of chronic bronchitis? *Chest.* 2002 Oct;122(4):1234–1239. doi:10.1378/chest.122.4.1234.
  42. Turner MC, Mehlum IS. Greater coordination and harmonisation of European occupational cohorts is needed. *Occup Environ Med.* 2018 Jul;75(7):475–476. doi:10.1136/oemed-2017-104955.
  43. ISAAC Steering Committee. *International Study of Asthma and Allergies in Childhood. ISAAC Phase One Manual.* Auckland/Munster; 1993. <http://isaac.auckland.ac.nz/phases/phaseone/phaseonemanual.pdf>
  44. Caplin B, Jakobsson K, Glaser J, et al. International collaboration for the epidemiology of eGFR in low and middle income populations – rationale and core protocol for the disadvantaged populations eGFR epidemiology study (DEGREE). *BMC Nephrol.* 2017 Jan 3;18(1):1. doi:10.1186/s12882-016-0417-1.
  45. Kapsos S, Sparreboom T. *The Challenge of a Jobs Recovery.* Geneva: International Labour Office; 2011.