



Targeted LC-ESI-MS² characterization of human milk oligosaccharide diversity at 6 to 16 weeks post-partum reveals clear staging effects and distinctive milk groups

Marko Mank¹ · Hans Hauner^{2,3} · Albert J. R. Heck^{4,5} · Bernd Stahl^{1,6}

Received: 18 April 2020 / Revised: 3 June 2020 / Accepted: 14 July 2020 / Published online: 14 August 2020
© The Author(s) 2020

Abstract

Many molecular components in human milk (HM), such as human milk oligosaccharides (HMOs), assist in the healthy development of infants. It has been hypothesized that the functional benefits of HM may be highly dependent on the abundance and individual fine structures of contained HMOs and that distinctive HM groups can be defined by their HMO profiles. However, the structural diversity and abundances of individual HMOs may also vary between milk donors and at different stages of lactations. Improvements in efficiency and selectivity of quantitative HMO analysis are essential to further expand our understanding about the impact of HMO variations on healthy early life development. Hence, we applied here a targeted, highly selective, and semi-quantitative LC-ESI-MS² approach by analyzing 2 × 30 mature human milk samples collected at 6 and 16 weeks post-partum. The analytical approach covered the most abundant HMOs up to hexasaccharides and, for the first time, also assigned blood group A and B tetrasaccharides. Principal component analysis (PCA) was employed and allowed for automatic grouping and assignment of human milk samples to four human milk groups which are related to the maternal Secretor (Se) and Lewis (Le) genotypes. We found that HMO diversity varied significantly between these four HM groups. Variations were driven by HMOs being either dependent or independent of maternal genetic Se and Le status. We found preliminary evidence for an additional HM subgroup within the Se- and Le-positive HM group I. Furthermore, the abundances of 6 distinct HMO structures (including 6'-SL and 3-FL) changed significantly with progression of lactation.

Keywords Human milk oligosaccharides (HMOs) · Stages of lactation · Human milk groups/milktyping · Targeted LC-MS² · Principal component analysis (PCA) · Secretor (Se) and Lewis (Le) gene-dependent or Secretor (Se) and Lewis (Le) gene-independent HMOs · 3-FL · 6'-SL

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00216-020-02819-x>) contains supplementary material, which is available to authorized users.

✉ Marko Mank
marko.mank@danone.com

¹ Danone Nutricia Research, Uppsalalaan 12, 3584 CT Utrecht, The Netherlands

² Else Kröner-Fresenius Center for Nutritional Medicine, Klinikum rechts der Isar, Technische Universität München, Ismaninger Straße 22, 81675 Munich, Germany

³ Nutritional Medicine Unit, Research Center for Nutrition and Food Sciences (ZIEL), Technische Universität München, Weihenstephaner Berg 1, 85354 Freising, Germany

⁴ Biomolecular Mass Spectrometry and Proteomics, Bijvoet Center for Biomolecular Research and Utrecht Institute for Pharmaceutical Sciences, University of Utrecht, Padualaan 8, 3584 CH Utrecht, The Netherlands

⁵ Netherlands Proteomics Center, Padualaan 8, 3584 CH Utrecht, The Netherlands

⁶ Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Universiteitsweg 99, 3584 CG Utrecht, The Netherlands

Introduction

Human milk (HM) is largely defined by the abundance of 5 major constituents comprising lactose, lipids, oligosaccharides, proteins, and water. Published values for lactose, lipid, and protein concentrations in mature milk 1–2 months post-partum (pp) are 67–78 g/l, 32–36 g/l, and 9–12 g/l [1] respectively. Even broader ranges in concentration of HM components apply if also colostrum and transitional milks (5–14 days pp) are considered [2]. The reported concentrations for total HMOs in mature milk range between ~3 and 18 g/l [3–8]. Until now, more than 150 HMO structures have been revealed and confirmed by various analytical approaches including HPLC, mass spectrometry, and NMR to name a few [9–13]. Abundant individual HMOs like 2'-fucosyllactose (2'-FL), lacto-N-tetraose (LNT), lacto-N-neotetraose (LNnT), lacto-N-fucopentaose I (LNFP I), difucosyl-lacto-N-hexaose II (DF-LNH II), trifucosyl-lacto-N-hexaose (TF-LNH), 6'-sialyllactose (6'-SL), and disialyl-lacto-N-tetraose (DSLNT) may be present in the low gram per liter range [3]. This set of known HMO structures represents only a few of more than 1000 compositional or structural variants as anticipated from MALDI-MS data of high molecular weight HMO fractions with molecular masses > 8 kDa. The latter mass range translates into a degree of polymerization (DP) for particular HMOs > 43 [4].

Additional variations in human milk oligosaccharide profiles can be introduced by preterm delivery (lower lactose concentrations compared with term birth [14]) or are observed at different stages of lactation. In fact, literature reports an overall decrease of HMOs [15–17] over time during stages of lactation and the highest concentration of approximately 20 g/l total HMOs for term delivered infants is described to appear on around day 4 post-partum [18]. Staging effects are also observed for other mammalian milks like bovine or caprine milk [19–22]. Finally, also regional differences may have an influence on HMO variations exemplified by 100% presence of 2'-FL in milks of 156 Mexican women versus only 46% presence in milks of 26 Philippine donors [23]. Beyond their presence in human milk, HMOs have also been found in blood samples of both, mothers (gestational week 10–35) and breastfed offspring [24–28] and are even found in the amniotic fluid [29].

Bovine milk which is the basis for many infant milk formulations contains on average less lactose (approximately 40–50 g/l lactose [30]) and only minute amounts of bovine milk oligosaccharides (BMOs). BMOs comprise about 40 structures [31]. Among these BMOs, 3'-sialyllactose and 6'-sialyllactose are prevalent, but also galactosyllactoses (Gal-Lacs) may be of interest. As reported by Nakamura et al. in 2003 [20], sialyllactose concentrations in bovine milk samples from 3 days post-partum range between 0.1 g/L for 3'-SL and < 0.05 g/L for 6'-SL. Mature bovine milk contains approximately 0.03–0.12 g/L 3'-SL and 0.03–0.09 g/L 6'-SL [32]

which is below the mean concentrations found in term human milk for 3'-SL (0.16 g/L) and 6'-SL (0.6 g/L) [3].

Biosynthesis of HMOs is governed by a limited set of glycosyltransferases present in the mammary gland and human milk itself [12]. Therefore, with only a few exceptions, the structural arrangement of most human milk oligosaccharides can be described by the following general scheme (see also Fig. 1): The sequence of each HMO molecule starts with a lactose building block (Gal β 1–4Glc) introducing a reducing end which is a common feature of all known HMOs described until now. This lactose building block can be extended with a dimeric residue consisting of galactose and *N*-acetylglucosamine (Gal-GlcNAc). As suggested by Kobata et al. 2010 [42], the transfer of Gal-GlcNAc to the terminal Gal of lactose may be accomplished by consecutive action of either β -1,3-*N*-acetylglucosaminyltransferase (iGnT) [43] and β 1,4-galactosyltransferases (β 4GalTs) [44] or iGnT and β 1,3-galactosyltransferases (β 3GalTs) [45]. Thereby, two basic types of HMO core structures can be created: a type I core (Gal β 1–3GlcNAc β 1–3Gal β 1–4Glc, LNT) or a type II core molecule (Gal β 1–4GlcNAc β 1–3Gal β 1–4Glc, LNnT). In HM, type I core HMOs are predominant over type II structures [46]. This feature distinguishes HM from other species including apes and monkeys. According to Urashima et al. [46], one possible explanation for this phenomenon is the role of type I oligosaccharides as important substrates for beneficial bifidobacteria. Moreover, the two types of neutral core entities (i.e., LNT or LNnT) may be further extended by successive addition of more Gal-GlcNAc blocks. If Gal-GlcNAc residues are appended to the core structures via β 1,3-glycosidic linkage, no further elongation will occur after this Gal-GlcNAc block. If, in contrast, Gal-GlcNAc dimers are added via action of β 6*N*-acetylglucosaminyltransferase (iGNT) [47] through β 1,6-glycosidic linkages, branching will occur and further elongation of the β 1,6-glycosidically linked branches is possible. Thereby, up to 19 complex HMO core structures can be created [38]. Finally, these core structures or lactose itself may be furthermore decorated by additional fucoses (Fuc), and sialic acid (Neu5Ac) moieties through human fucosyl- [33] or sialyl-transferases [34]. However, it remains to be explored which of the known human sialyl- and fucosyltransferases are actually actively involved in the mammary gland biosynthesis of HMOs. More certainty already exists with respect to HMO fucosylation exerted by the *Se*-enzyme [35] and fucosyltransferase 3 [48] as explained in the next paragraph. Finally, exceptions from the general HMO building scheme outlined above may exist and are e.g. exemplified by the presence of galactosyllactoses (GL) in human milk. Here, only a single Gal is added to lactose. Literature describes several galactosyllactose isomers such as 6'-GL, 3'-GL, and 4'-GL. To date, information about GL concentrations in HM is still rare, but first published levels are in the milligram per liter range [49].




Fig. 1 Major HM HMO structures and related glycosyltransferases [9, 33–36] involved in their mammary gland biosynthesis starting from lactose as a precursor. Certain HMOs resemble ABH [37] or Se/Le system-related antigens [38, 39]. 3-Fucosyllactose (3'-FL), 2'-fucosyllactose (2'-FL), 3'-sialyllactose (3'-SL), 6'-sialyllactose (6'-SL), difucosyllactose (DFL), lacto-N-tetraose (LNT); lacto-N-neotetraose (LNnT); tentative blood group B tetrasaccharide (Hex)₃(Fuc)₁; tentative blood group A tetrasaccharide (Hex)₂(HexNAc)₁(Fuc)₁; lacto-N-

fucopentaose I (LNFP I); lacto-N-fucopentaose II (LNFP II); lacto-N-fucopentaose III (LNFP III); lacto-N-fucopentaose V (LNFP V), lacto-N-fucopentaose VI (LNFP VI = LNnFP V), lacto-N-neodifucohexaose I (LNDFH I), lacto-N-neodifucohexaose II (LNDFH II), difucosyllactose (DFL), lacto-N-tetraose (LNT); lacto-N-neotetraose (LNnT); tentative blood group B tetrasaccharide (Hex)₃(Fuc)₁; tentative blood group A tetrasaccharide (Hex)₂(HexNAc)₁(Fuc)₁; lacto-N-

Inherited maternal genetic predispositions may induce remarkable inter-individual differences in oligosaccharide synthesis and resulting individual HMO profiles. For instance, Secretor (Se) and Lewis (Le) genes on chromosome 19 encode for two different fucosyltransferases: the Se-enzyme and fucosyltransferase 3 (Fuc-TIII) [33]. The Se-enzyme is expressed in the case of an active Se gene (FUT 2) and Fuc-TIII in the case of an active Le gene (FUT 3). The Se-enzyme specifically appends fucose (Fuc) via α1,2-glycosidic linkage to type 1 HMOs (e.g., lacto-N-tetraose (LNT)) or to type 2 HMOs (e.g., lacto-N neotetraose (LNnT)),

whereas in contrast Fuc-TIII catalyzes the formation of α1,3/4-fucosylated HMOs. Primarily catalyzed by the Se-enzyme and Fuc-TIII, 4 different ways of HMO-fucosylation at distinct monosaccharides and distinct sites within type I or II core HMO structures can occur [12]: α1,2-linkage to non-reducing end Gal, α1,3- or α1,4-linkage to N-acetylglucosamine (GlcNAc), or α1,3-linkage to reducing end Glc. Neutral or sialylated acidic HMOs with core structures smaller or larger than tetraoses can be decorated in the same way and lead to a multitude of known or hypothetically possible fucosylated HMO structures.

Figure 1 schematically exemplifies fucosylation of LNT (resembling a Lec-antigen [39]) and LNnT (Le I-antigen) by activity of Se-enzyme, FucTIII, or further α 1,3-fucosyltransferases [50]. Resulting HMOs are e.g. lacto-N-fucopentaoses (LNFPs) such as LNFP I, II, III, and V. Such fucosylated HMOs can resemble ABO- or Se/Le system-related antigens like the Le^d-antigen in case of LNFP I, the Le^a-antigen in case of LNFP II, or the Le^x-antigen in case of LNFP III [37, 39]. Additionally, LNFP VI (the type II core variant of LNFP V) may be formed, too. It was recently detected in human milk by Bao et al. [51] using negative ion mode LC-MS after conversion of HMOs into alditols. Other products of the Se-enzyme, Fuc-TIII (or other α 1,3-fucosyltransferases) are human milk oligosaccharides like fucosyllactoses (2'FL or 3-FL), difucosyllactose (DFL), the lacto-N-difucohexaoses LNDFH I (Le^b-antigen), LNnDFHI (Le^Y-antigen), LNDFH II and LNnDFH II, or higher molecular weight neutral and acidic HMOs with either linear or branched core structures.

Based on the presence, absence, or concomitance of α 1,2- and α 1,3/4-fucosylated HMOs like LNFP I and LNFP II in human milk, 4 different human milk groups (groups I, II, III, and IV) are currently distinguished [52, 53]. However, just recently, group I milks from Chinese donor were further subdivided into 2 new subgroups based on significantly different levels in e.g. 2'-FL, DFL, LNT, LNFP I, or F-LNO [54]. Assignment of human milk samples to one of the 4 main human milk groups I–IV can e.g. be performed by ESI-LC-MS or HPAEC HMO analysis and subsequent evaluation e.g. of the respective LNFP I and LNFP II patterns. Moreover, corresponding Le and Se gene product-dependent glycan epitopes may also be found in maternal blood, i.e., on erythrocytes or glycoproteins like mucins [55]. As mentioned before, other α 1,3-fucosyltransferases such as plasma α -1,3-fucosyltransferase (E.C. 2.4.1.152), different from Se-enzyme or Fuc-TIII and independent from the Le blood group system, may be active in human milk donors as well [39, 56, 57].

Table 1 summarizes the interdependence of maternal Lewis and Secretor genotypes, encoded fucosyltransferases, Lewis blood group phenotypes, related carbohydrate epitopes, resulting milk groups, and some prevalent HMO structures found in specific milk groups.

HMOs confer many intriguing functional properties of which some are likely to facilitate or influence healthy development of neonates [59–63]. Many of the published prebiotic, anti-infective, or immunomodulatory properties are highly structure specific [64–70]. They may vary between HMOs of the same monosaccharide composition, but different core types or fucosylation sites like LNFP I, LNFP II, or LNFP III. Also blood group A and B carbohydrate determinants can be present as free carbohydrates in HM [71]. Together with galactosyllactoses, they deviate from the HMO building scheme described above. Nevertheless, they may bear anti-

infective potential relevant for early life and beyond, e.g., by blocking the cholera toxin (CT) subunit B (CTB) of *Vibrio cholerae* from adhesion to mucins and intestinal epithelial cells of the small intestine [72].

As outlined above, different structure-effect relationships could be attributed to HMOs which are isomers but share the same monosaccharide composition. Thus, development and application of hyphenated analytical methods like LC-MS which can distinguish such subtle structural HMO features may be key to decipher still unknown (beneficial) biological functions of HMOs and other glycans. To accelerate the analysis of larger sample numbers available from e.g. human milk cohorts or clinical studies, analytical approaches with short run times and avoidance of tedious sample pre-treatment would be of advantage. In turn, this will help to improve statistical association of distinct HMO patterns with related health benefits for infants. In addition, (semi-)quantitative assays should be preferred over mere qualitative ones to allow for elucidation of concentration-dependent biological HMO effects.

Therefore, in this work, we applied a recently developed targeted semi-quantitative LC-ESI-MS² approach with high structural selectivity [73], easy sample preparation, and comparatively low run times of ~18 min per sample to 2 × 30 human milk specimen. These specimens were derived from a control group of the INFAT study which was not subjected to nutritional intervention. The used sample set comprised human milks collected at 6 and 16 weeks post-partum to elicit possible changes in HMO profiles between those 2 different stages of lactation.

Principal component analysis (PCA) was employed as a convenient and efficient strategy for mining of structural HMO information inherent in the acquired multiple reaction monitoring (MRM) LC-ESI-MS² data sets.

Experimental section

Human milk samples from INFAT study with unknown milk group and HMO status

A subset of 60 human milk samples derived from 30 healthy women resident in southern Germany and enrolled in a dietary intervention study called the INFAT study [74, 75] was subjected to MRM LC-ESI-MS² analysis. Every woman donated 2 milk samples at 6 and 16 weeks post-partum. Written consent from the donors was provided prior to analysis and approval was given by the ethical committee of the Faculty of Medicine of the Technical University of Munich, Germany (project number 1479/06 on 14 September 2012). The human milk samples used for this LC-MS investigation were derived from a control group not subjected to dietary intervention. Samples were obtained as described elsewhere [74]. Briefly, maternal breast milk samples were collected in the study

Table 1 Interdependence of maternal Lewis and Secretor genotypes, encoded fucosyltransferases (Fuc-Ts), Lewis blood groups, resulting carbohydrate epitopes, HMO structures, and milk groups. *2'-FL*, 2'-fucosyllactose; *LDFT*, lacto-Di-fucotetraose; *LNFP I*, lacto-N-fucopentaose I; *LNFP II*, lacto-N-fucopentaose II; *LNFP III*, lacto-N-fucopentaose III; *LNFP V*, lacto-N-fucopentaose V; *LNDFH I*, lacto-N-difucohexaose I; *LNDFH II*, lacto-N-difucohexaose II. Adapted from Oriol et al. 1986, Thurl et al. 1997, and Blank et al. 2011

Genotypes	Encoded fucosyl-transferases/gene loci	Resulting phenotypes		Human milk group	Prominent neutral HMO structures up to hexaoses ^a
		Erythrocytes	Human milk		
Secretor	Lewis		Fucosylated carbohydrate epitopes		
		Lewis	Prevalence in France [39]		
Se/-	Le/-	Se-enzyme/FUT2, Fuc-TIII/FUT3	a-b+c-d- 69	Fuc (α1-2)R, Fuc (α1-4)R, Fuc (α1-3)YR I	2'FL , 3'-FL, DFL , LNT, LNnT, LNFP I , LNFP II , LNFPIII, LNDFH I , LNDFH II
se/se	Le/-	Fuc-T III/FUT3	a+b-c-d- 20	Fuc (α1-4)R, Fuc (α1-3)YR II	3'-FL, LNT, LNnT, LNFP II , LNFPIII, LNDFH II ; (<i>2'FL</i> , <i>DFL</i> , <i>LNFP I</i> , <i>LNDFH I</i>)
Se/-	le/le	Se-enzyme/FUT2	a-b-c-d+ 9	Fuc (α1-2)R, Fuc (α1-3)YR III	2'FL , 3'-FL, DFL , LNT, LNnT, LNFP I , LNFPIII; (<i>LNFP II</i> , <i>LNDFH I</i> , <i>LNDFH II</i>)
se/se	le/le	Le- and Se- independent Fuc-Ts (e.g., Fuc-TVI/FUT6 [57])	a-b-c+d- 1	Fuc (α1-3)YR IV	3'-FL, LNT, LNnT, LNFPIII; (<i>2'FL</i> , <i>DFL</i> , <i>LNFP I</i> , <i>LNFP II</i> , <i>LNDFH I</i> , <i>LNDFH II</i>)

^a Bold black letters indicate major HMO structures present in human milks due to active Le or Se genes; italic letters in parentheses indicate HMO structures lacking or being present in very low concentrations in human milk due to fully or partly inactive Le or Se genes; regular black letters indicate prominent HMOs occurring in HM independent from maternal Le/Se activity

^b According to Hanfland et al. 1986 [58], there may be more than one Le^e antigenic determinant: Le^e epitopes may comprise non-fucosylated type I or type II glycan residues or extended unbranched or branched glycan residues with α1-3-fucosylation at the penultimate GlcNAc positioned before the non-reducing end of the glycan chain

^c Fuc (α1-3)-R epitope can also be catalyzed by Le/Se-independent FUT, e.g., FUT 6 [57]

center in the morning after an overnight fast in a standardized manner using a breast pump, aliquoted, and immediately stored at $-86\text{ }^{\circ}\text{C}$ until analysis. Samples were thawed once to take aliquots for different kinds of analysis and stored again at $-20\text{ }^{\circ}\text{C}$ until further usage for negative ion mode MRM LC-MS² analysis.

Chemicals, reagents, and disposables

Ammonium acetate (p.a.), acetonitrile (LC-MS grade), and ethanol (gradient grade for HPLC) were from Merck, Darmstadt, Germany. LC-MS grade H₂O (HiPerSolv Chromanorm) was from ProLabo, VWR International, Darmstadt, Germany. Alternatively, ultrapure lab water, 18 k Ω , < 5 ppm TOC produced by a MilliQ Advantage Ultrapure Water System from Merck Millipore (Darmstadt, Germany) was used. Clear glass 300- μl micro LC-MS vials were from Thermo Fisher Scientific (Waltham, MA, USA, #60180-507). Micro vial screw caps with pre-slit PTF/silicone membrane were from Thermo Scientific (Langerwehe, Germany). Two-milliliter polypropylene LoBind® test tubes were from Eppendorf AG (Hamburg, Germany, #022431102). Five hundred-microliter volume Amicon Ultra centrifugal filter devices with 3-kDa cutoff were from Millipore (Billerica, MA, USA).

Preparation of human milk samples and MRM LC-ESI-MS² analyses of HMOs

Pre-treatment of human milk samples, targeted semi-quantitative negative ion MRM LC-ESI-MS² analyses of HMOs, software settings, evaluation of data, and manual assignment of human milk groups were essentially conducted as recently published in more detail by Mank et al. [73]. Briefly, preparation of HM milk samples was accomplished by adding 15 μl of α -L-arabinopentaose internal standard (ITS) solution (0.05 mmol/l) to 135 μl of human milk. Frozen human milk samples had been thawed before at room temperature and were vortexed well. The combined sample-ITS-solution was further diluted 1:11 (v/v) in 2-ml Eppendorf test tubes by adding 1350 μl H₂O (LC-MS grade). Four hundred fifty microliters of the diluted mixture was transferred into a previously washed 500- μl Amicon Ultra centrifugal filter device with 3-kDa cutoff. Ultrafiltration (UF) was performed at 14,000g for 1 h. The UF permeate was transferred into a 300- μl glass LC-MS vial with screw top (Thermo Fisher Scientific, Waltham, MA, USA) and either stored at $-20\text{ }^{\circ}\text{C}$ until further use or directly subjected to LC-MS analysis. Sample preparation steps were conducted at room temperature.

For negative ion mode multiple reaction monitoring LC-ESI-MS² analyses, a 1100 series HPLC stack (Agilent, Santa Clara, CA, USA) with a solvent rack, degasser, binary solvent pump, high-performance autosampler, column oven, and DAD detector was coupled with a 3200 QTRAP triple

quadrupole linear ion trap MS instrument (Sciex, Framingham, MA, USA). A Hypercarb 2.1 \times 30 mm porous graphitized carbon (PGC) column (5- μm particle diameter) following a 2.1 \times 10 mm Hypercarb pre-column (both Thermo Fisher Scientific, Waltham, MA USA) served as the stationary phase. Column temperature was kept at 45 $^{\circ}\text{C}$, and sample temperature in the autosampler was 22 $^{\circ}\text{C}$. The DAD detector was used to monitor possible peptide or protein contaminations at 215- and 280-nm wavelengths. Five-microliter sample solutions prepared as described above were injected into the LC-MS system. Carry over was minimized by flushing the sample needle externally for 5 s with 20% aqueous isopropanol after each injection and by running a water and isopropanol blank between each HM sample. Elution of HMOs was achieved with a \sim 18 min H₂O-EtOH gradient at a flow rate of 400 $\mu\text{l}/\text{min}$ (see Table S1 in Electronic Supplementary Material, ESM). The solvents used to form the gradient were A (5 mM aqueous NH₄CH₃COO) and B (80% EtOH (v/v) with 5 mM aqueous NH₄CH₃COO). Analyst 1.4 software (Sciex, Framingham, USA) was used to control the complete LC-MS configuration and for further evaluation of LC-MS data.

Using this LC-MS platform, a first set of 10 distinct, specific, and partly isomeric HMO structures could be integrated into the targeted MRM LC-MS² method. HMO-specific MRM transitions were based on monoisotopic molecular masses of deprotonated precursor ions and respective diagnostic fragment ions. Unit resolution was chosen to conduct MRM analyses. MRM transitions were continuously monitored in negative ion mode during the entire MRM LC-MS² gradient elution (see [73]). One LC-MS run took approximately 18 min. HMO structures which were determined by this first set of (isomer-) specific MRM transitions included 2'-FL, 3-FL, 3'-SL, 6'-SL, DFL, LNT, LNnT, LNFP I, LNFP II, and LNFP III (see Table S2 in ESM). The correct identification of the latter HMO structures via their respective MRM transitions was proven by analysis of pure HMO standards (concentrations \sim 0.03 to 0.05 mg/ml) which were run together with the INFAT human milk study samples. The used pure HMO standards had been isolated from pooled human milk and were identical to those already described in more detail by Mank et al. 2019 [73].

Furthermore, a second set of 11 additional MRM transitions representing further OS structures could be included into the same LC-MS² approach as mentioned before (see Table S3 in ESM). Compared with the first MRM set, HMO monosaccharide compositions rather than exact structural features could be directly deduced from the fragment ions used in these MRM transitions (see [73]). Therefore, either one HMO composition or up to two different structures were assigned per MRM transition in set 2. Compounds which could be characterized by this 2nd set of MRM transitions were (Hex)₂ or lactose, (Hex)₃ or galactosyllactose, sialyllactose or (Hex)₂ (NeuAc)₁, (Hex)₄, α -arabinopentaose (internal

standard), tentative blood group B-tetrasaccharide or (Hex)₃(Fuc)₁, tentative blood group A-tetrasaccharide or (Hex)₂(HexNAc)₁(Fuc)₁, LNFP V or LNFP VI, LNDFH I or LNDFH II, and LNDFH II or LNDFH II. Despite the more putative assignment of the latter HMOs, at least LN(n)DFH I could also be distinguished from LN(n)DFH II due to a retention time delta of approximately 2 min as shown in Mank et al. 2019 [73]. All 60 human milk samples derived from the INFAT study were analyzed by the MRM LC-ESI-MS² approach described above in a randomized and anonymized manner (for exemplary MRM LC-ESI-MS² traces of some pure HMO standards (LNFP I, II, III, LNT, and LNnT) and of abundant HMOs in a HM study sample, please see Fig. S1a–f) in ESM).

If not otherwise mentioned, the settings for identification of HMOs, peak integration, and relative quantitation of HMOs using negative ion mode MRM LC-ESI-MS have been applied as described by Mank et al. [73]. Also, run to run variations for individual HMOs were determined as described in this reference by 10 repetitive injections of the same HM sample. Results were expressed as individual relative standard deviations (RSDs) for distinct HMOs: The resulting RSDs were less than 12% for LNFP I, LNFP II, LNFP III, LNT, and LNnT, up to 20% for total SL, (Hex)₃, 3'-SL, 6'-SL, and (Hex)₂/Lac, and maximally 38% for all other targeted HMOs. After integration of peak areas, dimensionless responses *R* were calculated for each HMO peak in a MRM trace by dividing its area under the curve by the area under the curve yielded for the internal standard α -arabinopentaose. The limit of detection (LOD) for individual HMOs was defined by a signal to noise (*S/N*) ratio of approximately 3:1 [76]. Peaks below LOD were not considered and reported as zero values.

For further processing and evaluation, original LC-MS raw data sets were exported from Analyst 1.4 software to Microsoft Excel version 14.0.7145.5000 (32 bit), Sciex Markerview version 1.2.1., or IBM SPSS version 19. Basic statistics (*t* test, Welsch's *t* test) were carried out within Markerview. Further statistical tests (see below) were accomplished with SPSS.

Manual assignment of HM samples to HM groups

Manual assignment of human milk INFAT study samples to one of the four commonly known human milk groups was based on the absence, presence, or concomitance of LNFP I and LNFP II. Biosynthesis of these two HMOs and other HMOs is governed by the maternal Se/Le status (see Table 1). If both LNFP I and LNFP II were detected in a sample, human milk group I was assigned. If LNFP I was absent but LNFP II was present, milk group II was assigned. If only LNFP I was detected but not LNFP II, group III was assigned and if neither LNFP I nor LNFP II could be detected, then milk group IV was assigned. Peaks were considered for

evaluation if the detected responses *R* exceeded the value of 0.2. The distribution of HM samples across human milk groups I, II, III, and IV was expressed in percentages and graphically visualized using SPSS software.

Automated HM group determination by principal component analyses (PCA) and statistical evaluation of lactation and HM group-specific HMO features

To relief future milk grouping attempts from tedious, time-consuming manual inspection of analytical LC-MS raw data, principal component analysis (PCA) was used as a means to automatically recognize different milk groups. Dedicated software (Markerview version 1.2.1. (Sciex)) facilitated PCA of 60 MRM LC-ESI-MS² HMO data sets derived from analyses of 2 × 30 HM samples (30 samples from 6 weeks and 30 samples from 16 weeks pp). The following parameters were used by the Markerview software to extract peak areas from MRM LC-ESI-MS² raw data: smoothing half width, 9 points; noise percentage, 50.0%; baseline subtraction window, 2.0 min; peak splitting factor, 2 points; retention time tolerance, 15.00 min; minimum peak intensity, 100 cps; minimum peak width, 4 points; minimum signal/noise ratio, 3:1. Prior to performing PCA, MRM LC-MS² peak areas of specific HMOs were first normalized against corresponding peak areas of the internal standard α -arabinopentaose. Then, normalized peak areas were converted into their natural logarithms and finally scaled by mean centering. Subsequently, PCA was performed. To investigate if abundances of individual HMO structures differed significantly between human milk groups as identified by PCA, non-parametric Kruskal-Wallis *H* testing [77] was performed with SPSS statistics software version 19.0.0 (IBM, Armonk, USA). Non-parametric Kruskal-Wallis *H* testing was applied as prior exploration of data sets by the Shapiro-Wilk test [78] indicated non-normal distribution of HMOs. The null hypothesis for non-parametric Kruskal-Wallis *H* testing was that medians of detected responses *R* for particular HMOs were the same between the milk groups. The considered significance level α was 0.05 and the confidence interval was 95%. The same LC-ESI-MS² HMO raw data sets as for PCA were used, but quality of integration was carefully inspected and manually re-adjusted if peaks were not fully integrated using automatic integration. Welch's *t* test [79] was applied to investigate if significant differences between HMO peak areas (normalized against the internal α -arabinopentaose) were existing between major HM groups and possible subgroups. Significance was reported if calculated *P* values were < 0.05 (α level of 0.05).

Moreover, variations in HMO abundances between HM samples from either 6 or 16 weeks pp were statistically elucidated by using Student's *t* test [80, 81] and Welch's *t* test. Prior to *t* testing, all data were interrogated by Levene's test for homoscedasticity [82]. Peak areas normalized against the

internal α -arabinopentose were used as input for calculations. Significance was stated if calculated P values were < 0.05 (α level of 0.05). Only HMO structures were considered which exhibited fold changes above 0.1 between the decadic logarithms of arithmetic mean at 6 compared with 16 weeks pp. For HMOs showing significant variations under these conditions, the differences between arithmetic mean values between 6 and 16 weeks pp of individual HMOs were expressed in percentages relative to the 6-week values. Results were displayed as bar charts created with Microsoft Excel for Office 365.

Results and discussion

Our MRM LC-ESI-MS² approach with enhanced specificity for fucose positional HMO isomers was applied to 60 unknown human milk samples. These milks originated from 30 donors resident of southern Germany which were enrolled in the INFAT study [74]. Specimens were randomly selected from a larger control group of women not subjected to dietary

intervention during the study. Per donor, two human milk samples from two different stages of lactation (6 and 16 weeks post-partum) were analyzed.

Figure 2 and Figure 3 summarize the resulting MRM LC-ESI-MS² responses R (see “Experimental section”) for HMO structures as detected across the 60 studied HM samples. These major HMO structures cover approximately $> 50\%$ of all HMOs contained in HM by quantity. It is noteworthy to mention that the applied semi-quantitative MRM LC-ESI-MS² approach does not yield equimolar responses for all investigated HMO structures. This is due to many factors including variations in ionization efficiency and fragmentation yields between specific HMOs. Nevertheless, relative differences in HMO concentration between samples or different stages of lactation can be accurately followed for individual HMO structures. Moreover, for some HMO isomers like LNFP I, II, and III, the MRM LC-ESI-MS² responses are similar enough to discriminate between biological abundances of these HMOs. This allows e.g. for reliable milk group assignment [73]. Also other HMOs like 3'-SL and 6'-SL show quite similar MRM LC-ESI-MS² responses which enable




Fig. 2 HMOs detected across 60 HM samples with LC-ESI-MS² ($N=30$ donors, 2 stages of lactation (6 weeks and 16 weeks) per donor). Responses R are not strictly equimolar and ranged between 0 and 132. HMO responses below LOD are reported with a zero value. Individual

milk donors are indicated by different capital letters A–EE, HM groups by Roman numbers I–IV, and stages of lactation by Roman numbers (VI= 6 weeks pp, XVI= 16 weeks pp)




Fig. 3 HMOs detected across 60 HM samples ($N = 30$ donors, 2 stages of lactation (6 weeks and 16 weeks) per donor) with MRM LC-ESI-MS². Responses R are not strictly equimolar and ranged between 0 and 7.7. HMO responses below LOD are reported with a zero value. Individual

milk donors are indicated by different capital letters A–EE, HM groups by Roman numbers I–IV, and stages of lactation by Roman numbers (VI = 6 weeks pp, XVI = 16 weeks pp)

reasonable direct comparisons of relative abundances. The same is true for LNT and LNnT.

Human milk groups shown in Fig. 2 and Fig. 3 were assigned following the “manual approach” (see “Experimental section”) based on the absence or presence of LNFP I and LNFP II. Hence, 77% of all HM samples (=23 donors) could be binned into HM group I, 13% (=4 donors) into HM group III, 7% (=2 donors) into HM group II, and 3% (=1 donor) into HM group IV. These proportions did not change when the PCA-based milk grouping approach was applied alternatively (see below).

Principal component analyses of HMO MRM LC-ESI-MS² data

As outlined in the “Experimental section”, yielded LC-MS data were also subjected to unsupervised

multivariate statistics using principal component analysis (PCA). This was done in an attempt to investigate three aspects:

1. the distinction of HM samples into (sub)groups not discovered by the manual approach,
2. the effect of progressing lactation on human milk group proportions between 6 and 16 weeks pp, and
3. the variations of Le/Se status-related or Le/Se status-independent HMOs between HM groups.

PCA-based grouping of HM milk samples

Figure 4 shows the resulting PCA score plot which could explain approximately 77% of variances in the MRM LC-ESI-MS² data set (PC 1, ~45%; PC 2, ~32%). Notably, four separate sample clusters distinguished from each other (blue, red, cyan, and orange dots). After post hoc inspection of LC-

Scores for PC1 (45,1 %) versus PC2 (31,8 %), Log | Mean Centre




Fig. 4 PCA evaluation of MRM LC-ESI-MS² data from 2 × 30 human milk samples (collected at 6 and 16 weeks pp). Every HM sample is represented by a dot in the PCA scoring plot depicted below. Blue and dark blue circles, cluster 1/group I human milks; red circles, cluster

2/group II human milks; cyan blue circles, cluster 3/group III human milks; and yellow circles, cluster 4/group IV human milks. Filled circles represent HM samples collected at 6 weeks pp. Open circles represent HM samples collected at 16 weeks pp

MS data sets for Se/Le gene-dependent milk group marker HMOs like 2'-FL, LNFP I, LNFP II, DFL, LNDH I, and LNDH II, each cluster could be unambiguously assigned to one of the 4 commonly known human milk groups (HM). Blue and dark blue circles in Fig. 4 represent samples belonging to HM group I, red circles HM group II, cyan circles milk group II, and orange circles milk group IV. Filled circles are HM samples collected at 6 weeks after delivery whereas open circles represent HM samples from 16 weeks pp.

Unexpectedly, within cluster I (blue circles, HM group I), a tendency towards formation of a possible subgroup (full and open dark blue circles in Fig. 4) became visible. We named this subgroup "HM group Ia." To investigate if responses of certain HMOs might differ significantly between HM group I and HM group Ia, we performed *t* testing as described in the "Experimental section." Thereby, the following findings were obtained: At 6 weeks post-partum, LNFP V/VI, LNFP II,

LNDFH II or LNnDFH II (C3-fragment), LNFP III, and 2'-FL varied significantly (all $P < 0.001$), and also 3'-SL ($P = 0.03$) and DFL ($P = 0.04$) were significantly different in abundance. At 16 weeks pp, LNFP V/VI, LNFP II, 3-FL, LNDFH II, or LNnDFH II (C3-fragment) exhibited significant abundance changes (all $P < 0.001$). Also levels of tentative blood group B-tetrasaccharide/(Hex)3(Fuc)1 ($P = 0.002$) and 2'-FL ($P = 0.003$) were found to be significantly different. In general, median values for Le+/FUT 3-related HMOs like LNFP II and LNDFH II were always lower in HM group Ia compared with HM group I. On the other hand, Se+-dependent HMOs like 2'-FL appeared in much higher levels in HM group Ia compared with HM group I.

The observed HM group Ia subset is concordant with similar findings from Dutch and Chinese human breast milks [54]. In 2018, Elwakiel et al. characterized HMO patterns derived from 30 HM from different Chinese donors via CE-LIF. Milk from

each donor was sampled between the first and 20th week after birth. For comparison, also 28 different Dutch breast milks which had been donated approximately at 4 weeks pp to a milk bank were analyzed. Hierarchical clustering of Chinese milk HMO patterns revealed a division of HM group I into two subgroups. These subgroups differed significantly in concentrations of 2'-FL, DFL, LNFP I, and LNFO. However, separation of Dutch Se+Le+ breast milks into subdivisions was driven by 2'-FL, LNT, and LNFO. Unlike in our study, 3-FL could not be evaluated as it was removed during sample pre-treatment. Also, no information was provided about the impact of LNFP II, LNFP III, LNFP V, or LNDFH II.

In sum, the “manual” approach for assignment of HM groups and the unsupervised PCA assignment resulted in the same 4 major milk groups with the same relative proportions of samples and also identical individual samples assigned to these major groups.

Figure 5 summarizes the resulting proportions of the 4 milk groups in percentages after characterization of HM samples by MRM LC-ESI-MS². HM group Ia (contributing 5% to the total HM group I + Ia) was not considered separately but accounted for in one HM group labeled as HM group I + Ia. This was done to be consistent with the number of milk groups published earlier [52] and to acknowledge the preliminary nature of HM group Ia identification: In our approach, this HM subgroup consisted of only 4 HM samples from 2 donors. Thus, future confirmation by analysis of much more HM

samples is warranted. The final PCA-based proportions of human milk groups (Fig. 5) were 77% (HM group I + Ia), 7% (HM group II), 13% (HM group III), and 3% (HM group IV). This result was identical to the proportions yielded by “manual” assignment relying on the presence or absence of LNFP I and LNFP II (see above). A comparable frequency of ~76% for human milk group I was reported by Tonon et al. in 2019 [83] for mature Brazilian human milks from 78 women. Noteworthy, the proportions for group II and group III of ~12% each differed from our results especially with regard to group II. This may be explained by regional differences or varying ethnicities between Germany and Brazil. Furthermore, in 2019, Samuel et al. [84] reported new findings for HM group distributions for milks collected during the first 4 months of lactation from 290 subjects across Europe (not including Germany). In their publication, 72% of donors could be assigned to HM group I, whereas ~18% of donors belonged to HM group II, ~7% to HM group 3, and ~4% to HM group IV. Again, differences between our findings and these numbers may be attributed to regional, ethnic, or yet unknown genetic or exogenous factors.

Similar proportions for human milk groups as found by us were also received by analyses of maternal Lewis blood group determinants directly in plasma of French (Oriol et al. 1986 [39]) or German (Thurl et al. in 2010 [16]) donors. Oriol reported values of 70% for Le (a-b+c-d-) phenotypes, 9% for Le (a-b-c-d+), and 1% for Le (a-b-c+d-). Thurl et al. tested




Fig. 5 HM group distribution in percentages based on MRM ESI-LCMS² analysis of 60 HM samples. Error bars represent 95% confidence intervals

blood specimen from 30 subjects but only for Le a and Le b epitopes and found the following distribution: 73% Le (a- b+), 17% Le (a+ b-), and 10% Le (a- b-). In general, the outcome of our study for HMOs and the results of Oriol et al. and Thurl et al. re-confirm the relationship of the Le blood group system and the maternal Le/Se status-dependent HM group system (see also Table 1). In this context, HM group II seems to be underrepresented in our data as only 7% of samples were assigned to this group. The corresponding Lewis blood group proportions published by Oriol et al. and Thurl et al. were 20% (Le a+b-c-d-) and 17% Le (a+b-), respectively. This difference may be attributed to the different geographies from which HM samples had been derived (Germany vs. France) or to different proportions of ethnicities in the probed sample sets. Also, variations introduced by the targeted biomolecules themselves (human milk oligosaccharides (this study) vs. erythrocytes) might be reasons for deviating findings. The set of fucosyltransferases (FUTs) affecting formation of Le epitopes on erythrocytes is related but not identical to the FUTs active in HM. Moreover, the relatively low number of 30 donors in our study might have contributed to the observed variations from results found by Oriol et al. in 1986, too. Therefore, in future, the parallel analysis of statistically meaningful numbers of blood and HM specimen from donors of the same geographies and ethnicities may be helpful to mount further evidence.

PCA-based investigation of effects related to progression of lactation

In the PCA score plot depicted in Fig. 4, only minor shifts in ordination of HM samples could be observed between sampling at either 6 (filled circles) or 16 weeks (open circles) after birth. On the contrary, the pronounced distances between HM group clusters I–IV did persist despite changes of lactational stages. Also the assignment of individual HM samples to HM groups I–IV stayed the same at 6 and 16 weeks pp. Consequently, no percentual change in milk group ratios could be attributed to progression of lactation. Thus, we conclude that at least between 6 and 16 weeks pp, progression of lactation has no influence on milk group distribution and milk group-related diversity (but not abundance) of particular HMOs.

Variation of Le/Se status-related or Le/Se status-independent HMOs between milk groups

The PCA loading plot displayed in Fig. 6 helped to reveal specific HMO compounds which were driving the separation into 4 major milk group clusters most: The highest impact on separation of milk groups was exerted by a set of MRM LC-MS² transitions specific for LNFP I, 2'-FL, and DFL (light blue rectangle). These HMOs are linked with a secretor-

positive (Se+) genetic maternal status. To a lower degree, also LNDFH I/LNnDFH I transitions (light red and blue rectangle), which are concatenated with a positive secretor and Lewis gene status (Se+/Le+), had a strong impact on HM grouping. HM group separation was less but still considerably influenced by MRM transitions of Le-positive (Le+) status-related HMOs like LNFP II and LNDFH II/LNnDFH II (light red rectangle). Intriguingly, another set of Se and Le gene-independent HMOs like LNFP V/VI, and LNnT, and also the tentative blood group A and B tetrasaccharides seemed to contribute to clustering as well, but to an even lesser extent than the Le/Se gene-linked glycans. Other compounds like LNFP III, lactose/(Hex)₂, (Hex)₃, and 6'-SL have almost no effect on group separation. This means that the latter structures are quite stable in presence and abundance across all probed HM samples. Variability of Le and Se gene-regulated HMOs is much higher.

The observations explained above were further substantiated by non-parametric statistics. The null hypothesis that distribution and abundance of particular HMOs are the same across the 4 human milk groups as distinguished by PCA was rejected for all measured Le/Se-dependent but also some Le/Se-independent HMOs by non-parametric statistics (Kruskal-Wallis method, see "Experimental section") as well. As a result, actually 12 HMO structures differed significantly in abundance between human milk groups I–IV (see Table 2). Moreover, some HMOs could only be detected in specific milk groups (see column 3 in Table 2 or Fig. 2 and Fig. 3). If HMOs were detectable in more than one milk group, the group with the highest median was highlighted by red bold letters in Table 2. For example, 2'-FL and LNFP I could both be detected in HM groups I and III but medians were higher in HM group III. Thus, Roman number III is displayed in bold red for both structures.

Noteworthy, HMOs related to a positive maternal genetic Se status (Se+), positive Le status (Le+), or positive Se and Le status (Se+Le+) such as 2'-FL, DFL, LNFP I, LNFP II, LN(n)DFH I, and LN(n)DFH II exhibited significantly different levels between milk groups (significance level $\alpha = 0.05$, see Table 2). The latter findings approximately corroborate data published e.g. by Thurl et al. in 2010 [16] for milk group-dependent abundances of distinct neutral, fucosylated HMOs.

This outcome complies with the well-accepted biological expectation that distribution of specific HMO structures should differ in HM samples due to known differences in maternal Le/Se status which can occur between individual milk donors: If the maternal Secretor and/or Lewis gene is active, it will encode expression of fucosyltransferases FUT 2 and/or FUT 3 and subsequent synthesis of either α 1,2- and/or α 1,4-fucosylated HMOs in the mammary gland.




Fig. 6 a PCA loading plot based on LC-ESI-MS² responses *R* for 21 MRM transitions of individual, partly isomeric HMOs as detected in 2 × 30 human milk samples collected at 6 and 16 weeks pp. HMO-specific MRM transitions are represented by individual dots including

information about precursor masses, fragment masses, and retention times. The order number of transition in MRM scan is reported in parentheses. **b** Magnification of gray shaded section in (a) depicting MRM transitions of HMOs with no or low contribution to clustering of HM

Table 2 Milk group-specific presence of individual HMOs. *P* values < α indicate statistically significant differences between milk groups I, II, III, and IV found for median values of 12 particular HMOs (nos. 1–12, gray shaded). $\alpha = 0.05$, non-parametric Kruskal-Wallis test, $N = 2 \times 30$

samples (from 6 and 16 weeks pp). Bold red letters in column 3 indicate elevated median values of specific HMOs in this particular milk group relative to other milk groups

No.	HMOs	Presence in milk group	P value
1	LNDFHI/LNnDFHI, (MRM transition C ₂ -fragment)	I+III	0.000
2	LNFP II	I+II	0.000
3	LNDFHI/LNnDFHI, (MRM transition C ₄ -fragment)	I	0.000
4	LNDFHII/LNnDFHII, (MRM transition C ₃ -fragment)	I+II	0.000
5	LNFPV/LNnFPV	I, II, III, IV	0.000
6	DFL	I+III	0.000
7	LNFP I	I+III	0.000
8	2'-FL	I+III	0.000
9	Tentative Blood Group B-tetrasaccharide/(Hex) ₃ (Fuc) ₁	I, II, III, IV	0.001
10	3-FL	I, II, III, IV	0.004
11	LNnT	I, II, III, IV	0.014
12	Tentative Blood Group A-tetrasaccharide/(Hex) ₂ (HexNAc) ₁ (Fuc) ₁	I, III	0.032
13	(Hex) ₃ /e.g. galactosyllactose	I, II, III, IV	0.158
14	(Hex) ₂ /Lac	I, II, III, IV	0.191
15	LNFP III	I, II, III, IV	0.213
16	LNT	I, II, III, IV	0.256
17	3'-SL	I, II, III, IV	0.284
18	Total SL	I, II, III, IV	0.542
19	6'-SL	I, II, III, IV	0.668
20	(Hex) ₄	I, II, III, IV	0.899
21	α -Arabinopentaose (internal standard)	I, II, III, IV	1.000

As a result, the presence or absence of α 1,2- and/or α 1,4-fucosylated human milk OS should lead to formation of 4 different milk groups defined by presence (or absence) of α 1,2- and/or α 1,4-fucosylated HMOs. As mentioned before,

not only the expected appearance of these different milk groups but also significant variations in α 1,2- and α 1,4-fucosylated oligosaccharides between the 4 HM groups could be confirmed after interrogation of the LC-MS results through

Table 3 Isomeric LNDFH structures, possible tandem MS fragment ion types, and related MRM transitions as utilized in the applied HMO LC-ESI-MS² method

No.	Oligosaccharide	Oligosaccharide Structure	MRM-Transitions		Possible Assignments Fragment ion types
			Precursor [m/z]	Fragment [m/z]	
1	LNDFH I or LNnDFH I		998.3569	325	
2	LNDFH I or LNnDFH I		998.3569	836	
3	LNDFH II or LNnDFH II		998.3569	690	

PCA and non-parametric statistics (see Table 2). More surprisingly and in addition, PCA and non-parametric statistical testing suggested that some HMOs such as α 1,3-fucosylated 3-FL and LN(n)FP V or e.g. non-fucosylated LNT which are all independent from maternal Le/Se status did also significantly vary between HM groups.

A further examination of milk groups for specific HMO structures revealed another unexpected finding: the appearance of the MRM signal for LNDFH I based on its C₂ fragment (see Table 3) in milk group III samples (with lower abundance than in milk group I). This finding is in contradiction with the published knowledge describing a lack or




Fig. 7 Variations of individual HMO levels between stages of lactation. **a** *P* values for variations in HMO mean values between 6 and 16 weeks pp were plotted over respective fold changes. Negative values for fold changes indicate an increase of HMO levels over time, and positive values decrease. **b** Section of (**a**) displaying only significantly changing

HMOs ($P < \alpha$). Red diamonds, Le⁺-dependent HMOs; blue diamonds, Se⁺-dependent HMOs; purple diamonds, Se⁺- and Le⁺-dependent HMOs; gray dots, Le/Se-independent HMOs; and black triangle, internal standard alpha-arabinopentaose




Fig. 8 Quantitative differences of HMO mean values with significant level changes (t test, $P < 0.05$) between 6 and 16 weeks pp. Level changes are expressed in percentages relative to the mean value at

6 weeks pp. Highest increase over time is found for 3-FL (57%), highest decrease for 6'-SL (-73%), followed by LNFP I (-37%)

strongly reduced activity of FUT 3 for Le-negative HM group III donors. FUT III would be needed in biosynthesis to append a fucose in $\alpha 1,4$ glycosidic linkage to the GlcNAc in the type I LNT backbone of LNDFH I. An explanation for this paradox may be provided by the fact that the used LNDFH I C₂-fragment-based MRM transition obviously picks up both the backbone type I and type II isomers of LNDFH I. Therefore, the LNDFH I variant determined in the group III milks is most probably LNneoDFH I, also named LNDFH III. LNDFH III consists of a type II LNnT backbone and the $\alpha 1,4$ -linked fucose present in LNDFH I is replaced by a $\alpha 1,3$ -linked fucose. The latter structure is FUT 3 independent and could therefore be present also in HM group III samples. In contrast, LNDFH I (with type I backbone) is detected only in group I milks as expected if the MRM transition of the LNDFH C₄-fragment is monitored. Thus, the latter transition is obviously more specific and suitable for correct identification of LNDFH I.

Relative HMO abundance changes between 2 stages of lactation (6 and 16 weeks post-partum)

Another important aspect which may contribute to functional adaptation of human milk during staging of lactation is the change of concentrations observed for particular HMOs. To further investigate such staging effects, we compared the human milks which were either collected at 6 or 16 weeks after parturition. This was done with no distinction between milk groups.

As shown in Fig. 7a and b, only six HMO structures were changing significantly in abundance between the lactational stages at 6 and 16 weeks post-partum (α level of 0.05; fold change threshold of ± 0.1). This finding could be confirmed for 6'-SL, 3-FL, LNFP I, and (Hex)₃ also with Wilcoxon signed-rank test for non-normally distributed, interdependent samples (data not shown). Among these significantly changing HMOs, only one glycan (LNFP I) was related to maternal Se+ status. LNFP I decreased in its relative abundance. Additional HMO compounds including 6'-SL, total SL, (Hex)₃, and LNT showed the same tendency and decreased. In opposition to this trend, 3-FL exhibited an increase in relative abundance between 6 and 16 weeks pp. Except for LNFP I, all the other 5 HMOs were independent from maternal Le or Se status.

Figure 8 visualizes quantitative abundance changes for the six selected HMO structures visible in Fig. 7b. Level changes were expressed in percentages relative to the original abundances measured at 6 weeks pp. Here, 6'-SL showed the strongest quantitative variation among all investigated HMOs with a decrease of 73%. On the contrary, 3'-FL increased by approximately 57% during the same time span. Interestingly, in 2016, Seppo et al. [85] published similar data with highly significant shifts of 6'-SL (decrease) and 3-FL (increase) between 0 and 6 months pp. They also showed a negative correlation between LNFP I concentration and increasing duration of breastfeeding. In contrast to our analytical LC-MS-based approach, these authors relied on HPLC analysis of HMOs after labelling with 2-aminobenzamide. Also other reports state a steady increase for 3'-FL levels either between approximately 0 and 8 months pp for HM collected from Chinese women [86] or (at least) between 0 and 4 months pp for

HM from Pan-European populations [84]. Furthermore, Austin et al. could show that 2'-FL, 3'-SL, 6'-SL, LNT, LNnT, LNFP-I, and LNFP-V decreased by trend over lactational stages. Although we analyzed human milks from a different geography (Germany), these trends were reflected in our data, too. Another publication by Coppa et al. [18] was in line with our results as well. In the latter study, 3'-SL decreased between 0 and 3 months pp in HM, but the differences in 3'-SL were not significant. Also the 6'-SL concentration declined over the same time range, but in this case and similar to our findings, the concentration changes were significant.

Conclusions

In this study, application of a novel LC-MRM-MS concept to 2×30 mature human milk samples collected at 6 and 16 weeks post-partum was demonstrated. This approach covered (partly isomeric) HMOs up to hexasaccharides and as a novelty also the (tentatively assigned) blood group A and B tetrasaccharides. More than approximately 50% of the HMO quantity present in HM could be characterized.

Principal component analysis (PCA) was successfully utilized for automated interrogation of LC-MS data. The chosen PCA approach allowed for unattended, effortless, and quick assignment of HM samples to one of the 4 currently known human milk groups. The additional advantage of PCA on top of advanced data processing speed and convenience was the immediate visualization and indication for possible HM subgroups. Taking these findings into account, unsupervised PCA seems to be well suited for convenient HM group recognition if applied as described in the “[Experimental section](#)”.

Intriguingly, mining of HMO LC-MS data via PCA and comparison with complementary non-parametric statistics revealed first evidence for possible distinction of HM group I into a further HM subgroup termed HM group Ia. Compared with other HM group I samples, HM group Ia specimen had significantly elevated levels of Se+-related HMOs (e.g., 2'-FL) and decreased levels of Le+ structures like LNFP II and LNDFH. Also Le/Se-independent HMOs such as LNFP V/VI or putatively assigned blood group B-tetrasaccharide/(Hex)₃(Fuc)₁ were found to differ.

Twelve HMO structures exhibited significantly different abundances between the major 4 HM groups. As expected and in line with contemporary literature, most of those 12 HMOs were related to either a positive maternal genetic Se status, a positive Le status, or a positive Se and Le status. However, also a second set of Le/Se status-independent α 1,3-fucosylated compounds like 3-FL and LN(n)FP V or non-fucosylated LNT did vary significantly.

During progression of lactation between 6 and 16 weeks pp, staging effects could be confirmed for 6 distinct

HMO structures. Most prominent level changes were revealed for 6'-SL (decrease by 73%) and 3-FL (increase by 57%).

It is challenging to find a rationale why there is a second set of Le/Se gene-independent HMOs, seemingly associated with milk grouping. Neither milk group-relevant active FUT2 or FUT3 genes are required for biosynthetic formation of those human milk glycans. Finally, and in accordance with first interpretation of PCA outcomes, Kruskal-Wallis ANOVA testing could confirm the existence of a third set of human milk oligosaccharides which were assessed to be permanently present across all samples without significant difference between the 4 human milk groups (see nos. 13–21 in Table 2).

Whether milk group-independent, ubiquitously present glycans like lactose/(Hex)₂, (Hex)₃, SL, LNT, or LNFP III can therefore be considered as more “essential” with regard to healthy infant development rather than the Le/Se milk group-dependent HMOs is yet still open. At least growing published evidence states their association with relevant early life health benefits like bifidogenic effect, anti-infective efficacy (e.g., directed against parasitic protozoans), and modulation of the developing immune system including less incidence of cows' milk allergy [68, 85, 87–91]. On the other hand, it may be hypothesized that the synthesis and secretion of the Le/Se-dependent HMOs like 2'-FL, DFL, LNFP I [92–96], or LNFP II [97] could also be interpreted as a complementary additional response of evolution to optimize protection of the infant against specific threats like pathogenic bacteria or viruses.

Diversity and abundance of these pathogens however probably did or does still vary by geography due to socioeconomic and environmental circumstances. Therefore, several tailor-made but geographically variable sets of HMOs (i.e., the Le/Se gene-regulated HMOs) may be required to optimally protect the infant with respect to geographically variable, specific early life threats. The profile of a HMO set fitting optimally to the infants needs may also depend on the infants' own Le/Se status. The latter aspect is important as the Le/Se status also co-defines glyco-epitopes expressed on gut epithelial linings or other inner surfaces [55]. Pathogens can adhere to these epitopes via glycan-specific receptors and subsequently cause infections e.g. of the gastrointestinal tract. An optimal free human milk OS pattern would therefore resemble all glycan epitopes possibly displayed by the host (the infant) to bacteria or viruses. Thereby, effective blocking of glycan receptors of these pathogens could be achieved and adhesion and subsequent infection could be avoided.

However, if pathogenic threats throughout human evolution exerted a selective pressure on HMO profiles, the evolutionary adaption should have manifested itself in geographically different Le/Se gene-dependent HMO patterns. These patterns may still be traceable today in diverse geographic regions or ethnicities. In this sense, some publications propose a possible link between history of human FUT 2

polymorphisms [98] or blood group variations with partly pathogen-driven selective pressures [99]. Moreover, there is at least a growing body of evidence published by e.g. Erney et al. [23] and other authors [100–103] that prevalence of the Se gene and related HMOs like 2'-FL differs in a geographically specific manner.

Whether the idea of “essential” HMOs which are always present across all human milk groups and other milk group-dependent “specialized” e.g. anti-infective HMOs can be substantiated by further studies remains to be elusive and an exciting question. The same applies for the influence of HMO abundances on infants’ healthy development which is significantly changing during lactational staging. In this publication, we investigated only the segment of human milk lactation between 6 and 16 weeks pp. An extended milk sampling including specimen from the first week until 6 or even 12 months pp and a much higher number of samples could reveal more insights in future.

Other biofluids than HM such as fecal slurries or plasma could probably also become amenable to sensitive MRM-LC-MS analyses by slight adaptations of the current sample pre-treatment procedure. For example, additional microfiltration steps or SPE could be exploited in addition or as an alternative to 3-kDa ultrafiltration. Thus in future, more holistic insights may be elicited by improved analyses with regard to systemic effects of HMOs on healthy infants’ development during the relevant first 1000 days of life.

Finally, we hope that the described multiple reaction monitoring (MRM) LC-ESI-MS² methodology might help to study HMO profiles of even larger human milk sample sets e.g. provided by prospective human milk cohorts. If combined with PCA, this in turn may contribute to reveal new associations between variations in HM glycan composition and their impact on healthy infant development and early life immunomodulation.

Acknowledgments We would like to thank P. Welsch and C. Pabst for their excellent technical assistance in human milk sample preparation and LC-MS analysis. We are also grateful to Bernadet Blijenberg who helped to improve quality of depicted HMO sketches and to Nana Bartke who facilitated the access to human milk study samples. In addition, we highly appreciate the contributions of S. Brunner who provided comprehensive background information about human milk samples from the INFAT study.

Compliance with ethical standards

The usage and analysis of the described human milk samples have been performed in accordance with ethical standards and guidelines as laid down in the Declaration of Helsinki. Ethical approval and written consents of donors were given as stated above for HM samples from the INFAT study.

Conflict of interest Marko Mank and Bernd Stahl are employees of Nutricia Research B.V., Utrecht, The Netherlands. Albert J.R. Heck is Professor at Utrecht University, Utrecht, The Netherlands. Hans Hauner is holding a professorship at the Technical University of Munich,

Munich, Germany. The present research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors. None of the authors have further conflicts of interest with regard to the content of this manuscript.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ballard O, Morrow AL. Human milk composition: nutrients and bioactive factors. *Pediatr Clin N Am*. 2013;60(1):49–74.
- Stam J, Sauer PJ, Boehm G. Can we define an infant's need from the composition of human milk? *Am J Clin Nutr*. 2013;98(2):521S–8S.
- Thurl S, Munzert M, Boehm G, Matthews C, Stahl B. Systematic review of the concentrations of oligosaccharides in human milk. *Nutr Rev*. 2017;75(11):920–33.
- Boehm G, Stahl B. 9 - Oligosaccharides. In: Mattila-Sandholm T, Saarela M, editors. *Functional Dairy Products*. Cambridge: Woodhead Publishing; 2003. p. 203–43.
- Viverge D, Grimmonprez L, Cassanas G, Bardet L, Bonnet H, Solere M. Variations of lactose and oligosaccharides in milk from women of blood types secretor A or H, secretor Lewis, and secretor H/nonsecretor Lewis during the course of lactation. *Ann Nutr Metab*. 1985;29(1):1–11.
- Montreuil J, Mullet S. Study of the variations of the glucide constituents of human milk during lactation. *Bull Soc Chim Biol (Paris)*. 1960;42:365–77.
- Kunz C, Rudloff S, Schad W, Braun D. Lactose-derived oligosaccharides in the milk of elephants: comparison with human milk. *Br J Nutr*. 1999;82(5):391–9.
- Hong Q, Ruhaak LR, Totten SM, Smilowitz JT, German JB, Lebrilla CB. Label-free absolute quantitation of oligosaccharides using multiple reaction monitoring. *Anal Chem*. 2014;86(5):2640–7.
- Urashima T, Kitaoka M, Terabayashi T, Fukuda K, Ohnishi M, Kobata A. Milk oligosaccharides. In: *Oligosaccharides: sources, properties and applications*: Nova Science Publishers, Inc.; 2011. p. 1–58.
- Wu S, Grimm R, German JB, Lebrilla CB. Annotation and structural analysis of sialylated human milk oligosaccharides. *J Proteome Res*. 2011;10(2):856–68.
- Wu S, Tao N, German JB, Grimm R, Lebrilla CB. Development of an annotated library of neutral human milk oligosaccharides. *J Proteome Res*. 2010;9(8):4138–51.
- Stahl B, Thurl S, Zeng J, Karas M, Hillenkamp F, Steup M, et al. Oligosaccharides from human milk as revealed by matrix-assisted laser desorption/ionization mass spectrometry. *Anal Biochem*. 1994;223(2):218–26.
- Xu G, Davis JC, Goonatilleke E, Smilowitz JT, German JB, Lebrilla CB. Absolute quantitation of human milk oligosaccharides reveals phenotypic variations during lactation. *J Nutr*. 2017;147(1):117–24.

14. Gabrielli O, Zampini L, Galeazzi T, Padella L, Santoro L, Peila C, et al. Preterm milk oligosaccharides during the first month of lactation. *Pediatrics*. 2011;128(6):e1520–31.
15. Chaturvedi P, Warren CD, Altaye M, Morrow AL, Ruiz-Palacios G, Pickering LK, et al. Fucosylated human milk oligosaccharides vary between individuals and over the course of lactation. *Glycobiology*. 2001;11(5):365–72.
16. Thurl S, Munzert M, Henker J, Boehm G, Muller-Werner B, Jelinek J, et al. Variation of human milk oligosaccharides in relation to milk groups and lactational periods. *Br J Nutr*. 2010;104(9):1261–71.
17. Binia A, Lefebvre G, Charpagne A, Vogel M, Kirsten T, Kiess W, et al. Time of lactation and maternal fucosyltransferase polymorphisms are main determinants of human milk oligosaccharide variation. *J Pediatr Gastroenterol Nutr*. 2018;66:871.
18. Coppa GV, Pierani P, Zampini L, Carloni I, Carlucci A, Gabrielli O. Oligosaccharides in human milk during different phases of lactation. *Acta Paediatr Suppl*. 1999;88(430):89–94.
19. Martin-Sosa S, Martin MJ, Garcia-Pardo LA, Hueso P. Sialyloligosaccharides in human and bovine milk and in infant formulas: variations with the progression of lactation. *J Dairy Sci*. 2003;86(1):52–9.
20. Nakamura T, Kawase H, Kimura K, Watanabe Y, Ohtani M, Arai I, et al. Concentrations of sialyloligosaccharides in bovine colostrum and milk during the prepartum and early lactation. *J Dairy Sci*. 2003;86(4):1315–20.
21. Tao N, DePeters EJ, German JB, Grimm R, Lebrilla CB. Variations in bovine milk oligosaccharides during early and middle lactation stages analyzed by high-performance liquid chromatography-chip/mass spectrometry. *J Dairy Sci*. 2009;92(7):2991–3001.
22. Martin-Ortiz A, Barile D, Salcedo J, Moreno FJ, Clemente A, Ruiz-Matute AI, et al. Changes in caprine milk oligosaccharides at different lactation stages analyzed by high performance liquid chromatography coupled to mass spectrometry. *J Agric Food Chem*. 2017;65(17):3523–31.
23. Erney RM, Malone WT, Skelding MB, Marcon AA, Kleman-Leyer KM, O’Ryan ML, et al. Variability of human milk neutral oligosaccharides in a diverse population. *J Pediatr Gastroenterol Nutr*. 2000;30(2):181–92.
24. Jantscher-Krenn E, Aigner J, Reiter B, Kofeler H, Csapo B, Desoye G, et al. Evidence of human milk oligosaccharides in maternal circulation already during pregnancy: a pilot study. *Am J Physiol Endocrinol Metab*. 2019;316(3):E347–E57.
25. Dotz V, Rudloff S, Blank D, Lochnit G, Geyer R, Kunz C. 13C-labeled oligosaccharides in breastfed infants’ urine: individual-, structure- and time-dependent differences in the excretion. *Glycobiology*. 2014;24(2):185–94.
26. Goehring KC, Marriage BJ, Oliver JS, Wilder JA, Barrett EG, Buck RH. Similar to those who are breastfed, infants fed a formula containing 2'-fucosyllactose have lower inflammatory cytokines in a randomized controlled trial. *J Nutr*. 2016;146(12):2559–66.
27. Marriage BJ, Buck RH, Goehring KC, Oliver JS, Williams JA. Infants fed a lower calorie formula with 2'FL show growth and 2'FL uptake like breast-fed infants. *J Pediatr Gastroenterol Nutr*. 2015;61(6):649–58.
28. Goehring KC, Kennedy AD, Prieto PA, Buck RH. Direct evidence for the presence of human milk oligosaccharides in the circulation of breastfed infants. *PLoS One*. 2014;9(7):e101692.
29. Wise A, Robertson B, Choudhury B, Rautava S, Isolauri E, Salminen S, et al. Infants are exposed to human milk oligosaccharides already in utero. *Front Pediatr*. 2018;6:270.
30. Mehra R, Kelly P. Milk oligosaccharides: structural and technological aspects. *Int Dairy J*. 2006;16(11):1334–40.
31. Tao N, DePeters EJ, Freeman S, German JB, Grimm R, Lebrilla CB. Bovine milk glycome. *J Dairy Sci*. 2008;91(10):3768–78.
32. Urashima T, Taufik E, Fukuda K, Asakuma S. Recent advances in studies on milk oligosaccharides of cows and other domestic farm animals. *Biosci Biotechnol Biochem*. 2013;77(3):455–66.
33. Costache M, Apoil PA, Cailleau A, Elmgren A, Larson G, Henry S, et al. Evolution of fucosyltransferase genes in vertebrates. *J Biol Chem*. 1997;272(47):29721–8.
34. Harduin-Lepers A, Vallejo-Ruiz V, Krzewinski-Recchi MA, Samyn-Petit B, Julien S, Delannoy P. The human sialyltransferase family. *Biochimie*. 2001;83(8):727–37.
35. Kumazaki T, Yoshida A. Biochemical evidence that secretor gene, Se, is a structural gene encoding a specific fucosyltransferase. *Proc Natl Acad Sci U S A*. 1984;81(13):4193–7.
36. Johnson PH, Watkins WM. Purification of the Lewis blood-group gene associated α -3/4-fucosyltransferase from human milk: an enzyme transferring fucose primarily to type 1 and lactose-based oligosaccharide chains. *Glycoconj J*. 1992;9(5):241–9.
37. Greenwell P. Blood group antigens: molecules seeking a function? *Glycoconj J*. 1997;14(2):159–73.
38. Urashima T, Hirabayashi J, Sato S, Kobata A. Human milk oligosaccharides as essential tools for basic and application studies on galectins. *Trends Glycosci Glycotechnol*. 2018;30(172):SE51–65.
39. Oriol R, Le Pendu J, Mollicone R. Genetics of ABO, H, Lewis, X and related antigens. *Vox Sang*. 1986;51(3):161–71.
40. Ceroni A, Maass K, Geyer H, Geyer R, Dell A, Haslam SM. GlycoWorkbench: a tool for the computer-assisted annotation of mass spectra of glycans. *J Proteome Res*. 2008;7(4):1650–9.
41. Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Marth JD, et al. Symbol nomenclature for glycan representation. *Proteomics*. 2009;9(24):5398–9.
42. Kobata A. Structures and application of oligosaccharides in human milk. *Proc Jpn Acad Ser B Phys Biol Sci*. 2010;86(7):731–47.
43. Sasaki K, Kurata-Miura K, Ujita M, Angata K, Nakagawa S, Sekine S, et al. Expression cloning of cDNA encoding a human beta-1,3-N-acetylglucosaminyltransferase that is essential for poly-N-acetylglucosamine synthesis. *Proc Natl Acad Sci U S A*. 1997;94(26):14294–9.
44. Almeida R, Amado M, David L, Levery SB, Holmes EH, Merckx G, et al. A family of human beta4-galactosyltransferases. Cloning and expression of two novel UDP-galactose:beta-n-acetylglucosamine beta1, 4-galactosyltransferases, beta4Gal-T2 and beta4Gal-T3. *J Biol Chem*. 1997;272(51):31979–91.
45. Isshiki S, Togayachi A, Kudo T, Nishihara S, Watanabe M, Kubota T, et al. Cloning, expression, and characterization of a novel UDP-galactose:beta-N-acetylglucosamine beta1,3-galactosyltransferase (beta3Gal-T5) responsible for synthesis of type 1 chain in colorectal and pancreatic epithelia and tumor cells derived therefrom. *J Biol Chem*. 1999;274(18):12499–507.
46. Urashima T, Asakuma S, Leo F, Fukuda K, Messer M, Oftedal OT. The predominance of type I oligosaccharides is a feature specific to human breast milk. *Adv Nutr*. 2012;3(3):473S–82S.
47. Ropp PA, Little MR, Cheng PW. Mucin biosynthesis: purification and characterization of a mucin beta 6N-acetylglucosaminyltransferase. *J Biol Chem*. 1991;266(35):23863–71.
48. Johnson PH, Donald AS, Feeney J, Watkins WM. Reassessment of the acceptor specificity and general properties of the Lewis blood-group gene associated alpha-3/4-fucosyltransferase purified from human milk. *Glycoconj J*. 1992;9(5):251–64.
49. Sumiyoshi W, Urashima T, Nakamura T, Arai I, Nagasawa T, Saito T, et al. Galactosylactoses in the milk of Japanese women: changes in concentration during the course of lactation. *Glycobiology*. 2004;14(11):1196.
50. Oriol R, Mollicone R, Cailleau A, Balanzino L, Breton C. Divergent evolution of fucosyltransferase genes from vertebrates, invertebrates, and bacteria. *Glycobiology*. 1999;9(4):323–34.

51. Bao Y, Chen C, Newburg DS. Quantification of neutral human milk oligosaccharides by graphitic carbon high-performance liquid chromatography with tandem mass spectrometry. *Anal Biochem.* 2013;433(1):28–35.
52. Thurl S, Henker J, Siegel M, Tovar K, Sawatzki G. Detection of four human milk groups with respect to Lewis blood group dependent oligosaccharides. *Glycoconj J.* 1997;14(7):795–9.
53. Coppa GV, Gabrielli O, Zampini L, Galeazzi T, Ficcidenti A, Padella L, et al. Oligosaccharides in 4 different milk groups, Bifidobacteria, and Ruminococcus obeum. *J Pediatr Gastroenterol Nutr.* 2011;53(1):80–7.
54. Elwakiel M, Hageman JA, Wang W, Szeto IM, van Goudoever JB, Hettinga KA, et al. Human milk oligosaccharides in colostrum and mature milk of Chinese mothers: Lewis positive secretor subgroups. *J Agric Food Chem.* 2018;66(27):7036–43.
55. Thomsson KA, Schulz BL, Packer NH, Karlsson NG. MUC5B glycosylation in human saliva reflects blood group and secretor status. *Glycobiology.* 2005;15(8):791–804.
56. Sarnesto A, Kohlin T, Hindsgaul O, Voegelé K, Blaszczyk-Thurin M, Thurin J. Purification of the beta-N-acetylglucosaminidase alpha 1—3-fucosyltransferase from human serum. *J Biol Chem.* 1992;267(4):2745–52.
57. Mollicone R, Gibaud A, Francois A, Ratcliffe M, Oriol R. Acceptor specificity and tissue distribution of three human alpha-3-fucosyltransferases. *Eur J Biochem.* 1990;191(1):169–76.
58. Hanfland P, Kordowicz M, Peter-Katalinic J, Pfannschmidt G, Crawford RJ, Graham HA, et al. Immunochemistry of the Lewis blood-group system: isolation and structures of Lewis-c active and related glycosphingolipids from the plasma of blood-group O Le(a-b-) nonsecretors. *Arch Biochem Biophys.* 1986;246(2):655–72.
59. Jantscher-Krenn E, Bode L. Human milk oligosaccharides and their potential benefits for the breast-fed neonate. *Minerva Pediatr.* 2012;64(1):83–99.
60. Musilova S, Rada V, Vlkova E, Bunesova V. Beneficial effects of human milk oligosaccharides on gut microbiota. *Benefic Microbes.* 2014;5(3):273–83.
61. Newburg DS. Neonatal protection by an innate immune system of human milk consisting of oligosaccharides and glycans. *J Anim Sci.* 2009;87(13 Suppl):26–34.
62. Kunz C, Rudloff S. Health promoting aspects of milk oligosaccharides. *Int Dairy J.* 2006;16(11):1341–6.
63. van Kooyk Y, Rabinovich GA. Protein-glycan interactions in the control of innate and adaptive immune responses. *Nat Immunol.* 2008;9(6):593–601.
64. Ayeche-Muruzabal V, van Stigt AH, Mank M, Willemsen LEM, Stahl B, Garssen J, et al. Diversity of human milk oligosaccharides and effects on early life immune development. *Front Pediatr.* 2018;6:239.
65. He Y, Lawlor NT, Newburg DS. Human milk components modulate toll-like receptor-mediated inflammation. *Adv Nutr.* 2016;7(1):102–11.
66. Newburg DS, Ko JS, Leone S, Nanthakumar NN. Human milk oligosaccharides and synthetic galactosyloligosaccharides contain 3'-, 4-, and 6'-galactosylactose and attenuate inflammation in human T84, NCM-460, and H4 cells and intestinal tissue *ex vivo*. *J Nutr.* 2016;146(2):358–67.
67. Newburg DS, He Y. Neonatal gut microbiota and human milk glycans cooperate to attenuate infection and inflammation. *Clin Obstet Gynecol.* 2015;58(4):814–26.
68. He Y, Liu S, Leone S, Newburg DS. Human colostrum oligosaccharides modulate major immunologic pathways of immature human intestine. *Mucosal Immunol.* 2014;7(6):1326–39.
69. He Y, Liu S, Kling DE, Leone S, Lawlor NT, Huang Y, et al. The human milk oligosaccharide 2'-fucosyllactose modulates CD14 expression in human enterocytes, thereby attenuating LPS-induced inflammation. *Gut.* 2016;65(1):33–46.
70. Kunz C, Kuntz S, Rudloff S. Bioactivity of human milk oligosaccharides. *Food Oligosaccharides: Wiley Blackwell;* 2014. p. 1–20.
71. Heggelund JE, Burschowsky D, Bjornestad VA, Hodnik V, Anderlüh G, Krenzel U. High-resolution crystal structures elucidate the molecular basis of cholera blood group dependence. *PLoS Pathog.* 2016;12(4):e1005567.
72. Heggelund JE, Varrot A, Imberty A, Krenzel U. Histo-blood group antigens as mediators of infections. *Curr Opin Struct Biol.* 2017;44:190–200.
73. Mank M, Welsch P, Heck AJR, Stahl B. Label-free targeted LC-ESI-MS(2) analysis of human milk oligosaccharides (HMOS) and related human milk groups with enhanced structural selectivity. *Anal Bioanal Chem.* 2019;411(1):231–50.
74. Much D, Brunner S, Vollhardt C, Schmid D, Sedlmeier EM, Bruderl M, et al. Breast milk fatty acid profile in relation to infant growth and body composition: results from the INFAT study. *Pediatr Res.* 2013;74(2):230–7.
75. Hauner H, Vollhardt C, Schneider KT, Zimmermann A, Schuster T, Amann-Gassner U. The impact of nutritional fatty acids during pregnancy and lactation on early human adipose tissue development. Rationale and design of the INFAT study. *Ann Nutr Metab.* 2009;54(2):97–103.
76. Brunetti B, Desimoni E. About estimating the limit of detection by the signal to noise approach. *Pharm. Anal. Acta.* 2015;06(04):355.
77. Kruskal WH, Wallis WA. Use of ranks in one-criterion variance analysis. *J Am Stat Assoc.* 1952;47(260):583–621.
78. Shapiro SS, Wilk MB. An analysis of variance test for normality (complete samples). *Biometrika.* 1965;52(3–4):591–611.
79. Welch BL. The generalisation of Student's problems when several different population variances are involved. *Biometrika.* 1947;34(1–2):28–35.
80. Student. The probable error of a mean. *Biometrika.* 1908;6(1):1–25.
81. Zabell SL. On Student's 1908 article "the probable error of a mean". *J Am Stat Assoc.* 2008;103(481):1–7.
82. Olkin I. In: Ghurye SG, Hoeffding W, Madow WG, Mann HB, editors. *Contribution to probability and statistics: essays in honor of Harold Hotelling.* Palo Alto: Stanford University Press; 1960.
83. Tonon KM, de Moraes MB, Abrão ACFV, Miranda A, Moraes TB. Maternal and infant factors associated with human milk oligosaccharides concentrations according to secretor and Lewis phenotypes. *Nutrients.* 2019;11(6):1358.
84. Samuel TM, Binia A, de Castro CA, Thakkar SK, Billeaud C, Agosti M, et al. Impact of maternal characteristics on human milk oligosaccharide composition over the first 4 months of lactation in a cohort of healthy European mothers. *Sci Rep.* 2019;9(1):11767.
85. Seppo AE, Autran CA, Bode L, Jarvinen KM. Human milk oligosaccharides and development of cow's milk allergy in infants. *J Allergy Clin Immunol.* 2017;139(2):708–11 e5.
86. Austin S, De Castro CA, Benet T, Hou Y, Sun H, Thakkar SK, et al. Temporal change of the content of 10 oligosaccharides in the milk of Chinese urban mothers. *Nutrients.* 2016;8(6):346.
87. Fuhrer A, Sprenger N, Kurakevich E, Borsig L, Chassard C, Hennet T. Milk sialyllactose influences colitis in mice through selective intestinal bacterial colonization. *J Exp Med.* 2010;207(13):2843–54.
88. Wada J, Ando T, Kiyohara M, Ashida H, Kitaoka M, Yamaguchi M, et al. Bifidobacterium bifidum lacto-N-biosidase, a critical enzyme for the degradation of human milk oligosaccharides with a type 1 structure. *Appl Environ Microbiol.* 2008;74(13):3996–4004.
89. Urashima T, Fukuda K, Messer M. Evolution of milk oligosaccharides and lactose: a hypothesis. *Animal.* 2012;6(3):369–74.

90. Bode L, Jantscher-Krenn E. Structure-function relationships of human milk oligosaccharides. *Adv Nutr*. 2012;3(3):383S–91S.
91. Atochina O, Dadara AA, Walker M, Harn DA. The immunomodulatory glycan LNFPIII initiates alternative activation of murine macrophages in vivo. *Immunology*. 2008;125(1):111–21.
92. Morrow AL, Ruiz-Palacios GM, Jiang X, Newburg DS. Human-milk glycans that inhibit pathogen binding protect breast-feeding infants against infectious diarrhea. *J Nutr*. 2005;135(5):1304–7.
93. Newburg D, Ruiz-Palacios G, Altaye M, Chaturvedi P, Meinen-Derr J, Guerrero M. Innate protection conferred by fucosylated oligosaccharides of human milk against diarrhea in breastfed infants. *Glycobiology*. 2004;14:253–63.
94. Ruiz-Palacios GM, Cervantes LE, Ramos P, Chavez-Munguia B, Newburg DS. *Campylobacter jejuni* binds intestinal H(O) antigen (Fuc alpha 1, 2Gal beta 1, 4GlcNAc), and fucosyloligosaccharides of human milk inhibit its binding and infection. *J Biol Chem*. 2003;278(16):14112–20.
95. Naarding MA, Ludwig IS, Groot F, Berkhout B, Geijtenbeek TB, Pollakis G, et al. Lewis X component in human milk binds DC-SIGN and inhibits HIV-1 transfer to CD4+ T lymphocytes. *J Clin Invest*. 2005;115(11):3256–64.
96. Hong P, Ninonuevo MR, Lee B, Lebrilla C, Bode L. Human milk oligosaccharides reduce HIV-1-gp120 binding to dendritic cell-specific ICAM3-grabbing non-integrin (DC-SIGN). *Br J Nutr*. 2009;101(4):482–6.
97. Wu AM, Wu JH, Singh T, Liu JH, Tsai MS, Gilboa-Garber N. Interactions of the fucose-specific *Pseudomonas aeruginosa* lectin, PA-IIL, with mammalian glycoconjugates bearing polyvalent Lewis(a) and ABH blood group glycotopes. *Biochimie*. 2006;88(10):1479–92.
98. Ferrer-Admetlla A, Sikora M, Laayouni H, Esteve A, Roubinet F, Blancher A, et al. A natural history of FUT2 polymorphism in humans. *Mol Biol Evol*. 2009;26(9):1993–2003.
99. Fumagalli M, Cagliani R, Pozzoli U, Riva S, Comi GP, Menozzi G, et al. Widespread balancing selection and pathogen-driven selection at blood group antigen genes. *Genome Res*. 2009;19(2):199–212.
100. Castanys-Munoz E, Martin MJ, Prieto PA. 2'-Fucosyllactose: an abundant, genetically determined soluble glycan present in human milk. *Nutr Rev*. 2013;71(12):773–89.
101. Musumeci M, Simapore J, D'Agata A, Sotgiu S, Musumeci S. Oligosaccharides in colostrum of Italian and Burkinabe women. *J Pediatr Gastroenterol Nutr*. 2006;43(3):372–8.
102. Sumiyoshi W, Urashima T, Nakamura T, Arai I, Saito T, Tsumura N, et al. Determination of each neutral oligosaccharide in the milk of Japanese women during the course of lactation. *Br J Nutr*. 2003;89(1):61–9.
103. Leo F, Asakuma S, Fukuda K, Senda A, Urashima T. Determination of sialyl and neutral oligosaccharide levels in transition and mature milks of Samoan women, using anthranilic derivatization followed by reverse phase high performance liquid chromatography. *Biosci Biotechnol Biochem*. 2010;74(2):298–303.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.