



The role of oxytocin in male and female reproductive behavior

J.G. Veening^{a,c,*}, T.R. de Jong^b, M.D. Waldinger^a, S.M. Korte^a, B. Olivier^a

^a Department of Psychopharmacology, Division of Pharmacology, University of Utrecht, Utrecht, The Netherlands

^b Department of Behavioral and Molecular Neurobiology, University of Regensburg, 93053 Regensburg, Germany

^c Department of Anatomy, Radboud University Medical Center, Nijmegen, The Netherlands

ARTICLE INFO

Article history:

Accepted 24 July 2014

Available online 1 August 2014

Keywords:

Oxytocin

Reproductive behavior

Maternal behavior

Paternal behavior

Ejaculation

Ventromedial hypothalamus

ABSTRACT

Oxytocin (OT) is a nonapeptide with an impressive variety of physiological functions. Among them, the 'prosocial' effects have been discussed in several recent reviews, but the direct effects on male and female sexual behavior did receive much less attention so far. As our contribution to honor the lifelong interest of Berend Olivier in the control mechanisms of sexual behavior, we decided to explore the role of OT in the present review. In the successive sections, some physiological mechanisms and the 'pair-bonding' effects of OT will be discussed, followed by sections about desire, female appetitive and copulatory behavior, including lordosis and orgasm. At the male side, the effects on erection and ejaculation are reviewed, followed by a section about 'premature ejaculation' and a possible role of OT in its treatment. In addition to OT, serotonin receives some attention as one of the main mechanisms controlling the effects of OT. In the succeeding sections, the importance of OT for 'the fruits of labor' is discussed, as it plays an important role in both maternal and paternal behavior. Finally, we pay attention to an intriguing brain area, the ventrolateral part of the ventromedial hypothalamic nucleus (VMHvl), apparently functioning in both sexual and aggressive behavior, which are at first view completely opposite behavioral systems.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

"There is no simple correspondence between an adaptive behavior and any single peptide" and "there is no necessary unitary relation between a limbic peptide and a single pattern of behavior" (Herbert, 1993). These statements were applied on a

wide range of neuropeptides, including oxytocin (OT), and remain just as true as they were about two decades ago. Despite our considerably increased knowledge about locations and mechanisms of OT-release, the physiological and behavioral effects of OT on reproductive behavior in mammals (including humans) remain a complex matter of "whispered secrets and public announcements" (Leng and Ludwig, 2008).

Oxytocin (OT) is a nonapeptide with a molecular weight of 1007 Da, isolated and characterized in 1953 (Du Vigneaud et al., 1953). OT is produced in magno- and parvocellular neurons of the paraventricular hypothalamic nucleus (PVH), in the supraoptic hypothalamic nucleus (SON) as well as in accessory hypothalamic neurons, located in between these nuclei as well as in the bed nucleus of the stria terminalis (BNST) and medial preoptic area (MPOA) and mostly surrounding hypothalamic blood vessels (Armstrong and Hatton, 1980; Armstrong et al., 1980; Bealer et al., 2010; Kelly and Swanson, 1980; Sawchenko and Swanson, 1982b; Swanson, 1987; Swanson and McKellar, 1979a). The magnocellular neurons of PVH and SON send their fibers to the posterior pituitary, where the content is released in the vasculature to enter the general circulation to get access to all receptive peripheral organs (Bargmann, 1949; Kelly and Swanson, 1980; Swanson, 1987). The magnocellular projections form the hypothalamo-neurohypophyseal tract, which bends laterally and ventrally around the fornix and ventromedial hypothalamic nucleus

Abbreviations: 5-HT, serotonin; 8-OH-DPAT, Dipropylamino-5,6,7,8-tetrahydro-naphthalen-1-ol; AOB, accessory olfactory bulb; AVP, vasopressin; BNST, bed nucleus of the stria terminalis; CNS, central nervous system; EB, estradiol benzoate; ER α -IR, estrogen receptor- α immuno-reactive; Fos-IR, Fos immuno-reactive; GABA, gamma aminobutyric acid; GnRH, gonadotropin-releasing-hormone; HAA, 'hypothalamic attack area'; HPA-axis, hypothalamus-pituitary-adrenal-axis; ICV, intra-cerebro-ventricular; IST, isotocin; LG, 'licking-and-grooming'; MEApd, medial amygdaloid nucleus, posterodorsal part; MEApv, medial amygdaloid nucleus, posteroventral part; MPOA, medial preoptic area; mRNA, messenger ribonucleic acid; OT, oxytocin; P, progesterone; PAG, periaqueductal gray; PAGdl, periaqueductal gray, dorsolateral part; PAGvl, periaqueductal gray, ventrolateral part; PE, premature ejaculation; PVH, paraventricular hypothalamic nucleus; SON, supraoptic hypothalamic nucleus; SNP, single nucleotide polymorphism; VCS, vaginocervical stimulation; VMH, ventromedial hypothalamic nucleus; VMHdl, ventromedial hypothalamic nucleus, dorsolateral part; VMHdm, ventromedial hypothalamic nucleus, dorsomedial part; VMHvl, ventromedial hypothalamic nucleus, ventrolateral part

* Correspondence to: Department of Anatomy (109), Radboud University Medical Center, P.O. Box 9101, 6500 HB, Nijmegen, The Netherlands. Tel.: +31 263813742.

E-mail address: jan.veening@radboudumc.nl (J.G. Veening).

<http://dx.doi.org/10.1016/j.ejphar.2014.07.045>

0014-2999/© 2014 Elsevier B.V. All rights reserved.

(VMH) before entering the internal lamina of the median eminence. The PVH, the SON and the accessory groups each contribute about one third of the fibers entering the posterior pituitary (Rhodes et al., 1981; Swanson, 1987). In addition to the pituitary-directed projections, the PVH projects to target areas within the central nervous system (CNS). It was long thought that these projections originated exclusively from parvocellular OT-neurons located in the dorsal PVH, however, some recent studies showed that magnocellular OT-neurons project to CNS-target areas as well (Knobloch et al., 2012; Ross et al., 2009; Ross and Young, 2009). The destinations of central OT-projections vary from the olfactory bulbs to the lumbosacral spinal cord and involves numerous limbic and brainstem regions (Buijs, 1978b; Buijs et al., 1978; Knobloch et al., 2012; Ono et al., 1978; Ross et al., 2009; Ross and Young, 2009; Sawchenko and Swanson, 1982b; Swanson and McKellar, 1979a; Yu et al., 1996b).

Concerning the functional aspects of the OT-neurons in PVH and SON, many studies have shown their involvement in a variety of physiological and behavioral functions. A major challenge in this field of research is the difficulty to manipulate central oxytocin neurotransmission, since both OT and the currently available OT-receptor antagonists do not readily cross the blood–brain barrier. A currently ongoing debate centers around the question as to whether peripherally or intranasally administered OT can affect central neurotransmission and thereby influence behavior. Recent studies and reviews have come to an affirmative conclusion and intranasally administered OT is currently applied in humans for the treatment of mental illness (Churchland and Winkelman, 2012; Kagerbauer et al., 2013; Neumann et al., 2013; Veening and Olivier, 2013).

Originally most applications for humans were in the clinical realm, in the form of treatments induce uterine contractions during labor or the milk flow in the lactation period, frequently by intranasal administration (Hendricks and Gabel, 1960; Huntingford, 1961). Later on OT was also applied for the treatment of other diseases and over the last decade the effects of OT in social interactions in humans ('mind-reading', trust, 'face-processing', autism and fear-reduction) are being extensively explored (Behnia et al., 2014; Koch et al., 2014), see (Veening and Olivier, 2013) for a recent review. In animal experiments the effects of OT on social and affiliative behavior were also evident (Insel, 1992) as were the effects on parental behavior, food intake, grooming behavior and pain relief (Lee et al., 2009; Neumann and Landgraf, 2012; Veening et al., 2010; Veening and Olivier, 2013). The behavioral effects of OT are 'prosocial' (Lukas et al., 2011; Shelley et al., 2006) and are not typical for humans and rodents, since similar 'nonapeptide-effects' have been observed in a variety of other species, from mollusks via teleosts and birds to primates, including man (Churchland and Winkelman, 2012; Insel, 2010; Knobloch and Grinevich, 2014; Snowdon et al., 2010).

From the obvious 'prosocial' effects of OT it is not a major step to study the effects of OT on male and female social behavior. A remarkable effect of OT was observed in voles and other rodents where monogamously living species were clearly different from the polygamous species in the density and distribution of OT-fibers and -receptors (Carter, 2014; Carter et al., 1995; Carter and Porges, 2013; Gimpl and Fahrenholz, 2001; Insel, 2010; Insel and Shapiro, 1992; Knobloch and Grinevich, 2014; Lee et al., 2009; Manning et al., 2012; Neumann, 2008; Neumann and Landgraf, 2012; Pedersen and Tomaszewski, 2012; Veenema and Neumann, 2008; Weisman et al., 2012; Witt et al., 1990; Young et al., 2005).

In our present review we will present an overview of the known effects of OT on male and female reproductive behavior. Concerning the physiological and pharmacological details of the neuronal circuitry, a large amount of additional information has become available over the last decades but many questions remain to be answered. For some recent reviews, see (Snoeren et al., 2013a, 2013b; Veening and Coolen, 2013; Veening et al., 2013).

2. Getting together: role of OT in pair bonding

In monogamous pairs, either 'romantic' human couples (Borrow and Cameron, 2012; Carter, 2014; Carter and Porges, 2013; Grewen et al., 2005; Light et al., 2005) or long-term attached pairs of rodents like prairie voles (Cho et al., 1999; Williams et al., 1992a; Williams et al., 1992b), the affiliative effects of OT have been extensively documented (Carter, 2014; Carter et al., 1995; Insel, 2010; Insel and Shapiro, 1992; Lukas et al., 2011; Neumann, 2008; Ross et al., 2009; Snowdon et al., 2010; Young et al., 2005). Specific differences in the distribution of OT-receptors have been observed between the monogamous prairie voles and polygamous species like montane voles (Lee et al., 2009; Lee et al., 2010; Ross et al., 2009; Ross and Young, 2009; Young et al., 2005) and analogous differences in the 'OT-systems' have been observed in other animals, including primates (Crockford et al., 2013; Lee et al., 2009; Moscovice and Ziegler, 2012; Pedersen and Tomaszewski, 2012; Schneiderman et al., 2012; Snowdon et al., 2010). Pair bonds can be very strong and losing the preferred partner may have detrimental effects on the surviving animal or human individual (Carter and Porges, 2013; Insel, 2010; Young et al., 2005; Zellner et al., 2011). Calling it 'love', either directed to a young or to a partner of the opposite sex, appears to be the most appropriate description (Komisaruk and Whipple, 1998). "Love is deeply biological ... and has a profound effect on our mental and physical state" (Carter and Porges, 2013), and OT is strongly involved in these feelings (balanced by AVP (Neumann and Landgraf, 2012; Pedersen and Boccia, 2006; Veenema and Neumann, 2008)). Panksepp included a role of OT in two of his 'Basic Emotional systems' (Panksepp, 2011) and noted the remarkable similarities between social attachment and addiction. "Attachment is a primary form of addiction, or perhaps more accurately, addiction is a deranged form of attachment" (Zellner et al., 2011).

From these data it is clear that the 'prosocial' effects of OT stimulate physical contacts and inter-individual interactions (Carter, 1992). These changes are also reflected in diminished fear and anxiety behavior (Lukas et al., 2011; McCarthy et al., 1992; Neumann and Landgraf, 2012), increased trust (Alvares et al., 2010; Baumgartner et al., 2008; Lukas et al., 2011) and at least in male individuals in increased risk-taking behavior (Kavaliers et al., 2012; Waldherr and Neumann, 2007).

3. Doing the deed: role of OT in female sexual behavior

Whereas the level of pair-bonding and affiliate behavior varies widely among mammalian species (and the associated neuroanatomy and dynamics of the OT system as well), male and female sexual behavior is much more stereotyped. Although the temporal pattern of copulation differs between species and even within rodent species (for example, rats are much faster to initiate and complete copulation than mice), all mammals display a similar list of sexual behaviors (for example mounts and intromissions), postures (for example lordosis) and physiological reflexes (for example erection and ejaculation). OT has been found to play a role in many aspects of these stereotyped sexual parameters.

3.1. Role of OT in desire and appetitive behavior

In the appetitive, precopulatory phase of the sexual sequence the male and female rodent start their interactions (Emery, 1986; Pfaus et al., 2000; Pfaus et al., 1999; Veening, 1975; Veening and Coolen, 2013; Veening et al., 2013). An estrous female will rapidly start to display soliciting or proceptive behavior, 'hopping' and 'darting', to attract the attention of the male, to induce erection and to elicit a successful ejaculatory series in the copulatory phase

(Emery, 1986; Pedersen and Boccia, 2006; Pfaus et al., 1999; Veening et al., 2013).

The precopulatory phase is characterized by the phenomenon of 'desire' (Giraldi et al., 2004; Giraldi et al., 2013; Pfaus, 2009). OT contributes to 'sexual desire' and expectancy of future reward (Bancroft, 2005; Borrow and Cameron, 2012; Pfaus, 2009) and in the human this desire is temporarily suppressed after parturition, affected by OT and amygdaloid influences (Rupp et al., 2013). Pfaus (Pfaus, 2009) developed a dual control-model of inhibitory and excitatory effects on sexual behavior, and showed that dopamine functions at the excitatory side of the balance, together with OT and the melanocortins, while 5-HT (mostly, see (Mendelson and Gorzalka, 1985)), opioids and endocannabinoids work at the inhibitory side of the model (Pfaus, 2009). Sexual arousal (Bancroft, 2005) and desire are naturally related to the 'rewards' expected later in the behavioral sequence and therefore the important role of dopamine does not come as a surprise (Baskerville and Douglas, 2010; Frohmader et al., 2010; Hull, 2011; Hull et al., 2004; Pfaus, 2009; Pfaus et al., 1995; Pfaus et al., 2012). In this behavioral phase learning and conditioning processes play a very important role (Pfaus et al., 2012) and providing the female with the opportunity to 'pace' the succeeding copulatory events, enhances the 'reward value' of the copulatory phase (Emery, 1986; Nyuyki et al., 2011; Pfaus et al., 2000; Pfaus et al., 1999).

In the precopulatory phase the peripheral levels of OT start rising in both sexes, which may contribute to the preparation of the genital organs for the coming ejaculation or 'orgasm' (Borrow and Cameron, 2012; Giraldi et al., 2004; Giraldi et al., 2013; Komisaruk and Sansone, 2003). The presence of OT is not necessary for the coordination and performance of sexual behavior of male and female rodents. Transgenic OT-knockout as well OT receptor-knockout mice showed normal patterns of sexual behavior and pregnancy. Only after parturition problems arose, because lactation was heavily disturbed and all young animals would have died within 24 h, without proper precautions (Lee et al., 2010; Nishimori et al., 1996; Young et al., 1996). Do we have to conclude then that OT does not play a role in sexual activities? No, certainly not!

3.2. Role of OT in female copulation – anatomical substrate

Lordosis is specific dorsiflexed posture, observed in rodents and many other mammals, displayed by the female during successive mounts, intromissions and ejaculation by the male. The female keeps a fixed standing position, for a variable time-period, usually with additional details like lateroflexion of the tail. For additional details: see (Veening et al., 2013). The behavior can be observed only when the female is in estrus, under the proper hormonal conditions (Coirini et al., 1989; McEwen, 1988; Schumacher et al., 1989). In other circumstances she will actively reject a male companion trying sexual interactions.

In the female, the pelvic organs involved in (pre-)copulatory behavior are provided with OT receptors and OT fibers descending into the lumbosacral parts of the spinal cord (Buijs, 1978a; Normandin and Murphy, 2011; Swanson and McKellar, 1979b). By this combination of peripheral and central effects the pelvic organs can be prepared for the copulatory activities with regard to lubrication, muscular contractility and pain suppression for the coming vagino-cervical distension (Devost et al., 2008; Gelez et al., 2010; Giraldi et al., 2004; Magon and Kalra, 2011; Moos and Richard, 1975; Peters et al., 1987; Sansone and Komisaruk, 2001; Wilson et al., 2009).

In addition to the pelvic nerve (Moody et al., 1994), the vagal nerve and the 'dorsal vagal complex' appear to play an important role in the release of OT (Borrow and Cameron, 2012; Komisaruk

and Sansone, 2003; Komisaruk et al., 2004; Moos and Richard, 1975; Whipple and Komisaruk, 2002) while complex autonomic adaptations occur in the copulatory phase (Wilson et al., 2009).

For a recent overview of the spinal and brain structures involved in the control of the lordosis posture, see (Veening et al., 2013). For the moment, we will focus on a few 'main' areas in which OT is known to play a significant role: the PVH, the VMH, especially its ventrolateral part (VMHvl), the periaqueductal gray (PAG) region and the medial preoptic area (MPOA).

The PVH is one of the main sources of OT with projections to numerous peripheral and central destinations. Neuronal activation after sexual activities has been observed in females occasionally in both parvo- and magnocellular neurons (Cameron and Erskine, 2003; Flanagan et al., 1993; Pfaus and Heeb, 1997; Witt and Insel, 1994; Yang and Voogt, 2002). In rat pups, the parvocellular PVH neurons were activated by perineal stimulation and their activation produced a transient 89% increase in OT-levels in the lumbosacral spinal cord (Lenz and Sengelaub, 2010), which suggests that oxytocin is involved in the development and sensitization of the perineal part of the body, potentially supporting future copulatory activities in the male and female rat.

Concerning the VMH, this hypothalamic brain region is a crucial part of the motor circuitry producing the lordotic posture (Calizo and Flanagan-Cato, 2002; Flanagan-Cato, 2011; Mathews and Edwards, 1977a; Pfaff and Sakuma, 1979a; Veening et al., 2013). The VMH has been subdivided in several parts, a dorsomedial part (VMHdm) and a ventrolateral part (VMHvl), separated by a cell-poor intermediate area, and surrounded by cell-poor zone, especially prominent along its ventrolateral side (Canteras et al., 1994; Fahrbach et al., 1989; Flanagan-Cato, 2011; Millhouse, 1973a, 1973b, 1978; Van Houten and Brawer, 1978a, 1978b). In male rats the VMHvl is at least 25% larger than in females (Dugger et al., 2007; Madeira et al., 2001; Matsumoto and Arai, 1983). Millhouse (1973b, 1978) observed different kinds of neurons and most of them show about 3 primary dendrites and a few secondary ones (Ferri and Flanagan-Cato, 2012). The longest primary dendrite extends into the ventrolateral zone surrounding the VMHvl (Ferri and Flanagan-Cato, 2012; Flanagan-Cato, 2011). The longest dendrites, extending into the lateral cell-poor zone, are contacted by ascending and descending components of the medial forebrain bundle, traversing the intermediate hypothalamic area (Geeraedts et al., 1990a, b; Nieuwenhuys et al., 1982; Veening et al., 1982). OT-containing fibers are especially numerous in this area, along the VMHvl, which play an essential role in the estrus related behavioral changes and plasticity observed in this part of the VMH (Flanagan-Cato, 2011).

In the VMHvl at least 3 different cell types have been described: PAG-projecting neurons, ER α -IR neurons and neurons showing Fos-IR after mating, with only minimal functional overlap (Flanagan-Cato, 2011). The epaxial muscles (involved in the execution the lordosis response), have been injected with a retrograde (pseudorabies) virus and the pattern of labeling showed that the 'core' of the VMH and the estrogen-receptive neurons are hardly involved in these spinal projections and that by far the majority of the labeled neurons were observed in VMHvl (Daniels and Flanagan-Cato, 2000; Daniels et al., 1999).

Lordosis responses can be induced by electrical stimulation or disrupted by lesions of the VMHvl and this brain area has numerous relationships, among them a set of strong, mostly reciprocal, connections with the PAG (for a recent overview see (Veening et al., 2013)). In addition, the VMH receives olfactory, somatosensory and genitosensory stimuli from respectively the olfactory system, the flank region and the vagina and clitoris (Bennett et al., 2002; Bueno and Pfaff, 1976; Georgescu et al., 2009; Kirkpatrick and Merrill, 2011; Marson, 1995; Pfaus et al., 2006; Pfaus et al., 1996). In several studies neural activation was

observed in the female VMH after sexual activities, mostly but not exclusively in its ventrolateral part, the VMHvl (Blaustein et al., 1994; Coolen et al., 1996; Flanagan et al., 1993; Pfau et al., 1993; Rowe and Erskine, 1993; Tetel et al., 1993; Wersinger et al., 1993).

Interestingly, epigenetic influences have been observed in the VMHvl when females reared by low-licking-and-grooming (LG) mothers were compared with females reared by high-LG mothers. In the latter group lordosis rating was lower and more defensive behaviors were observed during mating, a lower number of ER α -IR but more Fos-IR neurons were observed in the VMHvl after mating and the pseudo pregnancy rate after vaginocervical stimulation (VCS) was lower, compared to the high-LG group (Cameron et al., 2011). These differences may have effects on lordosis termination (see below) but generally favor the reproductive success of the low-LG-reared females.

Concerning the PAG, control of lordosis is among its most conspicuous functions (Arendash and Gorski, 1983; Harlan et al., 1983; Ogawa et al., 1991; Ogawa et al., 1992; Pfaff and Lewis, 1974; Sakuma and Pfaff, 1979). This control is performed in a close but complex cooperation with the ventromedial hypothalamic nucleus (VMH), especially its ventrolateral part (VMHvl) (Flanagan-Cato, 2011; Flanagan-Cato et al., 2006; Flanagan-Cato and McEwen, 1995; Pfaff and Sakuma, 1979a; Pfaff and Sakuma, 1979c).

Concerning the MPOA, this region shows neural activation (Fos-IR) in female rats after copulatory activities (Cameron and Erskine, 2003; Coolen et al., 1996; Coria-Avila and Pfau, 2007; Flanagan-Cato and McEwen, 1995; Kirkpatrick and Merrill, 2011; Oboh et al., 1995; Parada et al., 2010; Pfau et al., 1994; Pfau et al., 2006; Polston and Erskine, 1995; Wersinger et al., 1993; Yang and Voogt, 2002), and the gonadotropin-releasing hormone (GnRH) neurons seem to be included in this activation pattern (Pfau et al., 1994; Wu et al., 1992). The GnRH-fibers are descending along the mesencephalic aqueduct, where they are probably involved in the induction of lordosis via the surrounding PAG (Buma, 1989; Sakuma and Pfaff, 1980; Samson et al., 1980; Sirinathsinghji, 1985; Veening and Barendregt, 2010; Veening et al., 2012). The involvement of the MPOA in the hormonal control mechanisms was detected early (Freeman and Banks, 1980; Gray et al., 1978; Hayashi and Orimo, 1975; Leipheimer and Gallo, 1985; Moss and Foreman, 1976; Sirinathsinghji, 1986) and so were the stimulating effects of OT on lordosis, about a decade later (Arletti and Bertolini, 1985; Caldwell et al., 1989; Kow and Pfaff, 1988; Schulze and Gorzalka, 1991), (though this stimulation depended on the test situation chosen (Whitney, 1986)).

3.3. OT effects on lordosis and orgasm

OT injected in the cerebral ventricles (ICV) stimulates proceptive behavior (and the lordosis posture) while AVP has the opposite effect and tends to increase agonistic reactions towards the male (Pedersen and Boccia, 2006). Icv-infusion of OT-antagonists had an inhibitory effects on female sexual behavior in rats (Pedersen and Boccia, 2002), similar to the effects of vasopressin (Pedersen and Boccia, 2006) as well as of peripheral administration in primates (Boccia et al., 2007).

In rodents, OT may be dispensable for the induction of lordosis in mice (Lee et al., 2010; Nishimori et al., 1996) but in rats clear facilitating effects have been observed after both systemic (Arletti and Bertolini, 1985) and central (Caldwell et al., 1986; Gorzalka and Lester, 1987; Schumacher et al., 1989) administration of OT. Since peripheral OT does hardly enter the CNS (Mens et al., 1983; Robinson and Jones, 1982), neural mechanisms must be involved in the coordination, and these have been explored by peripheral lesions like pelvic neurectomy (Moody et al., 1994) and hysterectomy (Moody and Adler, 1995) and by VCS in sheep (Kendrick

et al., 1993) and rats (Coolen et al., 1996; Northrop and Erskine, 2008; Pfau and Heeb, 1997; Veening and Coolen, 1998).

The generally facilitating effects of OT on lordosis (Arletti and Bertolini, 1985; Kow et al., 1991; Witt and Insel, 1992) need some additional comments, because they were not observed under all circumstances in rats (Caldwell et al., 1989; Schulze and Gorzalka, 1991; 1992; Schumacher et al., 1991) or prairie voles (Witt et al., 1990). These authors showed that the sensitivity of the neural substrate for ICV OT depends on the light/dark schedule, with the stronger responses in the dark period (Schumacher et al., 1991), that the exact hypothalamic location of the OT-infusion is important (Caldwell et al., 1989) and may influence lordosis frequency and lordosis duration separately (Schulze and Gorzalka, 1991), and that infusion of OT in a lateral ventricle may have an inhibitory effect on lordosis, not observed after an identical infusion into the 3rd ventricle (Schulze and Gorzalka, 1992). For recent overviews of the possible effects of OT in the CSF, see (Veening et al., 2010; Veening and Olivier, 2013).

In female human studies, it was observed that plasma OT levels are variable over the menstrual cycle (Salonia et al., 2005), with lowest levels during the luteal phase. During sexual arousal plasma levels increase (Blaicher et al., 1999) to reach peak levels at the moment of orgasm with, in multiorgasmic women, a clear correlation between OT-levels and subjective orgasm intensity (Carmichael et al., 1994). Concerning the arousal and orgasm-induced elevated levels of OT, it seems to be “still unclear whether OT increases sexual arousal or is a natural byproduct of it” (Borrow and Cameron, 2012). The presence of OT-receptors on multiple organs, especially on the male and female genital organs (Gimpl and Fahrenholz, 2001) suggests a possibly ‘preparatory role’ of OT for the later and final phases of the copulatory process, like ejaculation and orgasm, ‘preparing’ all necessary muscular contractions and lubrication effects. On the other hand, it has been suggested that the high levels of OT play a role in ‘sexual satiety’ following orgasm by flooding the OT-receptors and producing desensitization (Caldwell, 2002).

The physiological and behavioral effects of OT are strongly linked to other hormones like estrogen (Caldwell et al., 1988; Caldwell et al., 1986; Coirini et al., 1989; McCarthy, 1995; McEwen, 1988), progesterone (Caldwell et al., 1994; Gorzalka and Lester, 1987; Schumacher et al., 1989; Witt and Insel, 1991), prolactin (Donner and Neumann, 2009; Egli et al., 2010; Harlan et al., 1983; Northrop and Erskine, 2008; Sobrinho, 1993) as well as other factors (Beinfeld, 2001; Carter et al., 2003; Dellovade et al., 2000; Kow and Pfaff, 1988; Pedersen and Boccia, 2006). A discussion of their interactions would go far beyond the scope of the present review and unraveling these requires sophisticated experiments, for instance, concerning prolactin (Northrop and Erskine, 2008). Vaginocervical stimulation (VCS), induced by 15 intromissions, induces prolactin release and pseudopregnancy in control female rats. Local infusion of an OT-antagonist in the VMHvl prevented this pseudopregnancy, leading to the suggestion that “OT release in the VMHvl mediates the effects of VCS on the induction of the prolactin secretion needed to establish pseudopregnancy” (Northrop and Erskine, 2008). This experiment brings us from the peripheral sensory stimulation, the VCS, directly towards one of the main controlling brain areas, the VMHvl, involved in the induction of the lordosis posture (see above). A single treatment with OT or an OT-antagonist in the first postnatal week of young rats resulted in long-term changes in the expression of ER α and OT in the mediobasal hypothalamus (Perry et al., 2009).

OT has been shown to be strongly involved in the induction of lordosis, if estrogen and progesterone are present (Davis et al., 1979; Pleim and Barfield, 1988; Pleim et al., 1989; Rubin and Barfield, 1983). The facilitating effects of OT were especially prominent in the VMHvl (McCarthy et al., 1994; Schulze and

Gorzalka, 1991; Schumacher et al., 1989), and the physiological effects of OT on hypothalamic brain slices were clearly enhanced by the addition estradiol (Booth et al., 2010; Kow et al., 1991).

For an extensive overview of the neurochemistry of the VMH, see Flanagan-Cato (2011). A few points have to be mentioned, however, to understand the plasticity of the VMHvl and certain behavioral transitions discussed in a later section. Many of the long, laterally extending dendrites of the ER α -IR neurons express OT-receptors (Ferri and Flanagan-Cato, 2012; Flanagan-Cato, 2011; Flanagan-Cato et al., 2001). As we have seen, both estrogen and OT have a facilitating effect on female sexual behavior and lordosis, but other factors play a role as well, like GABA-ergic interneurons and glutamate, present in the OT-fibers (Flanagan-Cato, 2011; Flanagan-Cato et al., 2001; McCarthy et al., 1995; McCarthy et al., 1990; McCarthy et al., 1991b). Estradiol treatment combined with OT-administration enhances neuronal firing in the VMHvl (Booth et al., 2010; Kow et al., 1991). "Ovarian hormones enhance OT production, the electrical activity of OT neurons, and up-regulate the level of OT receptors in the VMH" (Ferri and Flanagan-Cato, 2012; Theodosios, 2002). The fiber plexus lateral to the VMHvl plays an important role in these interactions, because there VMH-dendrites and OT-fibers actually meet (Griffin et al., 2010; Schumacher et al., 1990). Estradiol treatment in female rats increases spine density and axospinous synapses, while these parameters decrease in male rats after the same treatment (Carrer and Aoki, 1982; Frankfurt et al., 1990; Lewis et al., 1995; Sa and Madeira, 2005). Estradiol also caused a marked shortening of the laterally extending dendrites, which effect could be reversed by a sequential progesterone treatment (Griffin et al., 2010; Madeira et al., 2001). So, estradiol regulates the number of contact in the zone lateral to the VMHvl. Remarkably, however, the VMH-dendrites showing OT-labeling, probably as a result of OT-internalization, were excluded from this shortening effect and remained intact (Griffin et al., 2010). These OT-estradiol interactions appear to be clearly involved in the plasticity occurring during the estrous cycle (4 or 5 days in the female rat) with its concurrent changes in behavior, especially the performance of the lordotic posture. But there are more factors playing a role here e.g. glutamate. The OT-fibers co-release glutamate, also in the axon-terminals in the contact area, lateral to the VMHvl, and glutamate (-agonists) have an inhibiting effect on female sexual behavior (Georgescu and Pfaus, 2006a; Georgescu et al., 2009; McCarthy et al., 1991a; McCarthy et al., 1995). In addition, glutamatergic neurons are present in the VMH and glutamate-transporters have been observed in the short dendrites of the ER α -IR VMHvl neurons (Flanagan-Cato, 2011; Georgescu and Pfaus, 2006a; Georgescu et al., 2009; McCarthy et al., 1991a; McCarthy et al., 1995). The situation is apparently very complex, with a mix of excitatory and inhibitory factors released together, and additional research is absolutely necessary to clarify these complex anatomical and neurochemical relationships (Ferri and Flanagan-Cato, 2012; Flanagan-Cato, 2011; McCarthy et al., 1991a). These complex interactions are not always or necessarily 'pro-sexual' or 'pro-lordosis'! Some of the mechanisms working in the area lateral to the VMHvl may not promote lordosis, but prepare the female for the termination of the estrous period (see Section 5).

3.4. OT and male copulation – anatomical substrate

Oxytocin (OT) strongly affects male sexual behavior. Evidence from male rats and a few other species have indicated that sexual cues and behaviors stimulate central and peripheral OT neurotransmission which, in turn, facilitate copulation (Argiolas and Melis, 2013; Gil et al., 2013). The neurocircuitry responsible for these interactions has been investigated using various techniques, including tract-tracing studies and immunohistochemistry to

identify how oxytocin neurons are connected with reproductive tissue and measurement of the activation of OT neurons and OT release in response to sexual interactions.

Parvocellular OT neurons project directly to the spinal cord (Gerendai et al., 2001). A series of tract-tracing studies, in which pseudorabies viruses were injected into the corpus cavernosum, penile muscles, epididymis and prostate of male rats, showed that the PVN was connected to each type of tissue (Gerendai et al., 2001; Marson and McKenna, 1996; Marson et al., 1993; Orr and Marson, 1998; Tang et al., 1998) putatively via the sacral parasympathetic nucleus (Tang et al., 1999). Varicosities containing OT or neurophysin (a carrier protein indicating the presence of either OT or AVP (Breslow and Burman, 1990)) have been found in close apposition to neurons in the sacral parasympathetic nucleus, ventral horn, and dorsal gray commissure that directly or indirectly control the genitals (Tang et al., 1998; Veronneau-Longueville et al., 1999; Wagner and Clemens, 1993). The (putatively OTergic) paraventricular neurons projecting to the spinal cord also provide collateral projections to the nucleus paragigantocellularis in the brainstem, which exerts a tonic serotonergic inhibition of penile reflexes (Bancila et al., 2002; Normandin and Murphy, 2011). Conversely, it was found that predominantly magnocellular OT neurons in the caudal portion of the PVN receive genitosensory input from the penis (Normandin and Murphy, 2011).

Indeed, Fos-expression is increased in both parvocellular and magnocellular oxytocinergic neurons of the PVH and SON in response to sexual behavior or sexual cues in rats (Caquineau et al., 2006; Nishitani et al., 2004; Pattij et al., 2005; Witt and Insel, 1994). Consistently, electrical stimulation of the dorsal penile nerve or glans penis produces excitation in OT cells in the PVH and SON of rats (Honda et al., 1999; Yanagimoto et al., 1996). Finally, sexual interaction with a female coincides with an increase in OT release in the PVH (measured by *in vivo* microdialysis) and in the cerebrospinal fluid (measured using cisterna magna sampling) in rats (Hughes et al., 1987; Waldherr and Neumann, 2007).

Taken together, a circuit emerges in which sensory information associated with sexual interactions reaches predominantly magnocellular neurons in the PVN and SON, which release OT locally as well as in the bloodstream. Local OT release activates nearby parvocellular OT neurons (Kita et al., 2006), which in turn facilitate genital reflexes via the sacral parasympathetic nucleus in the spinal cord. Of course this circuit is strongly influenced by other neuropeptides and neurotransmitters, and includes other brain areas besides the PVH, SON and spinal cord, which will be discussed (briefly) later in the review.

3.5. OT effects on erection and ejaculation

Infusions of OT (~1–90 ng) into the cerebral ventricles is generally known to facilitate copulatory reflexes by increasing non-contact erections (observed in sexually potent male rats in the presence of an inaccessible receptive female) and reducing ejaculation latencies and postejaculatory intervals (Argiolas et al., 1985; Arletti et al., 1985; Mizusawa et al., 2002). In squirrel monkeys, ICV OT (0.1 pg) increased the frequency of sexual interactions in dominant, but not subordinate squirrel monkeys (Winslow and Insel, 1991). However, ICV infusion of OT at doses higher than approximately 100–200 ng may not be effective or even inhibiting sexual reflexes, potentially by mimicking sexual satiety (Argiolas et al., 1989b; Mahalati et al., 1991; Stoneham et al., 1985). ICV administration of an OT receptor antagonist (2–1000 ng) shows inhibition of sexual behavior by increasing the intromission latency, decreasing copulatory activity and abolishing ejaculation in rats as well as inhibiting noncontact erections (Argiolas et al., 1989a; Argiolas et al., 1988; Arletti et al., 1992; Melis et al., 1999). It is of note here that the OT receptor antagonist used in these

studies showed an additional strong blockade of the vasopressin (AVP) V_{1A} receptor (Manning et al., 2012). Although the nonapeptide AVP has not been consistently implicated in male sexual behavior, at least one report described the significant role of peripheral V_{1A} receptors in the pro-ejaculatory effects of OT (Gupta et al., 2008).

Consistent with the inhibitory effects of OT receptor antagonists, reducing OT neurotransmission via selective lesions of the parvocellular neurons in the PVN, which reduce the number of OT-immunoreactive fibers in the lumbosacral spinal cord, causes a longer latency to noncontact erections, mounts and intromissions, and decreases seminal emission as well as the postejaculatory interval (Ackerman et al., 1997; Hughes et al., 1987; Liu et al., 1997). Lesions of both the magnocellular and parvocellular neurons in the PVH, which target both OT and AVP neurons (as well as other neurotransmitters and neuropeptides present in the PVH), inhibit noncontact erections and increase the mount frequency and latency to ejaculation during copulation (Liu et al., 1997).

The clear effects on male sexual behavior by changes in general OT neurotransmission have led to a search for projection areas in which OT exerts its effects. Intrathecal injection of OT at the lumbosacral, but not thoracolumbar, level of the spinal cord increased erection (Giuliano et al., 2001). In addition, local infusions of OT in the PVH, hippocampus, ventral tegmental area, and posteromedial amygdala all facilitate noncontact erections (Chen and Chang, 2001; Melis et al., 1986; Melis et al., 2007; Melis et al., 2009). Microinjection of OT into the medial preoptic area (MPOA) reduced the mount frequency and postejaculatory interval in sexually experienced rats, and reduced the ejaculation latency in naïve rats (Gil et al., 2011; 2013). Although local infusion of OT receptor antagonists could inhibit or block the effects of OT in these areas (Chen and Chang, 2001; Melis et al., 1986; Melis et al., 2007; Melis et al., 2009), and administration of an OT receptor antagonist to the spinal cord at the lumbosacral level reduced bulbospongiosus muscle contractions and ejaculation (Clement et al., 2008), the antagonists had no or very mild effects on sexual reflexes when administered in the MPOA or PVH alone (Gil et al., 2011; Melis et al., 1999).

As described above, the effects of OT on male sexual behavior are at least partly executed through central OT neurotransmission. However, a significant role of peripheral OT neurotransmission cannot be excluded. The considerable amount of OT that gets released into the bloodstream following sexual cues or interactions in rats, rabbits and humans (Carmichael et al., 1987; Hillegaart et al., 1998; Kruger et al., 2003; Murphy et al., 1987; Stoneham et al., 1985; Uckert et al., 2003) presumably has a function. Indeed, systemic (intravenous or intraperitoneal) injection of oxytocin facilitates ejaculation, as measured in rats, rabbits and bulls (Arletti et al., 1985; Arletti et al., 1990; 1992; Melin and Kihlstroem, 1963; Palmer et al., 2004; Stoneham et al., 1985). Interestingly, peripheral OT predominantly facilitates ejaculation whereas central OT predominantly affects erection. In fact, peripherally administered OT has been occasionally found to inhibit erection (Palmer et al., 2004; Zhang et al., 2005). Consistently, peripheral administration of OT receptor antagonists inhibited markers of ejaculation and facilitated markers of erection in anaesthetized rats (Clement et al., 2008; Zhang et al., 2005).

The pro-ejaculatory effects of systemic OT may be exerted via peripheral OT receptors that have been found in the testis, epididymis, ductus deferens, prostate and penis of rats, rams and humans, where they mediate the contractility of smooth muscle cells involved in ejaculation (Corona et al., 2012). However, systemic administration of OT also activated the ejaculatory motor pattern, which is exerted by motor neurons in the spinal cord (Carro-Juarez and Rodríguez-Manzo, 2005), suggesting that at least some of the effects of peripheral OT occur via central OT

receptors. Consistently, the novel non-peptidergic OT receptor antagonist, GSK557296, which readily crosses the blood–brain barrier, showed inhibition of various markers of ejaculation, with the strongest effects obtained following ICV and intrathecal infusion at the lumbar level compared to IV and thoracic level infusions (Clement et al., 2013).

Although it is clear that OT strongly affects male sexual behavior, it is much less understood whether variability in the endogenous OT system may underlie innate or experience-induced variability in sexual parameters. When male rats were divided into stable rapid or sluggish ejaculators, it was shown that OT neurons in the SON (but not PVH) were more activated following copulation and ejaculation in the first group compared to the latter (Pattij et al., 2005). In addition, the pro-ejaculatory effects of systemic OT were stronger in sexually sluggish rats compared to sexually potent rats (although OT could not rescue sexual behavior in fully impotent rats) (Arletti et al., 1992). Furthermore, sexually potent rats had a higher expression of OT mRNA in the magnocellular PVN compared to impotent rats (Arletti et al., 1997). These studies suggest that endogenous release of OT in response to sexual cues is naturally inhibited in sluggish rats, or naturally facilitated in sexually efficient rats. Interestingly, sexually efficient male rats had lower levels of OT receptor binding in the MPOA compared to inefficient animals (Gil et al., 2011). The authors suggested that OT receptors might have been internalized or transcriptionally down-regulated in efficient copulators as a response to higher OT levels.

Copulation on the day of sacrifice stimulated OT receptor-mRNA expression in the MPOA, with the highest levels observed in first-time copulators. Sexually experienced males had higher levels of OT receptor protein in the MPOA than sexually naive males and first-time copulators (Gil et al., 2013). These studies indicate that sexual experience, either immediately or over time, changes the endogenous OT system.

3.6. OT and premature ejaculation in men

The ejaculation and erection facilitating effects of oxytocin, as found in animal studies, may contribute to explain a thus far rather unrecognized clinical phenomenon in men with lifelong premature ejaculation (PE): the acute hypertonic or hypererotic state, which occurs when a man with lifelong PE is engaged in an erotic situation. It is characterized by an early ejaculation (ejaculation praecox), but also by an early or facilitated erection (erection praecox) and an immediately occurring detumescence of the penis after ejaculation (detumescencia praecox) (Waldinger, 2014). It has been postulated that the facilitated erection, facilitated ejaculation and facilitated penile detumescence are associated with centrally and peripherally increased oxytocin release in association with involvement of the central serotonergic system and other neurotransmitter and endocrinologic systems (Waldinger, 2014). Research into this triad of symptoms will presumably contribute to a better neurobiological understanding of lifelong PE and its delineation to the three other PE subtypes (Waldinger, 2013; Waldinger, 2008).

In a large sample of men ($n=1517$), no effects were detected of twelve OT receptor gene SNPs on ejaculatory function. However, a heterozygote effect on one SNP in the oxytocin receptor gene (rs75775) was found, so that individuals heterozygous for this SNP had a significantly elevated risk for premature ejaculation symptoms compared with carriers of either homozygote (Jern et al., 2012).

3.7. Other neurotransmitters involved in the OT- male sexual behavior circuit

Recent reviews have extensively discussed the role of various neurotransmitters and neuropeptides (aside from OT) in the

control of male sexual behavior and genital reflexes (Argiolas and Melis, 2013; Corona et al., 2012; Melis and Argiolas, 2011). For example, it is known that the pro-erectile effects of OT are supported by dopamine, glutamate and nitric oxide within the PVH as well as in the hippocampus, amygdala and ventral tegmental area. On the other hand, gamma-aminobutyric acid (GABA), opioid peptides and endocannabinoids inhibit the pro-sexual activity of OT neurons.

Since these reviews did not include a description of the potential role of serotonin (5-HT)–OT interactions (de Jong et al., 2007), we discuss that in more detail here. The role of 5-HT in male sexual behavior is extensive and complex, but in general, increased extracellular 5-HT levels inhibit sexual behavior and increase the ejaculatory threshold through the activation of 5-HT_{1B/2C} receptors (de Jong et al., 2006). Increased 5-HT levels appear to simultaneously counteract their own inhibition of sexual behavior via activation of 5-HT_{1A}-receptors, which facilitate ejaculation (Ahlenius et al., 1981; Snoeren et al., 2013b). Thus, high 5-HT levels (either caused by tonic endogenous release or artificially increased via SSRI administration) maintain a high ejaculatory threshold, and increased or decreased activation or sensitivity of 5-HT_{1A} receptors can adjust it up or downward (Ahlenius and Larsson, 1999; de Jong et al., 2005a; de Jong et al., 2005c; Giuliano and Clement, 2006; Looney et al., 2005). The current theory is that both 5-HT_{1A} autoreceptors (located on 5-HT neurons in the raphe nuclei) and 5-HT_{1A} heteroreceptors (located throughout the CNS) are involved in the facilitation of ejaculation (Snoeren et al., 2013b). Interestingly, 5-HT and OT interact with one another in various brain areas associated with male sexual behavior. In the PVH, 5-HT_{1A} receptors are expressed by oxytocinergic (OT) neurons in the hypothalamus (Zhang et al., 2004). Since 5-HT_{1A} receptor activation induces Fos-immunoreactivity in hypothalamic OT neurons (Coolen et al., 1997; de Jong et al., 2005b) and causes a strong peripheral (and potentially central) release of OT (Bjorkstrand et al., 1996; Van de Kar et al., 1998; Vicentic et al., 1998), this has been proposed as a possible pathway through which 5-HT_{1A} receptor agonists facilitate ejaculation (de Jong et al., 2007). Indeed, unpublished studies have indicated that the pro-ejaculatory effect of systemic injection with 8-OH-DPAT could be partly blocked by ICV administration of an OT receptor antagonist (unpublished observations dr. T.R. de Jong). Furthermore, in the nucleus paragigantocellularis in the brainstem, 5-HT neurons that project to the spinal cord tonically inhibit sexual reflexes and receive putatively inhibitory OT-ergic projections from the PVH (Bancila et al., 2002; Normandin and Murphy, 2011). The sacral parasympathetic nucleus in the spinal cord, which controls the smooth muscle cells in the genitals, receives projections from both inhibitory 5-HT-ergic and facilitatory OT-ergic neurons (Tang et al., 1998). Finally, in the medial preoptic area, OT release facilitates and 5-HT release inhibits male sexual behavior (Dominguez and Hull, 2010; Gil et al., 2011). Taken together, OT and 5-HT appear to counteract each other in various central nervous system regions involved in male sexual behavior, whereas OT appears to support the facilitation of ejaculation by 5-HT_{1A} receptor activation.

4. The fruits of labor: role of OT in parental care

The crucial role of OT in various physiological and behavioral aspects of the peri-partum period is well known. Most knowledge has been derived from female mammals, in particular rats and mice, but more recently other species have been investigated as well. There is an increasing interest in mammalian species in which the male plays a substantial role in the care for offspring. The role of OT in paternal behavior is therefore highlighted in this review.

4.1. Role of OT in maternal care

The most drastic effects of OT appear around parturition when OT supports the delivery of the fetus by stimulating the uterine contractions, and the myoepithelial cells of the mammary gland to induce milk ejection for the succeeding lactation period (Armstrong and Hatton, 2006; Crowley et al., 1992; Lincoln and Paisley, 1982; Summerlee, 1981; Summerlee and Lincoln, 1981). Afferent signals from the uterine cervix or from the nipples stimulate the magnocellular neurons of the PVH and the SON to start the secretion of OT via the posterior pituitary. However, the magnocellular neurons send their products not only into the peripheral circulation. Recently it was observed that the fibers, leaving the magnocellular PVH and SON, send collaterals to the nucleus accumbens region involved in reward mechanisms (Ross et al., 2009). In addition to the magnocellular neurons, the PVH contains numerous parvocellular OT neurons with projections widely dispersed throughout the CNS, expanding even into the lumbosacral parts of the spinal cord (Armstrong and Hatton, 1980; Armstrong et al., 1980; Buijs, 1978a; Buijs et al., 1983; Sawchenko and Swanson, 1982a; Swanson and Kuypers, 1980; Swanson and McKellar, 1979b). Since OT released in the peripheral circulation has virtually no access to the CNS (Mens et al., 1983; Robinson and Jones, 1982), the numerous behavioral changes occurring at the time around parturition must be the result of activation of the parvocellular OT-innervation, of course in cooperation with other hormonal -, transmitter- and peptidergic changes occurring in that period (Herbert, 1993; Rosenblatt et al., 1988). These changes can be labeled as ‘maternal behavior’ (Numan, 2007; Numan et al., 2005; Numan and Sheehan, 1997; Olazabal et al., 2013; Tsuneoka et al., 2013) and are more extensive and complex than just nursing the just born animals because they involve changes in the olfactory system, food intake, sexual behavior, (maternal) aggression and in cognitive and memory functions (Blevins et al., 2003; Blevins et al., 2004; Carter, 1992; 1998; Neumann, 2002; Uvnas-Moberg, 1998; Vaccari et al., 1998). The high CNS-levels of OT around parturition play a strong role in the ‘behavioral transformation’ occurring in the mother, in combination with other factors like estrogen levels and other neuromodulatory substances (McCarthy et al., 1992; Neumann, 2003; Neumann et al., 2003).

OT is present in the olfactory bulbs of the rat because fiber projections from the PVH innervate the bulbs as well as other parts of the olfactory system (Gimpl and Fahrenholz, 2001; Vaccari et al., 1998). There is evidence that some of the OT does not reach the bulbs directly via these neural projections (Yu et al., 1996a; Yu et al., 1996b), and transport of OT via the cerebrospinal fluid is possibly one of the mechanisms involved (Veening et al., 2010; Veening and Olivier, 2013). OT can directly affect neuronal processing in the bulb itself and in addition many amygdaloid and other limbic brain areas contain OT-receptors (Ghosh et al., 2011; Gimpl and Fahrenholz, 2001; Kang et al., 2009; 2011; Miyamichi et al., 2011; Nagayama et al., 2010; Sosulski et al., 2011) and may be influenced by a local release of OT. Similar mechanisms have been studied extensively in sheep (Kendrick, 2000; Kendrick et al., 1997; Kendrick et al., 1986; Kendrick et al., 1991). Functionally, the changes in ‘olfactory sensitivity’ and the succeeding changes in the ‘olfactory memory systems’ appear to serve two important functions: (1) to memorize who mated the female, important for ‘pair-bonding’, see Section 2; (2) to recognize her offspring which deserves her maternal care. Variations in maternal care may affect their offspring directly, including the activity of specific hypothalamic brain areas (Cameron et al., 2011; Champagne, 2008).

The behavioral changes occurring at parturition are complex because for properly feeding the young, adaptations in food intake maybe necessary. In addition specific protective postures are adopted during nursing and milk ejection. At the same time, however, occasional intruders interested in either reproduction

or the young animals themselves have to be kept at a distance to protect the young and a strong reaction in the form of ‘maternal aggression’ may be temporarily necessary. This list of behavioral changes occurring at parturition can be extended but suffices to show the complexity of the changes, involving not only interaction with the pups but also adaptation to additional nutritional requirements and even clear-cut ‘antisocial’ reactions to unwanted companionship. OT seems to play a role in each of these changes.

For nursing the young, OT plays an important role in the milk-ejection reflex, induced by the suction of the nipples (Armstrong, 2007). The number of suckling rat pups determines the amount of OT released in the peripheral blood and the amplitude of the milk ejection burst, but not the frequency of the milk ejections, about once per 5 min (Armstrong, 2007; Hatton and Wang, 2008; Lincoln and Wakerley, 1975a, b; Rossoni et al., 2008). Apparently, a central pattern generator is involved in the effects of continuous suckling stimuli, generating a massive synchronization of OT-neurons across the SON and PVH (Armstrong, 2007; Moos and Richard, 1989). The neural mechanisms behind the synchronization of OT-release turned out to be complex, involving coordinated somato-dendritic release of OT in the PVH and SON (Ludwig and Leng, 2006; Moos et al., 1989), effects of local glial cells (Brown et al., 2013; Wang and Hamilton, 2009; Wang and Hatton, 2009) as well as subcellular mechanisms (Brown et al., 2013; Zhang et al., 2010). Other hypothalamic areas participate in the synchronization mechanisms, like the mammillary region (Wang et al., 2013) and the dorsomedial hypothalamic nucleus (Honda and Higuchi, 2010a, b). The effects of the suckling-induced massive OT-release were also observed in the spinal cord (Ramos et al., 2008). This limited overview may suffice to indicate the extreme importance of OT for nursing the young.

Providing the young with nutrition for an extensive period of time implies that the mother has to take care of her own nutritional status, and OT appears to play a role in that respect as well, be it indirectly.

Lactation, as well as pregnancy itself, induces temporary hyperphagia in mother rats and other mammals, to increase caloric intake with adaptive changes in energy expenditure (Douglas et al., 2007; Fleming, 1976; Leng et al., 2008; Svennersten-Sjaunja and Olsson, 2005; Woodside et al., 2012). Interestingly, both central and peripheral administration of OT suppress food intake (Arletti et al., 1989; Blevins and Ho, 2013; Deblon et al., 2011; Douglas et al., 2007; Leng et al., 2008) with differentiated effects on carbohydrate and fat intake (Olszewski et al., 2010; Sabatier et al., 2013). In the human, OT may induce anorexia (Chaves et al., 2013; Demitrack et al., 1989; Maguire et al., 2013; Sabatier et al., 2013) and is of clinical interest for the treatment of obesity (Blevins and Ho, 2013; Maguire et al., 2013; Sabatier et al., 2013). Suppression of these negative OT-effects on food intake is apparently necessary during the lactation period. The mechanisms for this suppression are complex, as they involve a range of neuronal interactions, for instance related to the ventromedial hypothalamic nucleus and the ‘dorsal vagal complex’ in the caudal brainstem (Douglas et al., 2007; Eriksson et al., 1994; Mullis et al., 2013; Romano et al., 2013; Sabatier et al., 2013; Uvnas-Moberg, 1994). Actually, a large variety of hormonal and peptidergic factors is involved in the temporary suppression of the anorectic effects of OT (Atasoy et al., 2012; Brogan et al., 2000; Ghosh and Sladek, 1995; Linden, 1989; Menzies et al., 2012; Parker and Bloom, 2012; Perello and Raingo, 2013; Woodside et al., 2012; Zagoory-Sharon et al., 2008). The complexity of these control mechanisms makes it impossible to discuss them in more detail, but the evidence provided may suffice to make it clear that for the necessary maternal hyperphagia the regular effects of OT on food intake have to be suppressed temporarily.

In addition to the changes in food intake, the lactation period is characterized by the temporary suppression of ovulation and

reproductive activities. The underlying mechanisms have been investigated extensively in humans, cows, ewes and mares (Bishop, 2013; Gee et al., 2012; Liu et al., 1983; Lunn, 1992; Mann and Lammung, 2001; McNeilly, 2001; Tallam et al., 2000) and appear to involve factors like GnRH and prolactin. There is no reason to suppose that the mechanisms in rats are fundamentally different (Fukushima et al., 2006; Grattan, 2001; Russell et al., 2001).

‘Maternal aggression’ is another, and for the present purpose final example of temporary behavioral changes occurring in the dam at the time of parturition and lactation (Erskine et al., 1978; Gandelman and Simon, 1980; Takahashi and Lore, 1982). It took some time before the involvement of OT was consistently reported (Consiglio and Lucion, 1996; Ferris et al., 1992; Giovenardi et al., 1997; McCarthy et al., 1986; Rosenblatt et al., 1988) because most of the early investigations were aiming at sensory or other hormonal factors (Albert et al., 1992a, b, c; Fleming and Rosenblatt, 1974a, b; Kolunie and Stern, 1990; Mayer and Rosenblatt, 1987; Stern and Kolunie, 1993). OT is currently known as a ‘prosocial’ factor (Lukas et al., 2011), which means that again some of its regular effects have to be suppressed temporarily. It appeared that OT and lactation also have a clear effect on fear behavior (Bosch et al., 2005; Ferreira et al., 2002; Hansen and Ferreira, 1986; Hard and Hansen, 1985; Lonstein et al., 1998; Neumann et al., 2001). Later investigations showed that vasopressin and OT are actually involved in complex and extensive interactions and control together many aspects of ‘social’ behavior, including fear, anxiety and aggression (Beiderbeck et al., 2007; Bosch et al., 2005; Bosch and Neumann, 2012; Ferreira et al., 2002; Neumann, 2008; Neumann et al., 2001; Veenema and Neumann, 2008) and these effects have to be studied in each brain area involved separately (Veenema and Neumann, 2008). No surprise then that the ‘hypothalamo-pituitary-adrenal- (HPA-) axis’ is also involved in these interactions (Caldwell, 1992; Giovenardi et al., 2005; Klampfl et al., 2013; Neumann, 2001; Neumann et al., 2001; Vilela and Giusti-Paiva, 2011).

The ‘hypothalamic-attack-area’ (HAA) (Kruk, 1991; Kruk et al., 1984a; Roeling et al., 1994; Siegel et al., 1999), located in the intermediate hypothalamus and along the ventrolateral part of the ventromedial hypothalamic nucleus (VMHvl) (Geeraedts et al., 1990a, b; Nieuwenhuys et al., 1982; Veening et al., 1982) is apparently involved in maternal aggression (Motta et al., 2013; Hansen, 1989). Pregnancy and lactation had no effect on the effective thresholds of electrical brain stimulation in the HAA (Mos et al., 1987). However, the HAA (Roeling et al., 1994) and the PVH have a widely distributed network of neuronal connections and therefore the neuronal adaptations occurring in the period of maternal aggression may occur outside the HAA. Lesioning the PVH is effective (Consiglio and Lucion, 1996; Giovenardi et al., 1997; Olazabal and Ferreira, 1997) but many additional brain areas have been indicated in the literature, too many for the present discussion. We intend to make an exception, however, for a single brain area: the VMHvl. This particular part of the VMH plays an important role in lordosis behavior, as discussed in Section 3.3, (Chung et al., 1990; Flanagan-Cato, 2011; Flanagan-Cato et al., 2001; Pfaff and Sakuma, 1979d; Romano et al., 1988; Sakuma and Pfaff, 1981; Sakuma and Pfaff, 1982; Schwanzel-Fukuda et al., 1984). At the same time, however, this region of the hypothalamus forms part of the HAA, mentioned above. It has been observed that under certain circumstances mother rats show ambivalent behavior towards a male intruder (Agrati et al., 2011). Recently, this brain area has been investigated in the mouse (Anderson, 2012; Lin et al., 2011; Saper, 2011), with quite remarkable results for which the reader is referred to Section 5 for further details.

4.2. Role of OT in paternal care

Although the majority of mammalian species display uniparental care, with mothers performing the great majority of parental

tasks such as feeding, sheltering, and grooming pups, in a small percentage (~5%) of mammalian species the fathers (and/or related males) demonstrate a considerable amount of caring behavior. In addition to humans, some alloparental primate species and biparental rodent species have been used as experimental animal models to investigate paternal behavior. The most widely used rodent species include the obligatory paternal prairie voles (*Microtus ochrogaster*), mandarin voles (*Lasiopodomys mandarinus*), and California mice (*Peromyscus californicus*) as well as the facultatively paternal meadow voles (*Microtus pennsylvanicus*) and various outbred strains of laboratory house mice (*Mus musculus*). Although there have been various findings suggesting that OT influences paternal care, and/or that paternal care is associated with changes in the OT system, the evidence is somewhat anecdotal and indicates species specificity. In addition, since most biparental species form monogamous pair bonds as well, see Section 2, it can be difficult to distinguish between the effects of becoming a father and the unavoidable simultaneous socio-sexual experience of cohabitation with a female, which affects the OT system as well (see Section 2).

In various species, it has been shown that OT levels or OT release is correlated with paternal behavior or paternal state. In human fathers, plasma and salivary OT levels are positively correlated with social engagement, affect synchrony, and positive communicative sequences with their 4–6-month-old infant (Feldman et al., 2011; Gordon et al., 2010). In paternally experienced males of a biparental primate, the common marmoset (*Callithrix jacchus*), the release of OT from cultured explants of the hypothalamus is increased compared with naïve, paternally inexperienced males (Woller et al., 2012). In juvenile and adult prairie voles, pup exposure causes transient elevations in plasma OT levels and increased Fos-immunoreactivity in OT neurons (Kenkel et al., 2012). However, aside from the acute pup-induced activation of the OT system, neither OT gene expression in the hypothalamus nor OT receptor binding in a number of brain regions differs between sexually naïve and paternally experienced male prairie voles (Wang et al., 2000). In male California mice, males cohabitating with a pregnant female show increased plasma OT levels compared with virgin males, but OT levels decline prior to parturition, remain low throughout the postpartum period, and do not correlate with the level of paternal behavior (Gubernick et al., 1995). Mandarin vole fathers show increased OT-immunoreactivity in the PVH and SON compared with virgin males, but brief exposure to pups or cohabitation with a female without pups have the same effects (Song et al., 2010). In meadow voles, sexually and paternally experienced, paternally behaving males have significantly higher OT receptor binding in several brain regions (accessory olfactory nucleus, lateral septum, BNST, and lateral amygdala) compared to sexually and paternally inexperienced, non-paternally behaving males; however, it is possible that these differences are caused by sexual rather than paternal experience (Gil et al., 2013; Parker and Lee, 2001). In addition, male California mice with various levels of sexual and paternal experience do not show increases in Fos-immunoreactivity in response to auditory, olfactory and visual cues from an inaccessible newborn pup (de Jong et al., 2009). In one non-mammalian species, the monogamous and biparental Central American convict cichlid (*Amatitlania nigrofasciata*), males displaying most paternal care show (Gil et al., 2013) increased Fos-immunoreactivity induction in parvocellular preoptic isotocin (IST) neurons, IST being the homolog of OT in fish (O'Connell et al., 2012).

Taken together, pup exposure and paternal experience appear to coincide with changes in the OT system that differ in strength among species. Conversely, artificially increasing or decreasing OT neurotransmission affects paternal care in various species. Intranasal OT administration improves social-paternal interactions

between human fathers and their infant- or toddler-aged children (Naber et al., 2010; Weisman et al., 2012). In male common marmosets, ICV administration of OT increases food transfer from fathers to offspring (Saito and Nakamura, 2011). Conversely, in the Central American convict cichlid, systemic injection of an IST receptor antagonist blocks paternal care (O'Connell et al., 2012) and ICV administration of a combination of OT and AVP receptor antagonists (but not either antagonist alone) inhibits paternal behavior in paternally naïve prairie voles (Bales et al., 2004). In the same experiment, ICV administration of OT did not further increase the high levels of paternal care and impaired kyphosis (nursing posture).

In both humans and mice, genetic associations between the OT system and paternal behavior have been demonstrated. In human fathers (and mothers) genotyped with risk alleles for the OT receptor and for CD38, a membrane glycoprotein that facilitates OT secretion, plasma OT is reduced and the fathers show less parental touch. On the other hand, low-risk CD38 alleles and high plasma OT predict longer durations of parent-infant gaze synchrony (Feldman et al., 2012). Consistently, in facultatively paternal ICR-strain mice, paternal care is reduced following knockout of the CD38 gene, and ICV OT can rescue this impaired behavior. ICV OT further improves paternal care following reinstatement of the CD38 gene in the nucleus accumbens of CD38^{−/−} sires (Akther et al., 2013).

Taken together, it appears that increased activity in the OT system is associated with increased or improved paternal care, and decreased activity in the OT system with the opposite, which is similar to the association between OT and maternal care. However, much more than maternal care, the influence of the OT system in paternal care is highly variable and species-specific and appears to depend on a) the natural level of paternal care displayed, and b) the basal neurobiology of the OT system, in males of a given species.

5. The VMHvl: a puzzling brain region ...!

In the experiments of Cameron et al. (Cameron et al., 2011) it was observed that the less receptive high-LG females showed the greatest neural activation in the VMHvl after mating. Since less than 50% of the activated neurons show ER α -immunoreactivity (Calizo and Flanagan-Cato, 2003; Tetel et al., 1994a; Tetel et al., 1993; 1994b), it is possible that some of the mechanisms working in the area lateral to the VMHvl do not promote lordosis, but prepare the female for the termination of the estrous period. As studied in detail by Georgescu and Pfaus (Georgescu and Pfaus, 2006a, b; Georgescu et al., 2009), VCS induced a considerable number of activated glutamatergic neurons in the VMHvl, and these numbers decreased under influence of estradiol and/or progesterone. The authors concluded that “the effect of EB and P to blunt the activation of the inhibitory glutamate neurons in the VMH by VCS may therefore be a mechanism by which females can experience a requisite amount of VCS before estrus termination is engaged” (Georgescu et al., 2009) and “it may well be the case that glutamate release in the VMH increases active rejection responses, consistent with our previous findings for the effect of glutamate agonist and antagonist infusions to the VMH” (Georgescu and Pfaus, 2006a, b).

‘Estrous termination’ can be described as a temporary form of ‘sexual satiety’ starting when a certain amount of VCS has been obtained in a given period of time. It lies at the basis of the rewarding aspects of ‘pacing behavior’ (Pfaus et al., 2012; Pfaus et al., 2000). The female starts to reject the anogenital investigations of the male, normally occurring in the early precopulatory phase, as well as all copulatory activities. When the female gets

pregnant and starts feeding her young, in the lactation period, the female not only rejects the male but her behavior becomes openly aggressive, 'maternal aggression', as discussed before. Apparently, the 'neural substrate' for this aggressive behavior is present in the female brain as well and can be temporarily released from its regular constraints. From our preceding discussion, it is almost inevitable to conclude that the plastic neuronal changes occurring in the VMHvl, in the lateral extensions of long dendrites into the intermediate hypothalamic zone and the OT-fibers contacting them, are involved in the behavioral changes from lordosis, via rejection to aggression.

The male-female similarities in the neural substrate are also evident if we pay attention to the induction of lordosis in male rodents. Defeminization- and masculinization processes follow their own specific patterns, concerning timing and hormonal control mechanisms (Gladue and Clemens, 1982; Kalcheim et al., 1981; Ward, 1972). Prenatal stress induces lordosis in adult male rats (Ward, 1972). The defeminization-process is under control of a variety of factors like androgens (Diaz et al., 1995; Segovia et al., 2009; Vega Matuszczyk et al., 1988; Ward, 1977; Weisz and Ward, 1980; Whalen et al., 1986), estrogen and progesterone (Dohanich and Ward, 1980; Olster and Blaustein, 1991; Ward et al., 1977) and various brain areas like anterior hypothalamic and preoptic regions, lateral septum and PAG (Gordon et al., 1979; Kondo et al., 1993; Kondo et al., 1990; Tsukahara et al., 2003; Tsukahara and Yamanouchi, 2001; 2002; Whitney and Herrenkohl, 1977). No surprise, then, to find the VMHvl involved also in these processes (Romano et al., 1990; Segovia et al., 2009). We can safely conclude that the neural substrate involved in lordosis behavior is not only present in the female, but in the male rat brain as well, where it can be uncovered by experiments leading to only a partial defeminization.

But if 'lordosis behavior' is represented in the male brain, what about 'aggressive behavior' in the male and female brains? The question rises if they are also represented in identical brain areas.

In the male rat brain, aggressive behavior can be induced by low currents of intracranial brain stimulation and the location has been precisely defined (Kruk, 1991; Kruk et al., 1984a; Kruk and van der Poel, 1980; Kruk et al., 1979; Kruk et al., 1983; Kruk et al., 1998; Lammers et al., 1988; Roeling et al., 1994), in the intermediate hypothalamic region, rostral to and alongside the VMHvl (Geeraedts et al., 1990a, b; Kruk et al., 1983; Roeling et al., 1994; Veening, 1992). Of course additional brain areas are involved as well (septal regions, amygdala) but many of their fibers converge on this brain region, alongside the VMHvl. As discussed earlier, lactating mother rats show the same behavior, 'maternal aggression', arising from exactly the same brain regions (Kruk et al., 1984b; Mos et al., 1987). Apparently, the neural substrate for 'attack behavior' is organized also in the male and female rat brain in a very similar way, with hormonal changes being necessary to uncover this behavior in the female rat, see Section 4.1.

At first sight it is rather surprising that we have to conclude that two very different behaviors like lordosis and attack behavior, including their completely different postures, lordotic vs kyphotic respectively, may be induced by the same region of the hypothalamus. Very recently, however, new findings in the mouse brain were reported, shedding a completely new light on this puzzling situation. The findings will be discussed in the next section.

5.1. The VMHvl as multi-modal hub for the control of sexual and aggressive behavior

In female rats, stimulation of the VMH produces lordosis (Calizo and Flanagan-Cato, 2002; Flanagan-Cato, 2011; Mathews and Edwards, 1977b; Pfaff and Sakuma, 1979a). In contrast, damaging the VMH or the descending projections disrupts sexual

behavior (Hennessey et al., 1990; Mathews and Edwards, 1977a; Pfaff and Sakuma, 1979b, d; Pfeifle et al., 1980; Rajendren et al., 1991). Neural activation of the female ventrolateral VMH (VMHvl) can be observed, but not exclusively, during and after sexual behavior (Blaustein et al., 1994; Coolen et al., 1996; Flanagan et al., 1993; Pfaus et al., 1993; Rowe and Erskine, 1993; Tetel et al., 1993; Wersinger et al., 1993). Interestingly, damaging the lateral side of the VMH affects female aggression (Hansen, 1989). In accordance, GABA (and glutamate) in or near the VMHvl is involved in maternal aggression (Hansen and Ferreira, 1986; Lee and Gammie, 2009). Recently, it was shown that during maternal aggression, c-Fos-staining can be especially observed in the VMHvl (Motta et al., 2013).

The hormones c.q. neurotransmitters estrogen (Pfaff, 1980; Pfaff et al., 2008), progesterone (DeBold and Malsbury, 1989; Pleim et al., 1991) and oxytocin (Bale and Dorsa, 1995; Griffin et al., 2010) have a very stimulating effect on lordosis, when applied in the VMH, see Section 3.3. Interestingly, estrogens affect both the structure and function of the VMH (Flanagan-Cato, 2011; McCarthy and Konkle, 2005; Pfaff, 1980; 2011). Importantly, the estrogen-alpha receptors (ER-alpha) and oxytocin receptors are present in the VMHvl, but not in the VMHdm (Flanagan-Cato, 2011). Remarkably, under natural conditions, both the levels of estradiol and progesterone fluctuate across the estrous cycle, and during these fluctuations dendritic connections are reorganized and numbers of dendritic spines are increased (Carrer and Aoki, 1982; Frankfurt et al., 1990; Frankfurt and McEwen, 1991). In addition, these gonadal hormones turn on genes in the hypothalamus, thereby making sexual behavior possible (Ogawa et al., 1997; Pfaff, 2011).

For a long time it was puzzling how the same brain area, the VMH, could be involved in both (maternal) aggression and sexual behavior (Hansen, 1989; Hansen and Ferreira, 1986; Hrabovszky et al., 2005; Kruk et al., 1979; Kruk et al., 1983; Lammers et al., 1988; Luiten et al., 1985; Olivier and Wiepkema, 1974; Roeling et al., 1994; Veening et al., 2005). Recently, experiments in male mice from the research group of David Anderson at Caltech, have shed new light on the problem (Lin et al., 2011).

In male mice, optogenetic stimulation of the dorsomedial hypothalamus (VMHdm) causes flight and freezing behavior, while VMHvl causes aggressive attacks of opponent males, females and even inanimate objects (plastic glove). Pharmacogenetic silencing of VMHvl neurons reversibly inhibits this aggression (Lin et al., 2011). In agreement, it has been shown by pharmacogenetic neural inhibition in freely behaving male mice, that the VMHdl is involved in fear of a predator, while the VMHvl is involved in aggression and social fear, but not in fear of foot shock (Silva et al., 2013). Again, suggesting an important role for the VMHvl in social encounters.

The research group of Anderson also showed that electrophysiological recording of firing rate of VMHvl neurons resulted in 40% of the neurons being more activated by a male intruder, about 12% exclusively during attack, and about 25% by a female mating partner. About 50% of the recorded neurons responded initially to both a male and a female intruder, suggesting that the neurons share some common input, to differentiate between the male and female later during the encounters (Lin et al., 2011). Most intriguingly, VMHvl neurons are activated during attack and are inhibited during interactions with a female, suggesting that the VMHvl neurons mediating aggression are actively suppressed during mating. Special attention has to be paid to the fact that many VMHvl neurons are activated during both male-male and male-female encounters (Anderson, 2012). It was hypothesized that these dual-activated neurons could be involved in the organization of motor programs common to both attack and mating, such as approach behavior to a conspecific or ano-genital investigation as

part of appetitive sexual behavior (Anderson, 2012). In agreement, recently it was shown that predator cat odor induces approach behavior in parasite *Toxoplasma gondii*-infected male rats, whereas healthy rats show avoidance behavior (House et al., 2011). In detail, exposure to cat urine induces avoidance behavior associated with c-Fos-staining, reflecting neural activation, in the accessory olfactory bulb (AOB), posteroventral medial amygdala (MEApv), and VMHdm. Bizarrely, the brain parasite infected rats show approach behavior towards the predator cat odors, associated with c-Fos staining in AOB, posterodorsal medial amygdala (MEApd). A similar staining pattern can be seen, when healthy male rats approach female rat odor (House et al., 2011). Thus, during evolution the parasite has managed to change brain functioning after infection, thereby perturbing the “avoidance brain pathway”, shifting neural activity to the nearby “approach brain pathway” (see above).

Apparently, sex and aggression in rodents not only share olfactory stimuli to ‘chase’ between the agonistic and mating patterns of behavior, but the VMHvl also appears to be a much more intricate point of sensory convergence before the decision ‘either-or’ can be taken. Altogether, it can be summarized that the VMH is a multimodal-hub for the control of social encounters. The projections from the VMHdm to the dorsolateral periaqueductal gray (PAGdl) serve the behavioral avoidance (flight) response with underlying physiology (increase in heart rate and blood pressure), while projections from the VMHvl to the ventrolateral periaqueductal gray (PAGvl) serve the behavioral approach response (Canteras, 2002; Keay and Bandler, 2001). Remarkably, in female rats this projection to the PAGvl may be very important during sexual behavior, when males grip the female tightly with their fore paws to enable intromission, because this pathway increases the nociceptive thresholds (Keay et al., 1994). Projections from the VMH to PAG are also important for the motor control of lordosis and vocalizations, including the ultrasounds used by rodents for courtship behavior (Floody and DeBold, 2004). Both neurons in the VMHvl and PAGvl contain high amounts of ER- α receptors, suggesting that both areas are under the control of gonadal hormones (Flanagan-Cato, 2011; Gerrits et al., 2009; Murphy and Hoffman, 2001).

In the seventies Millhouse (Millhouse, 1973b) showed with golgi-staining that a fiber capsule is entirely surrounding the VMH, which encompasses a dense terminal field. These fibers all have boutons en passant and short collaterals, which terminate in the capsule. Also oxytocin fibers descending from the magnocellular part of the PVH run along the VMHvl (Griffin et al., 2010). Furthermore, oxytocin-containing neurons are activated by sexual behavior (Arletti and Bertolini, 1985; Caldwell et al., 1994; Flanagan et al., 1993), their dendritic contacts are reorganized by ovarian hormones (Ferri and Flanagan-Cato, 2012; Griffin et al., 2010) and the effects of oxytocin are ‘pro-social’ (Macdonald and Macdonald, 2010; Meyer-Lindenberg, 2008; Striepens et al., 2011), including ‘pro-sexual’ for males and females (Flanagan-Cato, 2011) and ‘anti-aggressive’ (Calcagnoli et al., 2013; Caldwell et al., 1994; DeVries et al., 1997; Dhakar et al., 2012; Harmon et al., 2002; Neumann, 2008; Pedersen et al., 1994; Ragnauth et al., 2005). Thus, OT-fibers that are part of the VMHvl fiber capsule are in an excellent position to innervate long dendrites with OT receptors of interneurons in the VMHvl (Flanagan-Cato, 2011), thereby influencing the potential ‘sex-aggression balance’ towards appetitive sexual behavior.

6. Summary and conclusions

The conclusion seems warranted that under normal circumstances OT plays an important facilitating role during both male

and female reproductive behavior, in rodents (Pedersen and Boccia, 2002) and other mammals including man and the ‘unifying principle’ of the OT action in the brain “is to facilitate social encounters by reducing the associated anxiety” (McCarthy, 1995). Much research is needed to further delineate the complex organization of the neural networks involved in reproductive and sexual behaviors in male and female mammals including man. OT seems to play an important role in many of these networks but its precise role needs further experimentation and hypotheses.

References

- Ackerman, A.E., Lange, G.M., Clemens, L.G., 1997. Effects of paraventricular lesions on sex behavior and seminal emission in male rats. *Physiol. Behav.* 63, 49–53.
- Agrati, D., Fernandez-Guasti, A., Ferrero, M., Ferreira, A., 2011. Coexpression of sexual behavior and maternal aggression: the ambivalence of sexually active mother rats toward male intruders. *Behav. Neurosci.* 125, 446–451.
- Ahlenius, S., Larsson, K., 1999. Synergistic actions of the 5-HT_{1A} receptor antagonist WAY-100635 and citalopram on male rat ejaculatory behavior. *Eur. J. Pharmacol.* 379, 1–6.
- Ahlenius, S., Larsson, K., Svensson, L., Hjorth, S., Carlsson, A., Lindberg, P., Wikstrom, H., Sanchez, D., Arvidsson, L.E., Hacksell, U., Nilsson, J.L., 1981. Effects of a new type of 5-HT receptor agonist on male rat sexual behavior. *Pharmacol. Biochem. Behav.* 15, 785–792.
- Akther, S., Korshnova, N., Zhong, J., Liang, M., Cherepanov, S.M., Lopatina, O., Komleva, Y.K., Salmina, A.B., Nishimura, T., Fakhrul, A.A., Hirai, H., Kato, I., Yamamoto, Y., Takasawa, S., Okamoto, H., Higashida, H., 2013. CD38 in the nucleus accumbens and oxytocin are related to paternal behavior in mice. *Mol. Brain* 6, 41.
- Albert, D.J., Jonik, R.H., Walsh, M.L., 1992a. Hormone-dependent aggression in male and female rats: experiential, hormonal, and neural foundations. *Neurosci. Biobehav. Rev.* 16, 177–192.
- Albert, D.J., Jonik, R.H., Walsh, M.L., 1992b. Interaction of estradiol, testosterone, and progesterone in the modulation of hormone-dependent aggression in the female rat. *Physiol. Behav.* 52, 773–779.
- Albert, D.J., Jonik, R.H., Walsh, M.L., 1992c. Ovariectomy does not attenuate aggression by primiparous lactating female rats. *Physiol. Behav.* 52, 1043–1046.
- Alvares, G.A., Hickie, I.B., Guastella, A.J., 2010. Acute effects of intranasal oxytocin on subjective and behavioral responses to social rejection. *Exp. Clin. Psychopharmacol.* 18, 316–321.
- Anderson, D.J., 2012. Optogenetics, sex, and violence in the brain: implications for psychiatry. *Biol. Psychiatry* 71, 1081–1089.
- Arendash, G.W., Gorski, R.A., 1983. Suppression of lordotic responsiveness in the female rat during mesencephalic electrical stimulation. *Pharmacol. Biochem. Behav.* 19, 351–357.
- Argiolas, A., Collu, M., D'Aquila, P., Gessa, G.L., Melis, M.R., Serra, G., 1989a. Apomorphine stimulation of male copulatory behavior is prevented by the oxytocin antagonist d(CH₂)₅ Tyr(Me)-Orn⁸-vasotocin in rats. *Pharmacol. Biochem. Behav.* 33, 81–83.
- Argiolas, A., Collu, M., Gessa, G.L., Melis, M.R., Serra, G., 1988. The oxytocin antagonist d(CH₂)₅ Tyr(Me)-Orn⁸-vasotocin inhibits male copulatory behaviour in rats. *Eur. J. Pharmacol.* 149, 389–392.
- Argiolas, A., Melis, M.R., 2013. Neuropeptides and central control of sexual behaviour from the past to the present: a review. *Prog. Neurobiol.* 108, 80–107.
- Argiolas, A., Melis, M.R., Gessa, G.L., 1985. Intraventricular oxytocin induces yawning and penile erection in rats. *Eur. J. Pharmacol.* 117, 395–396.
- Argiolas, A., Melis, M.R., Stancampiano, R., Gessa, G.L., 1989b. Penile erection and yawning induced by oxytocin and related peptides: structure–activity relationship. *Peptides* 10, 559–563.
- Arletti, R., Bazzani, C., Castelli, M., Bertolini, A., 1985. Oxytocin improves male copulatory performance in rats. *Horm. Behav.* 19, 14–20.
- Arletti, R., Benelli, A., Bertolini, A., 1989. Influence of oxytocin on feeding behavior in the rat. *Peptides* 10, 89–93.
- Arletti, R., Benelli, A., Bertolini, A., 1990. Sexual behavior of aging male rats is stimulated by oxytocin. *Eur. J. Pharmacol.* 179, 377–381.
- Arletti, R., Benelli, A., Bertolini, A., 1992. Oxytocin involvement in male and female sexual behavior. *Ann. N. Y. Acad. Sci.* 652, 180–193.
- Arletti, R., Bertolini, A., 1985. Oxytocin stimulates lordosis behavior in female rats. *Neuropeptides* 6, 247–253.
- Arletti, R., Calza, L., Giardino, L., Benelli, A., Cavazzuti, E., Bertolini, A., 1997. Sexual impotence is associated with a reduced production of oxytocin and with an increased production of opioid peptides in the paraventricular nucleus of male rats. *Neurosci. Lett.* 233, 65–68.
- Armstrong, W.E., 2007. The neurophysiology of neurosecretory cells. *J. Physiol.* 585, 645–647.
- Armstrong, W.E., Hatton, G.I., 1980. The localization of projection neurons in the rat hypothalamic paraventricular nucleus following vascular and neurohypophyseal injections of HRP. *Brain Res. Bull.* 5, 473–477.
- Armstrong, W.E., Hatton, G.I., 2006. The puzzle of pulsatile oxytocin secretion during lactation: some new pieces. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 291, R26–R28.

- Armstrong, W.E., Warach, S., Hatton, G.I., McNeill, T.H., 1980. Subnuclei in the rat hypothalamic paraventricular nucleus: a cytoarchitectural, horseradish peroxidase and immunocytochemical analysis. *Neuroscience* 5, 1931–1958.
- Atasoy, D., Betley, J.N., Su, H.H., Sternson, S.M., 2012. Deconstruction of a neural circuit for hunger. *Nature* 488, 172–177.
- Bale, T.L., Dorsa, D.M., 1995. Sex differences in and effects of estrogen on oxytocin receptor messenger ribonucleic acid expression in the ventromedial hypothalamus. *Endocrinology* 136, 27–32.
- Bales, K.L., Kim, A.J., Lewis-Reese, A.D., Sue Carter, C., 2004. Both oxytocin and vasopressin may influence alloparental behavior in male prairie voles. *Horm. Behav.* 45, 354–361.
- Bancila, M., Giuliano, F., Rampin, O., Mailly, P., Brisorgueil, M.J., Calas, A., Verge, D., 2002. Evidence for a direct projection from the paraventricular nucleus of the hypothalamus to putative serotonergic neurons of the nucleus paragigantocellularis involved in the control of erection in rats. *Eur. J. Neurosci.* 16, 1240–1248.
- Bancroft, J., 2005. The endocrinology of sexual arousal. *J. Endocrinol.* 186, 411–427.
- Bargmann, W., 1949. Über die Neurosekretorische Verknüpfung von Hypothalamus und Neurohypophyse. *Z. Zellforsch. mikrosk. Anat.* 34, 610–634.
- Baskerville, T.A., Douglas, A.J., 2010. Dopamine and oxytocin interactions underlying behaviors: potential contributions to behavioral disorders. *CNS Neurosci. Ther.* 16, e92–123.
- Baumgartner, T., Heinrichs, M., Vonlanthen, A., Fischbacher, U., Fehr, E., 2008. Oxytocin shapes the neural circuitry of trust and trust adaptation in humans. *Neuron* 58, 639–650.
- Bealer, S.L., Armstrong, W.E., Crowley, W.R., 2010. Oxytocin release in magnocellular nuclei: neurochemical mediators and functional significance during gestation. *Am. J. Physiol. Regul., Integr. Comp. Physiol.* 299, R452–R458.
- Behnia, B., Heinrichs, M., Bergmann, W., Jung, S., Germann, J., Schedlowski, M., Hartmann, U., Kruger, T.H., 2014. Differential effects of intranasal oxytocin on sexual experiences and partner interactions in couples. *Horm. Behav.* 65, 308–318.
- Beiderbeck, D.I., Neumann, I.D., Veenema, A.H., 2007. Differences in intermale aggression are accompanied by opposite vasopressin release patterns within the septum in rats bred for low and high anxiety. *Eur. J. Neurosci.* 26, 3597–3605.
- Beinfeld, M.C., 2001. An introduction to neuronal cholecystokinin. *Peptides* 22, 1197–1200.
- Bennett, A.L., Greco, B., Blasberg, M.E., Blaustein, J.D., 2002. Response to male odours in progesterone receptor- and oestrogen receptor-containing cells in female rat brain. *J. Neuroendocrinol.* 14, 442–449.
- Bishop, C.V., 2013. Progesterone inhibition of oxytocin signaling in endometrium. *Front. Neurosci.* 7, 138.
- Bjorkstrand, E., Ahlenius, S., Smedh, U., Uvnäs-Moberg, K., 1996. The oxytocin receptor antagonist 1-deamino-2-D-Tyr-(OEt)-4-Thr-8-Orn-oxytocin inhibits effects of the 5-HT_{1A} receptor agonist 8-OH-DPAT on plasma levels of insulin, cholecystokinin and somatostatin. *Regul. Pept.* 63, 47–52.
- Blaicher, W., Gruber, D., Bieglmayer, C., Blaicher, A.M., Knogler, W., Huber, J.C., 1999. The role of oxytocin in relation to female sexual arousal. *Gynecol. Obstet. Invest.* 47, 125–126.
- Blaustein, J.D., Tetel, M.J., Ricciardi, K.H., Delville, Y., Turcotte, J.C., 1994. Hypothalamic ovarian steroid hormone-sensitive neurons involved in female sexual behavior. *Psychoneuroendocrinology* 19, 505–516.
- Blevins, J.E., Eakin, T.J., Murphy, J.A., Schwartz, M.W., Baskin, D.G., 2003. Oxytocin innervation of caudal brainstem nuclei activated by cholecystokinin. *Brain Res.* 993, 30–41.
- Blevins, J.E., Ho, J.M., 2013. Role of oxytocin signaling in the regulation of body weight. *Rev. Endocr. Metab. Disord.* 14, 311–329.
- Blevins, J.E., Schwartz, M.W., Baskin, D.G., 2004. Evidence that paraventricular nucleus oxytocin neurons link hypothalamic leptin action to caudal brain stem nuclei controlling meal size. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 287, R87–R96.
- Boccia, M.L., Goursaud, A.P., Bachevalier, J., Anderson, K.D., Pedersen, C.A., 2007. Peripherally administered non-peptide oxytocin antagonist, L368,899, accumulates in limbic brain areas: a new pharmacological tool for the study of social motivation in non-human primates. *Horm. Behav.* 52, 344–351.
- Booth, C., Wayman, C.P., Jackson, V.M., 2010. An ex vivo multi-electrode approach to evaluate endogenous hormones and receptor subtype pharmacology on evoked and spontaneous neuronal activity within the ventromedial hypothalamus; translation from female receptivity. *J. Sex. Med.* 7, 2411–2423.
- Borrow, A.P., Cameron, N.M., 2012. The role of oxytocin in mating and pregnancy. *Horm. Behav.* 61, 266–276.
- Bosch, O.J., Meddle, S.L., Beiderbeck, D.I., Douglas, A.J., Neumann, I.D., 2005. Brain oxytocin correlates with maternal aggression: link to anxiety. *J. Neurosci.: Off. J. Soc. Neurosci.* 25, 6807–6815.
- Bosch, O.J., Neumann, I.D., 2012. Both oxytocin and vasopressin are mediators of maternal care and aggression in rodents: from central release to sites of action. *Horm. Behav.* 61, 293–303.
- Breslow, E., Burman, S., 1990. Molecular, thermodynamic, and biological aspects of recognition and function in neurophysin-hormone systems: a model system for the analysis of protein-peptide interactions. *Adv. Enzymol. Relat. Areas Mol. Biol.* 63, 1–67.
- Brogan, R.S., Grove, K.L., Smith, M.S., 2000. Differential regulation of leptin receptor but not orexin in the hypothalamus of the lactating rat. *J. Neuroendocrinol.* 12, 1077–1086.
- Brown, C.H., Bains, J.S., Ludwig, M., Stern, J.E., 2013. Physiological regulation of magnocellular neurosecretory cell activity: integration of intrinsic, local and afferent mechanisms. *J. Neuroendocrinol.* 25, 678–710.
- Bueno, J., Pfaff, D.W., 1976. Single unit recording in hypothalamus and preoptic area of estrogen-treated and untreated ovariectomized female rats. *Brain Res.* 101, 67–78.
- Buijs, R.M., 1978a. Intra- and extrahypothalamic vasopressin and oxytocin pathways in the rat. Pathways to the limbic system, medulla oblongata and spinal cord. *Cell Tissue Res.* 192, 423–435.
- Buijs, R.M., 1978b. Intra- and extrahypothalamic vasopressin and oxytocin pathways in the rat. Pathways to the limbic system, medulla oblongata and spinal cord. *Cell Tissue Res.* 192, 423–435.
- Buijs, R.M., De Vries, G.J., Van Leeuwen, F.W., Swaab, D.F., 1983. Vasopressin and oxytocin: distribution and putative functions in the brain. *Prog. Brain Res.* 60, 115–122.
- Buijs, R.M., Swaab, D.F., Dogterom, J., van Leeuwen, F.W., 1978. Intra- and extrahypothalamic vasopressin and oxytocin pathways in the rat. *Cell Tissue Res.* 186, 423–433.
- Buma, P., 1989. Characterization of luteinizing hormone-releasing hormone fibres in the mesencephalic central grey substance of the rat. *Neuroendocrinology* 49, 623–630.
- Calcagnoli, F., de Boer, S.F., Althaus, M., den Boer, J.A., Koolhaas, J.M., 2013. Antiaggressive activity of central oxytocin in male rats. *Psychopharmacology (Berl)* 229, 639–651.
- Caldwell, J.D., 1992. Central oxytocin and female sexual behavior. *Ann. N. Y. Acad. Sci.* 652, 166–179.
- Caldwell, J.D., 2002. A sexual arousability model involving steroid effects at the plasma membrane. *Neurosci. Biobehav. Rev.* 26, 13–30.
- Caldwell, J.D., Jirikowski, G.F., Greer, E.R., Pedersen, C.A., 1989. Medial preoptic area oxytocin and female sexual receptivity. *Behav. Neurosci.* 103, 655–662.
- Caldwell, J.D., Jirikowski, G.F., Greer, E.R., Stumpf, W.E., Pedersen, C.A., 1988. Ovarian steroids and sexual interaction alter oxytocinergic content and distribution in the basal forebrain. *Brain Res.* 446, 236–244.
- Caldwell, J.D., Johns, J.M., Fagg, B.M., Senger, M.A., Pedersen, C.A., 1994. Infusion of an oxytocin antagonist into the medial preoptic area prior to progesterone inhibits sexual receptivity and increases rejection in female rats. *Horm. Behav.* 28, 288–302.
- Caldwell, J.D., Prange Jr., A.J., Pedersen, C.A., 1986. Oxytocin facilitates the sexual receptivity of estrogen-treated female rats. *Neuropeptides* 7, 175–189.
- Calizo, L.H., Flanagan-Cato, L.M., 2002. Estrogen-induced dendritic spine elimination on female rat ventromedial hypothalamic neurons that project to the periaqueductal gray. *J. Comp. Neurol.* 447, 234–248.
- Calizo, L.H., Flanagan-Cato, L.M., 2003. Hormonal-neural integration in the female rat ventromedial hypothalamus: triple labeling for estrogen receptor- α , retrograde tract tracing from the periaqueductal gray, and mating-induced Fos expression. *Endocrinology* 144, 5430–5440.
- Cameron, N., Erskine, M.S., 2003. c-FOS expression in the forebrain after mating in the female rat is altered by adrenalectomy. *Neuroendocrinology* 77, 305–313.
- Cameron, N.M., Soehngen, E., Meaney, M.J., 2011. Variation in maternal care influences ventromedial hypothalamus activation in the rat. *J. Neuroendocrinol.* 23, 393–400.
- Canteras, N.S., 2002. The medial hypothalamic defensive system: hodological organization and functional implications. *Pharmacol., Biochem. Behav.* 71, 481–491.
- Canteras, N.S., Simerly, R.B., Swanson, L.W., 1994. Organization of projections from the ventromedial nucleus of the hypothalamus: a Phaseolus vulgaris-leucoagglutinin study in the rat. *J. Comp. Neurol.* 348, 41–79.
- Caqueneau, C., Leng, G., Guan, X.M., Jiang, M., Van der Ploeg, L., Douglas, A.J., 2006. Effects of alpha-melanocyte-stimulating hormone on magnocellular oxytocin neurones and their activation at intromission in male rats. *J. Neuroendocrinol.* 18, 685–691.
- Carmichael, M.S., Humbert, R., Dixen, J., Palmisano, G., Greenleaf, W., Davidson, J.M., 1987. Plasma oxytocin increases in the human sexual response. *J. Clin. Endocrinol. Metab.* 64, 27–31.
- Carmichael, M.S., Warburton, V.L., Dixen, J., Davidson, J.M., 1994. Relationships among cardiovascular, muscular, and oxytocin responses during human sexual activity. *Arch. Sexual Behav.* 23, 59–79.
- Carrer, H.F., Aoki, A., 1982. Ultrastructural changes in the hypothalamic ventromedial nucleus of ovariectomized rats after estrogen treatment. *Brain Res.* 240, 221–233.
- Carro-Juarez, M., Rodriguez-Manzo, G., 2005. Evidence for the presence and functioning of the spinal generator for ejaculation in the neonatal male rat. *Int. J. Impot. Res.* 17, 270–276.
- Carter, C.S., 1992. Oxytocin and sexual behavior. *Neurosci. Biobehav. Rev.* 16, 131–144.
- Carter, C.S., 1998. Neuroendocrine perspectives on social attachment and love. *Psychoneuroendocrinology* 23, 779–818.
- Carter, C.S., 2014. Oxytocin pathways and the evolution of human behavior. *Annu. Rev. Psychol.* 65, 17–39.
- Carter, C.S., DeVries, A.C., Getz, L.L., 1995. Physiological substrates of mammalian monogamy: the prairie vole model. *Neurosci. Biobehav. Rev.* 19, 303–314.
- Carter, C.S., Porges, S.W., 2013. The biochemistry of love: an oxytocin hypothesis. *EMBO Rep.* 14, 12–16.
- Carter, S.J., Richardson, C.M., Wakerley, J.B., 2003. Excitatory effects of oxytocin and cholecystokinin on oxytocin neurones: differences between virgin, pregnant and lactating rats. *Neurosci. Lett.* 351, 13–16.

- Champagne, F.A., 2008. Epigenetic mechanisms and the transgenerational effects of maternal care. *Front. Neuroendocrinol.* 29, 386–397.
- Chaves, V.E., Tilelli, C.Q., Brito, N.A., Brito, M.N., 2013. Role of oxytocin in energy metabolism. *Peptides* 45, 9–14.
- Chen, K., Chang, L.S., 2001. Oxytocinergic neurotransmission at the hippocampus in the central neural regulation of penile erection in the rat. *Urology* 58, 107–112.
- Cho, M.M., DeVries, A.C., Williams, J.R., Carter, C.S., 1999. The effects of oxytocin and vasopressin on partner preferences in male and female prairie voles (*Microtus ochrogaster*). *Behav. Neurosci.* 113, 1071–1079.
- Chung, S.K., Pfaff, D.W., Cohen, R.S., 1990. Projections of ventromedial hypothalamic neurons to the midbrain central gray: an ultrastructural study. *Neuroscience* 38, 395–407.
- Churchland, P.S., Winkelman, P., 2012. Modulating social behavior with oxytocin: how does it work? What does it mean? *Horm. Behav.* 61, 392–399.
- Clement, P., Bernabe, J., Compagnie, S., Alexandre, L., McCallum, S., Giuliano, F., 2013. Inhibition of ejaculation by the non-peptide oxytocin receptor antagonist GSK557296: a multi-level site of action. *Br. J. Pharmacol.* 169, 1477–1485.
- Clement, P., Peeters, M., Bernabe, J., Denys, P., Alexandre, L., Giuliano, F., 2008. Brain oxytocin receptors mediate ejaculation elicited by 7-hydroxy-2-(di-N-propylamino) tetralin (7-OH-DPAT) in anesthetized rats. *Br. J. Pharmacol.* 154, 1150–1159.
- Coirini, H., Johnson, A.E., McEwen, B.S., 1989. Estradiol modulation of oxytocin binding in the ventromedial hypothalamic nucleus of male and female rats. *Neuroendocrinology* 50, 193–198.
- Consiglio, A.R., Lucion, A.B., 1996. Lesion of hypothalamic paraventricular nucleus and maternal aggressive behavior in female rats. *Physiol. Behav.* 59, 591–596.
- Coolen, L.M., Olivier, B., Peters, H.J., Veening, J.G., 1997. Demonstration of ejaculation-induced neural activity in the male rat brain using 5-HT_{1A} agonist 8-OH-DPAT. *Physiol. Behav.* 62, 881–891.
- Coolen, L.M., Peters, H.J., Veening, J.G., 1996. Fos immunoreactivity in the rat brain following consummatory elements of sexual behavior: a sex comparison. *Brain Res.* 738, 67–82.
- Coria-Avila, G.A., Pfau, J.G., 2007. Neuronal activation by stimuli that predict sexual reward in female rats. *Neuroscience* 148, 623–632.
- Corona, G., Jannini, E.A., Vignozzi, L., Rastrelli, G., Maggi, M., 2012. The hormonal control of ejaculation. *Nat. Rev. Urol.* 9, 508–519.
- Crockford, C., Wittig, R.M., Langergraber, K., Ziegler, T.E., Zuberbühler, K., Deschner, T., 2013. Urinary oxytocin and social bonding in related and unrelated wild chimpanzees. *Proc. Biol. Sci./R. Soc.* 280, 20122765.
- Crowley, W.R., Parker, S.L., Armstrong, W.E., Spinolo, L.H., Grosvenor, C.E., 1992. Neurotransmitter and neurohormonal regulation of oxytocin secretion in lactation. *Ann. N. Y. Acad. Sci.* 652, 286–302.
- Daniels, D., Flanagan-Cato, L.M., 2000. Functionally-defined compartments of the lordosis neural circuit in the ventromedial hypothalamus in female rats. *J. Neurobiol.* 45, 1–13.
- Daniels, D., Miselis, R.R., Flanagan-Cato, L.M., 1999. Central neuronal circuit innervating the lordosis-producing muscles defined by transneuronal transport of pseudorabies virus. *J. Neurosci.: Off. J. Soc. Neurosci.* 19, 2823–2833.
- Davis, P.G., McEwen, B.S., Pfaff, D.W., 1979. Localized behavioral effects of tritiated estradiol implants in the ventromedial hypothalamus of female rats. *Endocrinology* 104, 898–903.
- de Jong, T.R., Chauke, M., Harris, B.N., Saltzman, W., 2009. From here to paternity: neural correlates of the onset of paternal behavior in California mice (*Peromyscus californicus*). *Horm. Behav.* 56, 220–231.
- de Jong, T.R., Pattij, T., Veening, J.G., Dederen, P.J., Waldinger, M.D., Cools, A.R., Olivier, B., 2005a. Citalopram combined with WAY 100635 inhibits ejaculation and ejaculation-related Fos immunoreactivity. *Eur. J. Pharmacol.* 509, 49–59.
- de Jong, T.R., Pattij, T., Veening, J.G., Dederen, P.J., Waldinger, M.D., Cools, A.R., Olivier, B., 2005b. Effects of chronic paroxetine pretreatment on (+/–)-8-hydroxy-2-(di-n-propyl-amino)tetralin induced c-fos expression following sexual behavior. *Neuroscience* 134, 1351–1361.
- de Jong, T.R., Pattij, T., Veening, J.G., Waldinger, M.D., Cools, A.R., Olivier, B., 2005c. Effects of chronic selective serotonin reuptake inhibitors on 8-OH-DPAT-induced facilitation of ejaculation in rats: comparison of fluvoxamine and paroxetine. *Psychopharmacology* 179, 509–515.
- de Jong, T.R., Snaphaan, L.J., Pattij, T., Veening, J.G., Waldinger, M.D., Cools, A.R., Olivier, B., 2006. Effects of chronic treatment with fluvoxamine and paroxetine during adolescence on serotonin-related behavior in adult male rats. *Eur. Neuropsychopharmacol.: J. Eur. Coll. Neuropsychopharmacol.* 16, 39–48.
- de Jong, T.R., Veening, J.G., Olivier, B., Waldinger, M.D., 2007. Oxytocin involvement in SSRI-induced delayed ejaculation: a review of animal studies. *J. Sex. Med.* 4, 14–28.
- Deblon, N., Veyrat-Durebex, C., Bourgoin, L., Caillon, A., Bussier, A.L., Petrosino, S., Piscitelli, F., Legros, J.J., Geenen, V., Foti, M., Wahli, W., Di Marzo, V., Rohner-Jeanrenaud, F., 2011. Mechanisms of the anti-obesity effects of oxytocin in diet-induced obese rats. *PLoS One* 6, e25565.
- DeBold, J.F., Malsbury, C.W., 1989. Facilitation of sexual receptivity by hypothalamic and midbrain implants of progesterone in female hamsters. *Physiol. Behav.* 46, 655–660.
- Dellovade, T.L., Chan, J., Vennstrom, B., Forrest, D., Pfaff, D.W., 2000. The two thyroid hormone receptor genes have opposite effects on estrogen-stimulated sex behaviors. *Nat. Neurosci.* 3, 472–475.
- Demitrack, M.A., Lesem, M.D., Brandt, H.A., Pigott, T.A., Jimerson, D.C., Altemus, M., Gold, P.W., 1989. Neurohypophyseal dysfunction: implications for the pathophysiology of eating disorders. *Psychopharmacol. Bull.* 25, 439–443.
- Devost, D., Wrzal, P., Zingg, H.H., 2008. Oxytocin receptor signalling. *Prog. Brain Res.* 170, 167–176.
- DeVries, A.C., Young 3rd, W.S., Nelson, R.J., 1997. Reduced aggressive behaviour in mice with targeted disruption of the oxytocin gene. *J. Neuroendocrinol.* 9, 363–368.
- Dhakar, M.B., Rich, M.E., Reno, E.L., Lee, H.J., Caldwell, H.K., 2012. Heightened aggressive behavior in mice with lifelong versus postweaning knockout of the oxytocin receptor. *Horm. Behav.* 62, 86–92.
- Diaz, D.R., Fleming, D.E., Rhees, R.W., 1995. The hormone-sensitive early postnatal periods for sexual differentiation of feminine behavior and luteinizing hormone secretion in male and female rats. *Brain research. Dev. Brain Res.* 86, 227–232.
- Dohanich, G.P., Ward, I.L., 1980. Sexual behavior in male rats following intracerebral estrogen application. *J. Comp. Physiol. Psychol.* 94, 634–640.
- Dominguez, J.M., Hull, E.M., 2010. Serotonin impairs copulation and attenuates ejaculation-induced glutamate activity in the medial preoptic area. *Behav. Neurosci.* 124, 554–557.
- Donner, N., Neumann, I.D., 2009. Effects of chronic intracerebral prolactin on the oxytocinergic and vasopressinergic system of virgin ovariectomized rats. *Neuroendocrinology* 90, 315–322.
- Douglas, A.J., Johnstone, L.E., Leng, G., 2007. Neuroendocrine mechanisms of change in food intake during pregnancy: a potential role for brain oxytocin. *Physiol. Behav.* 91, 352–365.
- Du Vigneaud, V., Ressler, C., Trippett, S., 1953. The sequence of amino acids in oxytocin, with a proposal for the structure of oxytocin. *J. Biol. Chem.* 205, 949–957.
- Dugger, B.N., Morris, J.A., Jordan, C.L., Breedlove, S.M., 2007. Androgen receptors are required for full masculinization of the ventromedial hypothalamus (VMH) in rats. *Horm. Behav.* 51, 195–201.
- Egli, M., Leeners, B., Kruger, T.H., 2010. Prolactin secretion patterns: basic mechanisms and clinical implications for reproduction. *Reproduction* 140, 643–654.
- Emery, D.E., 1986. Effects of endocrine state on sociosexual behavior of female rats tested in a complex environment. *Behav. Neurosci.* 100, 71–78.
- Eriksson, M., Bjorkstrand, E., Smedh, U., Alster, P., Matthiesen, A.S., Uvnäs-Moberg, K., 1994. Role of vagal nerve activity during suckling. Effects on plasma levels of oxytocin, prolactin, VIP, somatostatin, insulin, glucagon, glucose and of milk secretion in lactating rats. *Acta Physiol. Scand.* 151, 453–459.
- Erskine, M.S., Denenberg, V.H., Goldman, B.D., 1978. Aggression in the lactating rat: effects of intruder age and test arena. *Behav. Biol.* 23, 52–66.
- Fahrbach, S.E., Morrell, J.L., Pfaff, D.W., 1989. Studies of ventromedial hypothalamic afferents in the rat using three methods of HRP application. *Exp. Brain Res. Exp. Hirnforsch.* 77, 221–233.
- Feldman, R., Gordon, I., Zagoory-Sharon, O., 2011. Maternal and paternal plasma, salivary, and urinary oxytocin and parent-infant synchrony: considering stress and affiliation components of human bonding. *Dev. Sci.* 14, 752–761.
- Feldman, R., Zagoory-Sharon, O., Weisman, O., Schneidman, I., Gordon, I., Maoz, R., Shalev, I., Ebstein, R.P., 2012. Sensitive parenting is associated with plasma oxytocin and polymorphisms in the OXTR and CD38 genes. *Biol. Psychiatry* 72, 175–181.
- Ferreira, A., Pereira, M., Agrati, D., Uriarte, N., Fernandez-Guasti, A., 2002. Role of maternal behavior on aggression, fear and anxiety. *Physiol. Behav.* 77, 197–204.
- Ferri, S.L., Flanagan-Cato, L.M., 2012. Oxytocin and dendrite remodeling in the hypothalamus. *Horm. Behav.* 61, 251–258.
- Ferris, C.F., Foote, K.B., Meltzer, H.M., Plenby, M.G., Smith, K.L., Insel, T.R., 1992. Oxytocin in the amygdala facilitates maternal aggression. *Ann. N. Y. Acad. Sci.* 652, 456–457.
- Flanagan, L.M., Pfau, J.G., Pfaff, D.W., McEwen, B.S., 1993. Induction of Fos immunoreactivity in oxytocin neurons after sexual activity in female rats. *Neuroendocrinology* 58, 352–358.
- Flanagan-Cato, L.M., 2011. Sex differences in the neural circuit that mediates female sexual receptivity. *Front. Neuroendocrinol.* 32, 124–136.
- Flanagan-Cato, L.M., Calizo, L.H., Daniels, D., 2001. The synaptic organization of VMH neurons that mediate the effects of estrogen on sexual behavior. *Horm. Behav.* 40, 178–182.
- Flanagan-Cato, L.M., Lee, B.J., Calizo, L.H., 2006. Co-localization of midbrain projections, progesterone receptors, and mating-induced fos in the hypothalamic ventromedial nucleus of the female rat. *Horm. Behav.* 50, 52–60.
- Flanagan-Cato, L.M., McEwen, B.S., 1995. Pattern of Fos and Jun expression in the female rat forebrain after sexual behavior. *Brain Res.* 673, 53–60.
- Fleming, A.S., 1976. Control of food intake in the lactating rat: role of suckling and hormones. *Physiol. Behav.* 17, 841–848.
- Fleming, A.S., Rosenblatt, J.S., 1974a. Olfactory regulation of maternal behavior in rats. I. Effects of olfactory bulb removal in experienced and inexperienced lactating and cycling females. *J. Comp. Physiol. Psychol.* 86, 221–232.
- Fleming, A.S., Rosenblatt, J.S., 1974b. Olfactory regulation of maternal behavior in rats. II. Effects of peripherally induced anosmia and lesions of the lateral olfactory tract in pup-induced virgins. *J. Comp. Physiol. Psychol.* 86, 233–246.
- Floody, O.R., DeBold, J.F., 2004. Effects of midbrain lesions on lordosis and ultrasound production. *Physiol. Behav.* 82, 791–804.
- Frankfurt, M., Gould, E., Woolley, C.S., McEwen, B.S., 1990. Gonadal steroids modify dendritic spine density in ventromedial hypothalamic neurons: a Golgi study in the adult rat. *Neuroendocrinology* 51, 530–535.
- Frankfurt, M., McEwen, B.S., 1991. 5,7-Dihydroxytryptamine and gonadal steroid manipulation alter spine density in ventromedial hypothalamic neurons. *Neuroendocrinology* 54, 653–657.

- Freeman, M.E., Banks, J.A., 1980. Hypothalamic sites which control the surges of prolactin secretion induced by cervical stimulation. *Endocrinology* 106, 668–673.
- Frohman, K.S., Pitchers, K.K., Balfour, M.E., Coolen, L.M., 2010. Mixing pleasures: review of the effects of drugs on sex behavior in humans and animal models. *Horm. Behav.* 58, 149–162.
- Fukushima, A., Yin, P., Ishida, M., Sugiyama, N., Arita, J., 2006. Pup removal suppresses estrogen-induced surges of LH secretion and activation of GnRH neurons in lactating rats. *J. Endocrinol.* 191, 339–348.
- Gandelman, R., Simon, N.G., 1980. Postpartum fighting in the rat: nipple development and the presence of young. *Behav. Neural Biol.* 28, 350–360.
- Gee, E.K., Gillespie, L., Bolwell, C.F., 2012. Effect of oxytocin on suppression of oestrus in mares exhibiting normal oestrous cycles. *N. Z. Vet. J.* 60, 189–193.
- Geeraedts, L.M., Nieuwenhuys, R., Veening, J.G., 1990a. Medial forebrain bundle of the rat: III. Cytoarchitecture of the rostral (telencephalic) part of the medial forebrain bundle bed nucleus. *J. Comp. Neurol.* 294, 507–536.
- Geeraedts, L.M., Nieuwenhuys, R., Veening, J.G., 1990b. Medial forebrain bundle of the rat: IV. Cytoarchitecture of the caudal (lateral hypothalamic) part of the medial forebrain bundle bed nucleus. *J. Comp. Neurol.* 294, 537–568.
- Gelez, H., Poirier, S., Facchinetti, P., Allers, K.A., Wayman, C., Alexandre, L., Giuliano, F., 2010. Neuroanatomical evidence for a role of central melanocortin-4 receptors and oxytocin in the efferent control of the rodent clitoris and vagina. *J. Sex. Med.* 7, 2056–2067.
- Georgescu, M., Pfau, J.G., 2006a. Role of glutamate receptors in the ventromedial hypothalamus in the regulation of female rat sexual behaviors I. Behavioral effects of glutamate and its selective receptor agonists AMPA, NMDA and kainate. *Pharmacol., Biochem. Behav.* 83, 322–332.
- Georgescu, M., Pfau, J.G., 2006b. Role of glutamate receptors in the ventromedial hypothalamus in the regulation of female rat sexual behaviors. II. Behavioral effects of selective glutamate receptor antagonists AP-5, CNQX, and DNQX. *Pharmacol., Biochem. Behav.* 83, 333–341.
- Georgescu, M., Sabongui, C., Del Corpo, A., Marsan, L., Pfau, J.G., 2009. Vagino-cervical stimulation induces Fos in glutamate neurons in the ventromedial hypothalamus: attenuation by estrogen and progesterone. *Horm. Behav.* 56, 450–456.
- Gerendai, I., Toth, I.E., Kocsis, K., Boldogkoi, Z., Rusvai, M., Halasz, B., 2001. Identification of CNS neurons involved in the innervation of the epididymis: a viral transneuronal tracing study. *Auton. Neurosci.: Basic Clin.* 92, 1–10.
- Gerrits, P.O., Krukerink, M., Veening, J.G., 2009. Columnar organization of estrogen receptor- α immunoreactive neurons in the periaqueductal gray projecting to the nucleus para-retroambiguus in the caudal brainstem of the female golden hamster. *Neuroscience* 161, 459–474.
- Ghosh, R., Sladek, C.D., 1995. Role of prolactin and gonadal steroids in regulation of oxytocin mRNA during lactation. *Am. J. Physiol.* 269, E76–E84.
- Ghosh, S., Larson, S.D., Hefzi, H., Marnoy, Z., Cutforth, T., Dokka, K., Baldwin, K.K., 2011. Sensory maps in the olfactory cortex defined by long-range viral tracing of single neurons. *Nature* 472, 217–220.
- Gil, M., Bhatt, R., Picotte, K.B., Hull, E.M., 2011. Oxytocin in the medial preoptic area facilitates male sexual behavior in the rat. *Horm. Behav.* 59, 435–443.
- Gil, M., Bhatt, R., Picotte, K.B., Hull, E.M., 2013. Sexual experience increases oxytocin receptor gene expression and protein in the medial preoptic area of the male rat. *Psychoneuroendocrinology* 38, 1688–1697.
- Gimpl, G., Fahrenholz, F., 2001. The oxytocin receptor system: structure, function, and regulation. *Physiol. Rev.* 81, 629–683.
- Giovenardi, M., de Azevedo, M.S., da Silva, S.P., Hermel Edo, E., Gomes, C.M., Lucion, A.B., 2005. Neonatal handling increases fear and aggression in lactating rats. *Physiol. Behav.* 86, 209–217.
- Giovenardi, M., Padoin, M.J., Cadore, L.P., Lucion, A.B., 1997. Hypothalamic paraventricular nucleus, oxytocin, and maternal aggression in rats. *Ann. N. Y. Acad. Sci.* 807, 606–609.
- Giraldi, A., Marson, L., Nappi, R., Pfau, J., Traish, A.M., Vardi, Y., Goldstein, I., 2004. Physiology of female sexual function: animal models. *J. Sex. Med.* 1, 237–253.
- Giraldi, A., Rellini, A.H., Pfau, J., Laan, E., 2013. Female sexual arousal disorders. *J. Sex. Med.* 10, 58–73.
- Giuliano, F., Bernabe, J., McKenna, K., Longueville, F., Rampin, O., 2001. Spinal proerectile effect of oxytocin in anesthetized rats. *Am. J. Physiol. Regul., Integr. Comp. Physiol.* 280, R1870–R1877.
- Giuliano, F., Clement, P., 2006. Serotonin and premature ejaculation: from physiology to patient management. *Eur. Urol.* 50, 454–466.
- Gladue, B.A., Clemens, L.G., 1982. Development of feminine sexual behavior in the rat: androgenic and temporal influences. *Physiol. Behav.* 29, 263–267.
- Gordon, I., Zagoory-Sharon, O., Leckman, J.F., Feldman, R., 2010. Oxytocin and the development of parenting in humans. *Biol. Psychiatry* 68, 377–382.
- Gordon, J.H., Nance, D.M., Wallis, C.J., Gorski, R.A., 1979. Effects of septal lesions and chronic estrogen treatment on dopamine, GABA and lordosis behavior in male rats. *Brain Res. Bull.* 4, 85–89.
- Gozaalka, B.B., Lester, G.L., 1987. Oxytocin-induced facilitation of lordosis behaviour in rats is progesterone-dependent. *Neuropeptides* 10, 55–65.
- Grattan, D.R., 2001. The actions of prolactin in the brain during pregnancy and lactation. *Prog. Brain Res.* 133, 153–171.
- Gray, G.D., Soderstein, P., Tallentire, D., Davidson, J.M., 1978. Effects of lesions in various structures of the suprachiasmatic-preoptic region on LH regulation and sexual behavior in female rats. *Neuroendocrinology* 25, 174–191.
- Grewen, K.M., Girdler, S.S., Amico, J., Light, K.C., 2005. Effects of partner support on resting oxytocin, cortisol, norepinephrine, and blood pressure before and after warm partner contact 67, 531–538. *Psychosom. Med.* 67, 531–538.
- Griffin, G.D., Ferri-Kolwicz, S.L., Reyes, B.A., Van Bockstaele, E.J., Flanagan-Cato, L.M., 2010. Ovarian hormone-induced reorganization of oxytocin-labeled dendrites and synapses lateral to the hypothalamic ventromedial nucleus in female rats. *J. Comp. Neurol.* 518, 4531–4545.
- Gubernick, D.J., Winslow, J.T., Jensen, P., Jeanotte, L., Bowen, J., 1995. Oxytocin changes in males over the reproductive cycle in the monogamous, biparental California mouse, *Peromyscus californicus*. *Horm. Behav.* 29, 59–73.
- Gupta, J., Russell, R., Wayman, C., Hurley, D., Jackson, V., 2008. Oxytocin-induced contractions within rat and rabbit ejaculatory tissues are mediated by vasopressin V1A receptors and not oxytocin receptors. *Br. J. Pharmacol.* 155, 118–126.
- Hansen, S., 1989. Medial hypothalamic involvement in maternal aggression of rats. *Behav. Neurosci.* 103, 1035–1046.
- Hansen, S., Ferreira, A., 1986. Food intake, aggression, and fear behavior in the mother rat: control by neural systems concerned with milk ejection and maternal behavior. *Behav. Neurosci.* 100, 64–70.
- Hard, E., Hansen, S., 1985. Reduced fearfulness in the lactating rat. *Physiol. Behav.* 35, 641–643.
- Harlan, R.E., Shivers, B.D., Pfaff, D.W., 1983. Midbrain microinfusions of prolactin increase the estrogen-dependent behavior, lordosis. *Science* 219, 1451–1453.
- Harmon, A.C., Huhman, K.L., Moore, T.O., Albers, H.E., 2002. Oxytocin inhibits aggression in female Syrian hamsters. *J. Neuroendocrinol.* 14, 963–969.
- Hatton, G.L., Wang, Y.F., 2008. Neural mechanisms underlying the milk ejection burst and reflex. *Prog. Brain Res.* 170, 155–166.
- Hayashi, S., Orimo, T., 1975. Sterility and delayed vaginal opening following electrolytic lesions placed neonatally in medial preoptic area in female rats. *Endocrinol. Jpn.* 22, 409–413.
- Hendricks, C.H., Gabel, R.A., 1960. Use of intranasal oxytocin in obstetrics. 1. A laboratory evaluation. *Am. J. Obstet. Gynecol.* 79, 780–788.
- Hennessey, A.C., Camak, L., Gordon, F., Edwards, D.A., 1990. Connections between the pontine central gray and the ventromedial hypothalamus are essential for lordosis in female rats. *Behav. Neurosci.* 104, 477–488.
- Herbert, J., 1993. Peptides in the limbic system: neurochemical codes for co-ordinated adaptive responses to behavioural and physiological demand. *Prog. Neurobiol.* 41, 723–791.
- Hillegaart, V., Alster, P., Uvnas-Moberg, K., Ahlenius, S., 1998. Sexual motivation promotes oxytocin secretion in male rats. *Peptides* 19, 39–45.
- Honda, K., Higuchi, T., 2010a. Effects of unilateral electrolytic lesion of the dorsomedial nucleus of the hypothalamus on milk-ejection reflex in the rat. *J. Reprod. Dev.* 56, 98–102.
- Honda, K., Higuchi, T., 2010b. Electrical activities of neurones in the dorsomedial hypothalamic nucleus projecting to the supraoptic nucleus during milk-ejection reflex in the rat. *J. Reprod. Dev.* 56, 336–340.
- Honda, K., Yanagimoto, M., Negoro, H., Narita, K., Murata, T., Higuchi, T., 1999. Excitation of oxytocin cells in the hypothalamic supraoptic nucleus by electrical stimulation of the dorsal penile nerve and tactile stimulation of the penis in the rat. *Brain Res. Bull.* 48, 309–313.
- House, P.K., Vyas, A., Sapolsky, R., 2011. Predator cat odors activate sexual arousal pathways in brains of *Toxoplasma gondii* infected rats. *PLoS One* 6, e23277.
- Hrabovszky, E., Halasz, J., Meelis, W., Kruk, M.R., Liposits, Z., Haller, J., 2005. Neurochemical characterization of hypothalamic neurons involved in attack behavior: glutamatergic dominance and co-expression of thyrotropin-releasing hormone in a subset of glutamatergic neurons. *Neuroscience* 133, 657–666.
- Hughes, A.M., Everitt, B.J., Lightman, S.L., Todd, K., 1987. Oxytocin in the central nervous system and sexual behaviour in male rats. *Brain Res.* 414, 133–137.
- Hull, E.M., 2011. Sex, drugs and glutony: how the brain controls motivated behaviors. *Physiol. Behav.* 104, 173–177.
- Hull, E.M., Muschamp, J.W., Sato, S., 2004. Dopamine and serotonin: influences on male sexual behavior. *Physiol. Behav.* 83, 291–307.
- Huntingford, P.J., 1961. Intranasal use of synthetic oxytocin in management of breast-feeding. *Br. Med. J.* 1, 709–711.
- Insel, T.R., 1992. Oxytocin—a neuropeptide for affiliation: evidence from behavioral, receptor autoradiographic, and comparative studies. *Psychoneuroendocrinology* 17, 3–35.
- Insel, T.R., 2010. The challenge of translation in social neuroscience: a review of oxytocin, vasopressin, and affiliative behavior. *Neuron* 65, 768–779.
- Insel, T.R., Shapiro, L.E., 1992. Oxytocin receptor distribution reflects social organization in monogamous and polygamous voles. *Proc. Natl. Acad. Sci. USA* 89, 5981–5985.
- Jern, P., Westberg, L., Johansson, A., Jonsson, L., Corander, J., Sandnabba, N.K., Santtila, P., 2012. Are single nucleotide polymorphisms in the oxytocin and vasopressin 1A/1B receptor genes likely candidates for variation in ejaculatory function? *BJU Int.* 110, E1173–E1180.
- Kalchauer, S., Szechtman, H., Koch, Y., 1981. Bisexual behavior in male rats treated neonatally with antibodies to luteinizing hormone-releasing hormone. *J. Comp. Physiol. Psychol.* 95, 36–44.
- Kagerbauer, S.M., Martin, J., Schuster, T., Blobner, M., Kochs, E.F., Landgraf, R., 2013. Plasma oxytocin and vasopressin do not predict neuropeptide concentrations in human cerebrospinal fluid. *J. Neuroendocrinol.* 25, 668–673.
- Kang, N., Baum, M.J., Cherry, J.A., 2009. A direct main olfactory bulb projection to the vomeronasal amygdala in female mice selectively responds to volatile pheromones from males. *Eur. J. Neurosci.* 29, 624–634.
- Kang, N., Baum, M.J., Cherry, J.A., 2011. Different profiles of main and accessory olfactory bulb mitral/tufted cell projections revealed in mice using an anterograde tracer and a whole-mount, flattened cortex preparation. *Chem. Senses* 36, 251–260.

- Kavaliers, M., Clipperton-Allen, A., Cragg, C.L., Gustafsson, J.A., Korach, K.S., Muglia, L., Choleris, E., 2012. Male risk taking, female odors, and the role of estrogen receptors. *Physiol. Behav.* 107, 751–761.
- Keay, K.A., Bandler, R., 2001. Parallel circuits mediating distinct emotional coping reactions to different types of stress. *Neurosci. Biobehav. Rev.* 25, 669–678.
- Keay, K.A., Clement, C.I., Owler, B., Depaulis, A., Bandler, R., 1994. Convergence of deep somatic and visceral nociceptive information onto a discrete ventrolateral midbrain periaqueductal gray region. *Neuroscience* 61, 727–732.
- Kelly, J., Swanson, L.W., 1980. Additional forebrain regions projecting to the posterior pituitary: preoptic region, bed nucleus of the stria terminalis, and zona incerta. *Brain Res.* 197, 1–9.
- Kendrick, K.M., 2000. Oxytocin, motherhood and bonding. *Exp. Physiol.* 85 Spec No, 111S–124S.
- Kendrick, K.M., Da Costa, A.P., Broad, K.D., Ohkura, S., Guevara, R., Levy, F., Keverne, E.B., 1997. Neural control of maternal behaviour and olfactory recognition of offspring. *Brain Res. Bull.* 44, 383–395.
- Kendrick, K.M., Fabre-Nys, C., Blache, D., Goode, J.A., Broad, K.D., 1993. The role of oxytocin release in the mediobasal hypothalamus of the sheep in relation to female sexual receptivity. *J. Neuroendocrinol.* 5, 13–21.
- Kendrick, K.M., Keverne, E.B., Baldwin, B.A., Sharman, D.F., 1986. Cerebrospinal fluid levels of acetylcholinesterase, monoamines and oxytocin during labour, parturition, vaginocervical stimulation, lamb separation and suckling in sheep. *Neuroendocrinology* 44, 149–156.
- Kendrick, K.M., Keverne, E.B., Hinton, M.R., Goode, J.A., 1991. Cerebrospinal fluid and plasma concentrations of oxytocin and vasopressin during parturition and vaginocervical stimulation in the sheep. *Brain Res. Bull.* 26, 803–807.
- Kenkel, W.M., Paredes, J., Yee, J.R., Pournajafi-Nazarloo, H., Bales, K.L., Carter, C.S., 2012. Neuroendocrine and behavioural responses to exposure to an infant in male prairie voles. *J. Neuroendocrinol.* 24, 874–886.
- Kirkpatrick, M.E., Merrill, L., 2011. Effect of systemic blockade of alpha(1)-noradrenergic receptors on sex behavior and vaginal-cervical stimulation-induced Fos in female rats. *Pharmacol., Biochem. Behav.* 97, 486–493.
- Kita, I., Yoshida, Y., Nishino, S., 2006. An activation of parvocellular oxytocinergic neurons in the paraventricular nucleus in oxytocin-induced yawning and penile erection. *Neurosci. Res.* 54, 269–275.
- Klampfl, S.M., Neumann, I.D., Bosch, O.J., 2013. Reduced brain corticotropin-releasing factor receptor activation is required for adequate maternal care and maternal aggression in lactating rats. *Eur. J. Neurosci.* 38, 2742–2750.
- Knobloch, H.S., Charlet, A., Hoffmann, L.C., Eliava, M., Khurlev, S., Cetin, A.H., Osten, P., Schwarz, M.K., Seeburg, P.H., Stoop, R., Grinevich, V., 2012. Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron* 73, 553–566.
- Knobloch, H.S., Grinevich, V., 2014. Evolution of oxytocin pathways in the brain of vertebrates. *Front. Behav. Neurosci.* 8, 31.
- Koch, S.B., van Zuiden, M., Nawijn, L., Frijling, J.L., Veltman, D.J., Olf, M., 2014. Intranasal oxytocin as strategy for medication-enhanced psychotherapy of PTSD: salience processing and fear inhibition processes. *Psychoneuroendocrinology* 40C, 242–256.
- Kolonie, J.M., Stern, J.M., 1990. Maternal aggression: disruption by perioral anesthesia in lactating Long-Evans rats (*Rattus norvegicus*). *J. Comp. Psychol.* 104, 352–360.
- Komisaruk, B.R., Sansone, G., 2003. Neural pathways mediating vaginal function: the vagus nerves and spinal cord oxytocin. *Scand. J. Psychol.* 44, 241–250.
- Komisaruk, B.R., Whipple, B., 1998. Love as sensory stimulation: physiological consequences of its deprivation and expression. *Psychoneuroendocrinology* 23, 927–944.
- Komisaruk, B.R., Whipple, B., Crawford, A., Liu, W.C., Kalnin, A., Mosier, K., 2004. Brain activation during vaginocervical self-stimulation and orgasm in women with complete spinal cord injury: fMRI evidence of mediation by the vagus nerves. *Brain Res.* 1024, 77–88.
- Kondo, Y., Koizumi, T., Arai, Y., Kakeyama, M., Yamanouchi, K., 1993. Functional relationships between mesencephalic central gray and septum in regulating lordosis in female rats: effect of dual lesions. *Brain Res. Bull.* 32, 635–638.
- Kondo, Y., Shinoda, A., Yamanouchi, K., Arai, Y., 1990. Role of septum and preoptic area in regulating masculine and feminine sexual behavior in male rats. *Horm. Behav.* 24, 421–434.
- Kow, L.M., Johnson, A.E., Ogawa, S., Pfaff, D.W., 1991. Electrophysiological actions of oxytocin on hypothalamic neurons in vitro: neuropharmacological characterization and effects of ovarian steroids. *Neuroendocrinology* 54, 526–535.
- Kow, L.M., Pfaff, D.W., 1988. Transmitter and peptide actions on hypothalamic neurons in vitro: implications for lordosis. *Brain Res. Bull.* 20, 857–861.
- Kruger, T.H., Haake, P., Chereath, D., Knapp, W., Janssen, O.E., Exton, M.S., Schedlowski, M., Hartmann, U., 2003. Specificity of the neuroendocrine response to orgasm during sexual arousal in men. *J. Endocrinol.* 177, 57–64.
- Kruk, M.R., 1991. Ethology and pharmacology of hypothalamic aggression in the rat. *Neurosci. Biobehav. Rev.* 15, 527–538.
- Kruk, M.R., Van der Laan, C.E., Meelis, W., Phillips, R.E., Mos, J., Van der Poel, A.M., 1984a. Brain-stimulation induced agonistic behaviour: a novel paradigm in ethopharmacological aggression research. *Prog. Clin. Biol. Res.* 167, 157–177.
- Kruk, M.R., Van der Laan, C.E., Mos, J., Van der Poel, A.M., Meelis, W., Olivier, B., 1984b. Comparison of aggressive behaviour induced by electrical stimulation in the hypothalamus of male and female rats. *Prog. Brain Res.* 61, 303–314.
- Kruk, M.R., van der Poel, A.M., 1980. Is there evidence for a neural correlate of an aggressive behavioural system in the hypothalamus of the rat? *Prog. Brain Res.* 53, 385–390.
- Kruk, M.R., van der Poel, A.M., de Vos-Frerichs, T.P., 1979. The induction of aggressive behaviour by electrical stimulation in the hypothalamus of male rats. *Behaviour* 70, 292–322.
- Kruk, M.R., Van der Poel, A.M., Meelis, W., Hermans, J., Mostert, P.G., Mos, J., Lohman, A.H., 1983. Discriminant analysis of the localization of aggression-inducing electrode placements in the hypothalamus of male rats. *Brain Res.* 260, 61–79.
- Kruk, M.R., Westphal, K.G., Van Erp, A.M., van Asperen, J., Cave, B.J., Slater, E., de Koning, J., Haller, J., 1998. The hypothalamus: cross-roads of endocrine and behavioural regulation in grooming and aggression. *Neurosci. Biobehav. Rev.* 23, 163–177.
- Lammers, J.H., Kruk, M.R., Meelis, W., van der Poel, A.M., 1988. Hypothalamic substrates for brain stimulation-induced attack, teeth-chattering and social grooming in the rat. *Brain Res.* 449, 311–327.
- Lee, G., Gammie, S.C., 2009. GABA(A) receptor signaling in the lateral septum regulates maternal aggression in mice. *Behav. Neurosci.* 123, 1169–1177.
- Lee, H.J., Macbeth, A.H., Pagani, J.H., Young 3rd, W.S., 2009. Oxytocin: the great facilitator of life. *Prog. Neurobiol.* 88, 127–151.
- Lee, H.J., Pagani, J., Young 3rd, W.S., 2010. Using transgenic mouse models to study oxytocin's role in the facilitation of species propagation. *Brain Res.* 1364, 216–224.
- Leipheimer, R.E., Gallo, R.V., 1985. Medial preoptic area involvement in norepinephrine-induced suppression of pulsatile luteinizing hormone release in ovariectomized rats. *Neuroendocrinology* 40, 345–351.
- Leng, G., Ludwig, M., 2008. Neurotransmitters and peptides: whispered secrets and public announcements. *J. Physiol.* 586, 5625–5632.
- Leng, G., Onaka, T., Caqueneau, C., Sabatier, N., Tobin, V.A., Takayanagi, Y., 2008. Oxytocin and appetite. *Progress in Brain Res.* 170, 137–151.
- Lenz, K.M., Sengelaub, D.R., 2010. Maternal care effects on the development of a sexually dimorphic motor system: the role of spinal oxytocin. *Horm. Behav.* 58, 575–581.
- Lewis, C., McEwen, B.S., Frankfurt, M., 1995. Estrogen-induction of dendritic spines in ventromedial hypothalamus and hippocampus: effects of neonatal aromatase blockade and adult GDx. *Brain Res. Dev. Brain Res.* 87, 91–95.
- Light, K.C., Grewen, K.M., Amico, J.A., 2005. More frequent partner hugs and higher oxytocin levels are linked to lower blood pressure and heart rate in premenopausal women. *Biol. Psychol.* 69, 5–21.
- Lin, D., Boyle, M.P., Dollar, P., Lee, H., Lein, E.S., Perona, P., Anderson, D.J., 2011. Functional identification of an aggression locus in the mouse hypothalamus. *Nature* 470, 221–226.
- Lincoln, D.W., Paisley, A.C., 1982. Neuroendocrine control of milk ejection. *J. Reprod. Fertil.* 65, 571–586.
- Lincoln, D.W., Wakerley, J.B., 1975a. Factors governing the periodic activation of supraoptic and paraventricular neurosecretory cells during suckling in the rat. *J. Physiol.* 250, 443–461.
- Lincoln, D.W., Wakerley, J.B., 1975b. Neurosecretory activation in the rat: correlation of the suckling stimulus with the pulsatile release of oxytocin. *J. Physiol.* 245, 42P–43P.
- Linden, A., 1989. Role of cholecystokinin in feeding and lactation. *Acta Physiol. Scand. Suppl.* 585 (i–vii), 1–49.
- Liu, J., Rebar, R.W., Yen, S.S., 1983. Neuroendocrine control of the postpartum period. *Clin. Perinatol.* 10, 723–736.
- Liu, Y.C., Salamone, J.D., Sachs, B.D., 1997. Impaired sexual response after lesions of the paraventricular nucleus of the hypothalamus in male rats. *Behav. Neurosci.* 111, 1361–1367.
- Lonstein, J.S., Simmons, D.A., Stern, J.M., 1998. Functions of the caudal periaqueductal gray in lactating rats: kyphosis, lordosis, maternal aggression, and fearfulness. *Behav. Neurosci.* 112, 1502–1518.
- Looney, C., Thor, K.B., Ricca, D., Marson, L., 2005. Differential effects of simultaneous or sequential administration of paroxetine and WAY-100,635 on ejaculatory behavior. *Pharmacol., Biochem., Behav.* 82, 427–433.
- Ludwig, M., Leng, G., 2006. Dendritic peptide release and peptide-dependent behaviours. *Nat. Rev. Neurosci.* 7, 126–136.
- Luiten, P.G., Koolhaas, J.M., de Boer, S., Koopmans, S.J., 1985. The cortico-medial amygdala in the central nervous system organization of agonistic behavior. *Brain Res.* 332, 283–297.
- Lukas, M., Toth, I., Reber, S.O., Slattery, D.A., Veenema, A.H., Neumann, I.D., 2011. The neuropeptide oxytocin facilitates pro-social behavior and prevents social avoidance in rats and mice. *Neuropsychopharmacol.: Off. Publ. Am. Coll. Neuropsychopharmacol.* 36, 2159–2168.
- Lunn, P.G., 1992. Breast-feeding patterns, maternal milk output and lactational infecundity. *J. Biosoc. Sci.* 24, 317–324.
- Macdonald, K., Macdonald, T.M., 2010. The peptide that binds: a systematic review of oxytocin and its prosocial effects in humans. *Harvard Rev. Psychiatry* 18, 1–21.
- Madeira, M.D., Ferreira-Silva, L., Paula-Barbosa, M.M., 2001. Influence of sex and estrus cycle on the sexual dimorphisms of the hypothalamic ventromedial nucleus: stereological evaluation and Golgi study. *J. Comp. Neurol.* 432, 329–345.
- Magon, N., Kalra, S., 2011. The orgasmic history of oxytocin: love, lust, and labor. *Indian J. Endocrinol. Metab.* 15 (Suppl 3), S156–S161.
- Maguire, S., O'Dell, A., Touyz, L., Russell, J., 2013. Oxytocin and anorexia nervosa: a review of the emerging literature. *Eur. Eat. Disord. Rev.: J. Eat. Disord. Assoc.* 21, 475–478.
- Mahalati, K., Okanoya, K., Witt, D.M., Carter, C.S., 1991. Oxytocin inhibits male sexual behavior in prairie voles. *Pharmacol., Biochem. Behav.* 39, 219–222.

- Mann, G.E., Lammings, G.E., 2001. Relationship between maternal endocrine environment, early embryo development and inhibition of the luteolytic mechanism in cows. *Reproduction* 121, 175–180.
- Manning, M., Misicka, A., Olma, A., Bankowski, K., Stoev, S., Chini, B., Durroux, T., Mouillac, B., Corbani, M., Guillon, G., 2012. Oxytocin and vasopressin agonists and antagonists as research tools and potential therapeutics. *J. Neuroendocrinol.* 24, 609–628.
- Marson, L., 1995. Central nervous system neurons identified after injection of pseudorabies virus into the rat clitoris. *Neurosci. Lett.* 190, 41–44.
- Marson, L., McKenna, K.E., 1996. CNS cell groups involved in the control of the ischiocavernosus and bulbospongiosus muscles: a transneuronal tracing study using pseudorabies virus. *J. Comp. Neurol.* 374, 161–179.
- Marson, L., Platt, K.B., McKenna, K.E., 1993. Central nervous system innervation of the penis as revealed by the transneuronal transport of pseudorabies virus. *Neuroscience* 55, 263–280.
- Mathews, D., Edwards, D.A., 1977a. Involvement of the ventromedial and anterior hypothalamic nuclei in the hormonal induction of receptivity in the female rat. *Physiol. Behav.* 19, 319–326.
- Mathews, D., Edwards, D.A., 1977b. The ventromedial nucleus of the hypothalamus and the hormonal arousal of sexual behaviors in the female rat. *Horm. Behav.* 8, 40–51.
- Matsumoto, A., Arai, Y., 1983. Sex difference in volume of the ventromedial nucleus of the hypothalamus in the rat. *Endocrinol. Jpn.* 30, 277–280.
- Mayer, A.D., Rosenblatt, J.S., 1987. Hormonal factors influence the onset of maternal aggression in laboratory rats. *Horm. Behav.* 21, 253–267.
- McCarthy, M.M., 1995. Estrogen modulation of oxytocin and its relation to behavior. *Adv. Exp. Med. Biol.* 395, 235–245.
- McCarthy, M.M., Bare, J.E., vom Saal, F.S., 1986. Infanticide and parental behavior in wild female house mice: effects of ovariectomy, adrenalectomy and administration of oxytocin and prostaglandin F₂ alpha. *Physiol. Behav.* 36, 17–23.
- McCarthy, M.M., Curran, G.H., Feder, H.H., 1991a. Excitatory amino acid modulation of lordosis in the rat. *Neurosci. Lett.* 126, 94–97.
- McCarthy, M.M., Kaufman, L.C., Brooks, P.J., Pfaff, D.W., Schwartz-Giblin, S., 1995. Estrogen modulation of mRNA levels for the two forms of glutamic acid decarboxylase (GAD) in female rat brain. *J. Comp. Neurol.* 360, 685–697.
- McCarthy, M.M., Kleopoulos, S.P., Mobbs, C.V., Pfaff, D.W., 1994. Infusion of antisense oligodeoxynucleotides to the oxytocin receptor in the ventromedial hypothalamus reduces estrogen-induced sexual receptivity and oxytocin receptor binding in the female rat. *Neuroendocrinology* 59, 432–440.
- McCarthy, M.M., Konkle, A.T., 2005. When is a sex difference not a sex difference? *Front. Neuroendocrinol.* 26, 85–102.
- McCarthy, M.M., Kow, L.M., Pfaff, D.W., 1992. Speculations concerning the physiological significance of central oxytocin in maternal behavior. *Ann. N. Y. Acad. Sci.* 652, 70–82.
- McCarthy, M.M., Malik, K.F., Feder, H.H., 1990. Increased GABAergic transmission in medial hypothalamus facilitates lordosis but has the opposite effect in preoptic area. *Brain Res.* 507, 40–44.
- McCarthy, M.M., Masters, D.B., Fiber, J.M., Lopez-Colome, A.M., Beyer, C., Komisaruk, B.R., Feder, H.H., 1991b. GABAergic control of receptivity in the female rat. *Neuroendocrinology* 53, 473–479.
- McEwen, B.S., 1988. Genomic regulation of sexual behavior. *J. Steroid Biochem.* 30, 179–183.
- McNeilly, A.S., 2001. Neuroendocrine changes and fertility in breast-feeding women. *Prog. Brain Res.* 133, 207–214.
- Melin, P., Kihlstrom, J.E., 1963. Influence of oxytocin on sexual behavior in male rabbits. *Endocrinology* 73, 433–435.
- Melis, M.R., Argiolas, A., 2011. Central control of penile erection: a re-visitation of the role of oxytocin and its interaction with dopamine and glutamic acid in male rats. *Neurosci. Biobehav. Rev.* 35, 939–955.
- Melis, M.R., Argiolas, A., Gessa, G.L., 1986. Oxytocin-induced penile erection and yawning: site of action in the brain. *Brain Res.* 398, 259–265.
- Melis, M.R., Melis, T., Cocco, C., Succu, S., Sanna, F., Pillolla, G., Boi, A., Ferri, G.L., Argiolas, A., 2007. Oxytocin injected into the ventral tegmental area induces penile erection and increases extracellular dopamine in the nucleus accumbens and paraventricular nucleus of the hypothalamus of male rats. *Eur. J. Neurosci.* 26, 1026–1035.
- Melis, M.R., Spano, M.S., Succu, S., Argiolas, A., 1999. The oxytocin antagonist d(CH₂)⁵Tyr(Me)²-Orn⁸-vasotocin reduces non-contact penile erections in male rats. *Neurosci. Lett.* 265, 171–174.
- Melis, M.R., Succu, S., Sanna, F., Boi, A., Argiolas, A., 2009. Oxytocin injected into the ventral subiculum or the posteromedial cortical nucleus of the amygdala induces penile erection and increases extracellular dopamine levels in the nucleus accumbens of male rats. *Eur. J. Neurosci.* 30, 1349–1357.
- Mendelson, S.D., Gorzalka, B.B., 1985. A facilitatory role for serotonin in the sexual behavior of the female rat. *Pharmacol. Biochem. Behav.* 22, 1025–1033.
- Mens, W.B., Witter, A., van Wimersma Greidanus, T.B., 1983. Penetration of neurohypophyseal hormones from plasma into cerebrospinal fluid (CSF): half-times of disappearance of these neuropeptides from CSF. *Brain Res.* 262, 143–149.
- Menzies, J.R., Skibicka, K.P., Dickson, S.L., Leng, G., 2012. Neural substrates underlying interactions between appetite stress and reward. *Obes. Facts* 5, 208–220.
- Meyer-Lindenberg, A., 2008. Impact of prosocial neuropeptides on human brain function. *Prog. Brain Res.* 170, 463–470.
- Millhouse, O.E., 1973a. Certain ventromedial hypothalamic afferents. *Brain Res.* 55, 89–105.
- Millhouse, O.E., 1973b. The organization of the ventromedial hypothalamic nucleus. *Brain Res.* 55, 71–87.
- Millhouse, O.E., 1978. Cytological observations on the ventromedial hypothalamic nucleus. *Cell Tissue Res.* 191, 473–491.
- Miyamichi, K., Amat, F., Moussavi, F., Wang, C., Wickersham, I., Wall, N.R., Taniguchi, H., Tasic, B., Huang, Z.J., He, Z., Callaway, E.M., Horowitz, M.A., Luo, L., 2011. Cortical representations of olfactory input by trans-synaptic tracing. *Nature* 472, 191–196.
- Mizusawa, H., Hedlund, P., Andersson, K.E., 2002. alpha-Melanocyte stimulating hormone and oxytocin induced penile erections, and intracavernous pressure increases in the rat. *J. Urol.* 167, 757–760.
- Moody, K.M., Adler, N.T., 1995. The role of the uterus and cervix in systemic oxytocin-PGE₂ facilitated lordosis behavior. *Horm. Behav.* 29, 571–580.
- Moody, K.M., Steinman, J.L., Komisaruk, B.R., Adler, N.T., 1994. Pelvic neurectomy blocks oxytocin-facilitated sexual receptivity in rats. *Physiol. Behav.* 56, 1057–1060.
- Moos, F., Poulain, D.A., Rodriguez, F., Guerne, Y., Vincent, J.D., Richard, P., 1989. Release of oxytocin within the supraoptic nucleus during the milk ejection reflex in rats. *Exp. Brain Res.* 76, 593–602.
- Moos, F., Richard, P., 1975. [Level of oxytocin release induced by vaginal dilatation (Ferguson reflex) and vagal stimulation (vago-pituitary reflex) in lactating rats (author's transl)]. *J. Physiol.* 70, 307–314.
- Moos, F., Richard, P., 1989. Paraventricular and supraoptic bursting oxytocin cells in rat are locally regulated by oxytocin and functionally related. *J. Physiol.* 408, 1–18.
- Mos, J., Olivier, B., Lammers, J.H., van der Poel, A.M., Kruk, M.R., Zethof, T., 1987. Postpartum aggression in rats does not influence threshold currents for EBS-induced aggression. *Brain Res.* 404, 263–266.
- Moscovice, L.R., Ziegler, T.E., 2012. Peripheral oxytocin in female baboons relates to estrous state and maintenance of sexual consortships. *Horm. Behav.* 62, 592–597.
- Moss, R.L., Foreman, M.M., 1976. Potentiation of lordosis behavior by intrahypothalamic infusion of synthetic luteinizing hormone-releasing hormone. *Neuroendocrinology* 20, 176–181.
- Motta, S.C., Guimaraes, C.C., Furigo, I.C., Sukikara, M.H., Baldo, M.V., Lonstein, J.S., Canteras, N.S., 2013. Ventral premammillary nucleus as a critical sensory relay to the maternal aggression network. *Proc. Natl. Acad. Sci. USA* 110, 14438–14443.
- Mullis, K., Kay, K., Williams, D.L., 2013. Oxytocin action in the ventral tegmental area affects sucrose intake. *Brain Res.* 1513, 85–91.
- Murphy, A.Z., Hoffman, G.E., 2001. Distribution of gonadal steroid receptor-containing neurons in the preoptic-periaqueductal gray-brainstem pathway: a potential circuit for the initiation of male sexual behavior. *J. Comp. Neurol.* 438, 191–212.
- Murphy, M.R., Seckl, J.R., Burton, S., Checkley, S.A., Lightman, S.L., 1987. Changes in oxytocin and vasopressin secretion during sexual activity in men. *J. Clin. Endocrinol. Metab.* 65, 738–741.
- Naber, F., van Ijzendoorn, M.H., Deschamps, P., van Engeland, H., Bakermans-Kranenburg, M.J., 2010. Intranasal oxytocin increases fathers' observed responsiveness during play with their children: a double-blind within-subject experiment. *Psychoneuroendocrinology* 35, 1583–1586.
- Nagayama, S., Enerva, A., Fletcher, M.L., Masurkar, A.V., Igarashi, K.M., Mori, K., Chen, W.R., 2010. Differential axonal projection of mitral and tufted cells in the mouse main olfactory system. *Front. Neural Circuits*, 4.
- Neumann, I.D., 2001. Alterations in behavioral and neuroendocrine stress coping strategies in pregnant, parturient and lactating rats. *Prog. Brain Res.* 133, 143–152.
- Neumann, I.D., 2002. Involvement of the brain oxytocin system in stress coping: interactions with the hypothalamo-pituitary-adrenal axis. *Prog. Brain Res.* 139, 147–162.
- Neumann, I.D., 2003. Brain mechanisms underlying emotional alterations in the peripartum period in rats. *Depress Anxiety* 17, 111–121.
- Neumann, I.D., 2008. Brain oxytocin: a key regulator of emotional and social behaviours in both females and males. *J. Neuroendocrinol.* 20, 858–865.
- Neumann, I.D., Bosch, O.J., Toschi, N., Torner, L., Douglas, A.J., 2003. No stress response of the hypothalamo-pituitary-adrenal axis in parturient rats: lack of involvement of brain oxytocin. *Endocrinology* 144, 2473–2479.
- Neumann, I.D., Landgraf, R., 2012. Balance of brain oxytocin and vasopressin: implications for anxiety, depression, and social behaviors. *Trends Neurosci.* 35, 649–659.
- Neumann, I.D., Toschi, N., Ohl, F., Torner, L., Kromer, S.A., 2001. Maternal defence as an emotional stressor in female rats: correlation of neuroendocrine and behavioural parameters and involvement of brain oxytocin. *Eur. J. Neurosci.* 13, 1016–1024.
- Neumann, I.D., Maloumby, R., Beiderbeck, D.I., Lukas, M., Landgraf, R., 2013. Increased brain and plasma oxytocin after nasal and peripheral administration in rats and mice. *Psychoneuroendocrinology* 38, 1985–1993.
- Nieuwenhuys, R., Geeraedts, L.M., Veening, J.G., 1982. The medial forebrain bundle of the rat. I. Gen. Introd. *J. Comp. Neurol.* 206, 49–81.
- Nishimori, K., Young, L.J., Guo, Q., Wang, Z., Insel, T.R., Matzuk, M.M., 1996. Oxytocin is required for nursing but is not essential for parturition or reproductive behavior. *Proc. Natl. Acad. Sci. USA* 93, 11699–11704.
- Nishitani, S., Moriya, T., Kondo, Y., Sakuma, Y., Shinohara, K., 2004. Induction of Fos immunoreactivity in oxytocin neurons in the paraventricular nucleus after female odor exposure in male rats: effects of sexual experience. *Cell. Mol. Neurobiol.* 24, 283–291.

- Normandin, J.J., Murphy, A.Z., 2011. Somatic genital reflexes in rats with a nod to humans: anatomy, physiology, and the role of the social neuropeptides. *Horm. Behav.* 59, 656–665.
- Northrop, L.E., Erskine, M.S., 2008. Selective oxytocin receptor activation in the ventrolateral portion of the ventromedial hypothalamus is required for mating-induced pseudopregnancy in the female rat. *Endocrinology* 149, 836–842.
- Numan, M., 2007. Motivational systems and the neural circuitry of maternal behavior in the rat. *Dev. Psychobiol.* 49, 12–21.
- Numan, M., Numan, M.J., Schwarz, J.M., Neuner, C.M., Flood, T.F., Smith, C.D., 2005. Medial preoptic area interactions with the nucleus accumbens-ventral pallidum circuit and maternal behavior in rats. *Behav. Brain Res.* 158, 53–68.
- Numan, M., Sheehan, T.P., 1997. Neuroanatomical circuitry for mammalian maternal behavior. *Ann. N. Y. Acad. Sci.* 807, 101–125.
- Nyuyki, K.D., Waldherr, M., Baeuml, S., Neumann, I.D., 2011. Yes, I am ready now: differential effects of paced versus unpaced mating on anxiety and central oxytocin release in female rats. *PLoS One* 6, e23599.
- O'Connell, L.A., Matthews, B.J., Hofmann, H.A., 2012. Isotocin regulates paternal care in a monogamous cichlid fish. *Horm. Behav.* 61, 725–733.
- Oboh, A.M., Paredes, R.G., Baum, M.J., 1995. A sex comparison of increments in FOS immunoreactivity in forebrain neurons of gonadectomized, testosterone-treated rats after mounting an estrous female. *Neurobiol. Learn. Mem.* 63, 66–73.
- Ogawa, S., Kow, L.-M., McCarthy, M.M., Pfaff, D., Schwartz-Giblin, S., W., 1991. Midbrain PAG control of female reproductive behavior: in vitro electrophysiological characterization of lordosis-relevant substances. In: Depaulis, A., Bandler, R. (Eds.), *The Midbrain Periaqueductal Gray Matter*. Plenum Press, New York, pp. 211–235.
- Ogawa, S., Kow, L.M., Pfaff, D.W., 1992. Effects of lordosis-relevant neuropeptides on midbrain periaqueductal gray neuronal activity in vitro. *Peptides* 13, 965–975.
- Ogawa, S., Lubahn, D.B., Korach, K.S., Pfaff, D.W., 1997. Behavioral effects of estrogen receptor gene disruption in male mice. *Proc. Natl. Acad. Sci. USA* 94, 1476–1481.
- Olazabal, D.E., Ferreira, A., 1997. Maternal behavior in rats with kainic acid-induced lesions of the hypothalamic paraventricular nucleus. *Physiol. Behav.* 61, 779–784.
- Olazabal, D.E., Pereira, M., Agrati, D., Ferreira, A., Fleming, A.S., Gonzalez-Mariscal, G., Levy, F., Lucion, A.B., Morrell, J.L., Numan, M., Uriarte, N., 2013. New theoretical and experimental approaches on maternal motivation in mammals. *Neurosci. Biobehav. Rev.* 37, 1860–1874.
- Olivier, B., Wiepkema, P.R., 1974. Behaviour changes in mice following electrolytic lesions in the median hypothalamus. *Brain Res.* 65, 521–524.
- Olster, D.H., Blaustein, J.D., 1991. Progesterone facilitates lordosis, but not LH release, in estradiol pulse-primed male rats. *Physiol. Behav.* 50, 237–242.
- Olszewski, P.K., Klockars, A., Schioth, H.B., Levine, A.S., 2010. Oxytocin as feeding inhibitor: maintaining homeostasis in consummatory behavior. *Pharmacol., Biochem. Behav.* 97, 47–54.
- Ono, T., Nishino, H., Sasaka, K., Muramoto, K., Yano, I., Simpson, A., 1978. Paraventricular nucleus connections to spinal cord and pituitary. *Neurosci. Lett.* 10, 141–146.
- Orr, R., Marson, L., 1998. Identification of CNS neurons innervating the rat prostate: a transneuronal tracing study using pseudorabies virus. *J. Auton. Nerv. Syst.* 72, 4–15.
- Palmer, C.W., Amundson, S.D., Brito, L.F., Waldner, C.L., Barth, A.D., 2004. Use of oxytocin and cloprostenol to facilitate semen collection by electroejaculation or transrectal massage in bulls. *Anim. Reprod. Sci.* 80, 213–223.
- Panksepp, J., 2011. The basic emotional circuits of mammalian brains: do animals have affective lives? *Neurosci. Biobehav. Rev.* 35, 1791–1804.
- Parada, M., Chamas, L., Censi, S., Coria-Avila, G., Pfau, J.G., 2010. Clitoral stimulation induces conditioned place preference and Fos activation in the rat. *Horm. Behav.* 57, 112–118.
- Parker, J.A., Bloom, S.R., 2012. Hypothalamic neuropeptides and the regulation of appetite. *Neuropharmacology* 63, 18–30.
- Parker, K.J., Lee, T.M., 2001. Social and environmental factors influence the suppression of pup-directed aggression and development of paternal behavior in captive meadow voles (*Microtus pennsylvanicus*). *J. Comp. Psychol.* 115, 331–336.
- Pattij, T., de Jong, T.R., Uitterdijk, A., Waldinger, M.D., Veening, J.G., Cools, A.R., van der Graaf, P.H., Olivier, B., 2005. Individual differences in male rat ejaculatory behaviour: searching for models to study ejaculation disorders. *Eur. J. Neurosci.* 22, 724–734.
- Pedersen, A., Tomaszycki, M.L., 2012. Oxytocin antagonist treatments alter the formation of pair relationships in zebra finches of both sexes. *Horm. Behav.* 62, 113–119.
- Pedersen, C.A., Boccia, M.L., 2002. Oxytocin maintains as well as initiates female sexual behavior: effects of a highly selective oxytocin antagonist. *Horm. Behav.* 41, 170–177.
- Pedersen, C.A., Boccia, M.L., 2006. Vasopressin interactions with oxytocin in the control of female sexual behavior. *Neuroscience* 139, 843–851.
- Pedersen, C.A., Caldwell, J.D., Walker, C., Ayers, G., Mason, G.A., 1994. Oxytocin activates the postpartum onset of rat maternal behavior in the ventral tegmental and medial preoptic areas. *Behav. Neurosci.* 108, 1163–1171.
- Perello, M., Raingo, J., 2013. Leptin activates oxytocin neurons of the hypothalamic paraventricular nucleus in both control and diet-induced obese rodents. *PLoS One* 8, e59625.
- Perry, A.N., Paramadilok, A., Cushing, B.S., 2009. Neonatal oxytocin alters subsequent estrogen receptor alpha protein expression and estrogen sensitivity in the female rat. *Behav. Brain Res.* 205, 154–161.
- Peters, L.C., Kristal, M.B., Komisaruk, B.R., 1987. Sensory innervation of the external and internal genitalia of the female rat. *Brain Res.* 408, 199–204.
- Pfaff, D.W., 1980. *Estrogens and Brain Function*. Springer Verlag, New York.
- Pfaff, D., W., 2011. *Man and Woman: An Inside Story*. Oxford University Press, USA.
- Pfaff, D., W., Sakuma, Y., 1979a. Facilitation of the lordosis reflex of female rats from the ventromedial nucleus of the hypothalamus. *J. Physiol.* 288, 189–202.
- Pfaff, D.W., Kieffer, B.L., Swanson, L.W., 2008. Mechanisms for the regulation of state changes in the central nervous system: an introduction. *Ann. N. Y. Acad. Sci.* 1129, 1–7.
- Pfaff, D.W., Lewis, C., 1974. Film analyses of lordosis in female rats. *Horm. Behav.* 5, 317–335.
- Pfaff, D.W., Sakuma, Y., 1979b. Deficit in the lordosis reflex of female rats caused by lesions in the ventromedial nucleus of the hypothalamus. *J. Physiol.* 288, 203–210.
- Pfaff, D.W., Sakuma, Y., 1979c. Deficit in the lordosis reflex of female rats caused by lesions in the ventromedial nucleus of the hypothalamus. *J. Physiol.* 288, 203–210.
- Pfaff, D.W., Sakuma, Y., 1979d. Facilitation of the lordosis reflex of female rats from the ventromedial nucleus of the hypothalamus. *J. Physiol.* 288, 189–202.
- Pfau, J.G., 2009. Pathways of sexual desire. *J. Sex. Med.* 6, 1506–1533.
- Pfau, J.G., Damsma, G., Wenkster, D., Fibiger, H.C., 1995. Sexual activity increases dopamine transmission in the nucleus accumbens and striatum of female rats. *Brain Res.* 693, 21–30.
- Pfau, J.G., Heeb, M.M., 1997. Implications of immediate-early gene induction in the brain following sexual stimulation of female and male rodents. *Brain Res. Bull.* 44, 397–407.
- Pfau, J.G., Jakob, A., Kleopoulos, S.P., Gibbs, R.B., Pfaff, D.W., 1994. Sexual stimulation induces Fos immunoreactivity within GnRH neurons of the female rat preoptic area: interaction with steroid hormones. *Neuroendocrinology* 60, 283–290.
- Pfau, J.G., Kippin, T.E., Coria-Avila, G.A., Gelez, H., Afonso, V.M., Ismail, N., Parada, M., 2012. Who, what, where, when (and maybe even why)? How the experience of sexual reward connects sexual desire, preference, and performance. *Arch. Sex. Behav.* 41, 31–62.
- Pfau, J.G., Kleopoulos, S.P., Mobbs, C.V., Gibbs, R.B., Pfaff, D.W., 1993. Sexual stimulation activates c-fos within estrogen-concentrating regions of the female rat forebrain. *Brain Res.* 624, 253–267.
- Pfau, J.G., Manitt, C., Coopersmith, C.B., 2006. Effects of pelvic, pudendal, or hypogastric nerve cuts on Fos induction in the rat brain following vaginocervical stimulation. *Physiol. Behav.* 89, 627–636.
- Pfau, J.G., Marcangione, C., Smith, W.J., Manitt, C., Abillamaa, H., 1996. Differential induction of Fos in the female rat brain following different amounts of vaginocervical stimulation: modulation by steroid hormones. *Brain Res.* 741, 314–330.
- Pfau, J.G., Smith, W.J., Byrne, N., Stephens, G., 2000. Appetitive and consummatory sexual behaviors of female rats in bilevel chambers. II. Patterns of estrus termination following vaginocervical stimulation. *Horm. Behav.* 37, 96–107.
- Pfau, J.G., Smith, W.J., Coopersmith, C.B., 1999. Appetitive and consummatory sexual behaviors of female rats in bilevel chambers. I. A correlational and factor analysis and the effects of ovarian hormones. *Horm. Behav.* 35, 224–240.
- Pfeifle, J.K., Shivers, M., Edwards, D.A., 1980. Parasagittal hypothalamic knife cuts and sexual receptivity in the female rat. *Physiol. Behav.* 24, 145–150.
- Pleim, E.T., Barfield, R.J., 1988. Progesterone versus estrogen facilitation of female sexual behavior by intracranial administration to female rats. *Horm. Behav.* 22, 150–159.
- Pleim, E.T., Baumann, J., Barfield, R.J., 1991. A contributory role for midbrain progesterone in the facilitation of female sexual behavior in rats. *Horm. Behav.* 25, 19–28.
- Pleim, E.T., Brown, T.J., MacLusky, N.J., Etgen, A.M., Barfield, R.J., 1989. Dilute estradiol implants and progestin receptor induction in the ventromedial nucleus of the hypothalamus: correlation with receptive behavior in female rats. *Endocrinology* 124, 1807–1812.
- Polston, E.K., Erskine, M.S., 1995. Patterns of induction of the immediate-early genes c-fos and egr-1 in the female rat brain following differential amounts of mating stimulation. *Neuroendocrinology* 62, 370–384.
- Ragnauth, A.K., Devidze, N., Moy, V., Finley, K., Goodwillie, A., Kow, L.M., Muglia, L.J., Pfaff, D.W., 2005. Female oxytocin gene-knockout mice, in a semi-natural environment, display exaggerated aggressive behavior. *Genes, Brain, Behav.* 4, 229–239.
- Rajendren, G., Dudley, C.A., Moss, R.L., 1991. Role of the ventromedial nucleus of hypothalamus in the male-induced enhancement of lordosis in female rats. *Physiol. Behav.* 50, 705–710.
- Ramos, E., Castilla, A., Navarro, N., Monasterio, N., Mena, F., Morales, T., 2008. Suckling-induced oxytocin increase in the spinal cord of the rat. *Brain Res.* 1236, 85–92.
- Rhodes, C.H., Morrell, J.L., Pfaff, D.W., 1981. Immunohistochemical analysis of magnocellular elements in rat hypothalamus: distribution and numbers of cells containing neurophysin, oxytocin, and vasopressin. *J. Comp. Neurol.* 198, 45–64.
- Robinson, I.C., Jones, P.M., 1982. Oxytocin and neurophysin in plasma and CSF during suckling in the guinea-pig. *Neuroendocrinology* 34, 59–63.
- Roeling, T.A., Veening, J.G., Kruk, M.R., Peters, J.P., Vermelis, M.E., Nieuwenhuys, R., 1994. Efferent connections of the hypothalamic “aggression area” in the rat. *Neuroscience* 59, 1001–1024.

- Romano, A., Potes, C.S., Tempesta, B., Cassano, T., Cuomo, V., Lutz, T., Gaetani, S., 2013. Hindbrain noradrenergic input to the hypothalamic PVN mediates the activation of oxytocinergic neurons induced by the satiety factor oleylethanolamide. *Am. J. Physiol. Endocrinol. Metab.* 305, E1266–E1273.
- Romano, G.J., Harlan, R.E., Shivers, B.D., Howells, R.D., Pfaff, D.W., 1988. Estrogen increases proenkephalin messenger ribonucleic acid levels in the ventromedial hypothalamus of the rat. *Mol. Endocrinol.* 2, 1320–1328.
- Romano, G.J., Mobbs, C.V., Lauber, A., Howells, R.D., Pfaff, D.W., 1990. Differential regulation of proenkephalin gene expression by estrogen in the ventromedial hypothalamus of male and female rats: implications for the molecular basis of a sexually differentiated behavior. *Brain Res.* 536, 63–68.
- Rosenblatt, J.S., Mayer, A.D., Giordano, A.L., 1988. Hormonal basis during pregnancy for the onset of maternal behavior in the rat. *Psychoneuroendocrinology* 13, 29–46.
- Ross, H.E., Cole, C.D., Smith, Y., Neumann, I.D., Landgraf, R., Murphy, A.Z., Young, L.J., 2009. Characterization of the oxytocin system regulating affiliative behavior in female prairie voles. *Neuroscience* 162, 892–903.
- Ross, H.E., Young, L.J., 2009. Oxytocin and the neural mechanisms regulating social cognition and affiliative behavior. *Front. Neuroendocrinol.* 30, 534–547.
- Rossoni, E., Feng, J., Tirozzi, B., Brown, D., Leng, G., Moos, F., 2008. Emergent synchronous bursting of oxytocin neuronal network. *PLoS Comput. Biol.* 4, e1000123.
- Rowe, D.W., Erskine, M.S., 1993. c-Fos proto-oncogene activity induced by mating in the preoptic area, hypothalamus and amygdala in the female rat: role of afferent input via the pelvic nerve. *Brain Res.* 621, 25–34.
- Rubin, B.S., Barfield, R.J., 1983. Induction of estrous behavior in ovariectomized rats by sequential replacement of estrogen and progesterone to the ventromedial hypothalamus. *Neuroendocrinology* 37, 218–224.
- Rupp, H.A., James, T.W., Ketterson, E.D., Sengelaub, D.R., Ditzen, B., Heiman, J.R., 2013. Lower sexual interest in postpartum women: relationship to amygdala activation and intranasal oxytocin. *Horm. Behav.* 63, 114–121.
- Russell, J.A., Douglas, A.J., Ingram, C.D., 2001. Brain preparations for maternity—adaptive changes in behavioral and neuroendocrine systems during pregnancy and lactation. An overview. *Prog. Brain Res.* 133, 1–38.
- Sa, S.I., Madeira, M.D., 2005. Estrogen modulates the sexually dimorphic synaptic connectivity of the ventromedial nucleus. *J. Comp. Neurol.* 484, 68–79.
- Sabatier, N., Leng, G., Menzies, J., 2013. Oxytocin, feeding, and satiety. *Front. Endocrinol.* 4, 35.
- Saito, A., Nakamura, K., 2011. Oxytocin changes primate paternal tolerance to offspring in food transfer. *J. Comp. Physiol. A, Neuroethol., Sens., Neural, Behav. Physiol.* 197, 329–337.
- Sakuma, Y., Pfaff, D., 1981. Electrophysiologic determination of projections from ventromedial hypothalamus to midbrain central gray: differences between female and male rats. *Brain Res.* 225, 184–188.
- Sakuma, Y., Pfaff, D.W., 1979. Facilitation of female reproductive behavior from mesencephalic central gray in the rat. *Am. J. Physiol.* 237, R278–R284.
- Sakuma, Y., Pfaff, D.W., 1980. LH-RH in the mesencephalic central grey can potentiate lordosis reflex of female rats. *Nature* 283, 566–567.
- Sakuma, Y., Pfaff, D.W., 1982. Properties of ventromedial hypothalamic neurons with axons to midbrain central gray. *Exp. Brain Res. Exp. Hirnforsch. Exp. cerebr.* 46, 292–300.
- Salonia, A., Nappi, R.E., Pontillo, M., Daverio, R., Smeraldi, A., Briganti, A., Fabbri, F., Zanni, G., Rigatti, P., Montorsi, F., 2005. Menstrual cycle-related changes in plasma oxytocin are relevant to normal sexual function in healthy women. *Horm. Behav.* 47, 164–169.
- Samson, W.K., McCann, S.M., Chud, L., Dudley, C.A., Moss, R.L., 1980. Intra- and extrahypothalamic luteinizing hormone-releasing hormone (LHRH) distribution in the rat with special reference to mesencephalic sites which contain both LHRH and single neurons responsive to LHRH. *Neuroendocrinology* 31, 66–72.
- Sansone, G.R., Komisaruk, B.R., 2001. Evidence that oxytocin is an endogenous stimulator of autonomic sympathetic preganglionic: the pupillary dilatation response to vaginocervical stimulation in the rat. *Brain Res.* 898, 265–271.
- Saper, C.B., 2011. Animal behaviour: the nexus of sex and violence. *Nature* 470, 179–181.
- Sawchenko, P.E., Swanson, L.W., 1982a. Immunohistochemical identification of neurons in the paraventricular nucleus of the hypothalamus that project to the medulla or to the spinal cord in the rat. *J. Comp. Neurol.* 205, 260–272.
- Sawchenko, P.E., Swanson, L.W., 1982b. Immunohistochemical identification of neurons in the paraventricular nucleus of the hypothalamus that project to the medulla or to the spinal cord in the rat. *J. Comp. Neurol.* 205, 260–272.
- Schneiderman, I., Zagoory-Sharon, O., Leckman, J.F., Feldman, R., 2012. Oxytocin during the initial stages of romantic attachment: relations to couples' interactive reciprocity. *Psychoneuroendocrinology* 37, 1277–1285.
- Schulze, H.G., Gorzalka, B.B., 1991. Oxytocin effects on lordosis frequency and lordosis duration following infusion into the medial pre-optic area and ventromedial hypothalamus of female rats. *Neuropeptides* 18, 99–106.
- Schulze, H.G., Gorzalka, B.B., 1992. Low concentrations of oxytocin suppress lordosis when infused into the lateral ventricle of female rats. *Endocr. Regul.* 26, 23–27.
- Schumacher, M., Coirini, H., Frankfurt, M., McEwen, B.S., 1989. Localized actions of progesterone in hypothalamus involve oxytocin. *Proc. Natl. Acad. Sci. USA* 86, 6798–6801.
- Schumacher, M., Coirini, H., Pfaff, D.W., McEwen, B.S., 1990. Behavioral effects of progesterone associated with rapid modulation of oxytocin receptors. *Science* 250, 691–694.
- Schumacher, M., Coirini, H., Pfaff, D.W., McEwen, B.S., 1991. Light-dark differences in behavioral sensitivity to oxytocin. *Behav. Neurosci.* 105, 487–492.
- Schwanzel-Fukuda, M., Morrell, J.I., Pfaff, D.W., 1984. Localization of forebrain neurons which project directly to the medulla and spinal cord of the rat by retrograde tracing with wheat germ agglutinin. *J. Comp. Neurol.* 226, 1–20.
- Segovia, S., Garcia-Falgueras, A., Perez-Laso, C., Pinos, H., Carrillo, B., Collado, P., Claro, F., Guillemon, A., 2009. The effects of partial and complete masculinization on the sexual differentiation of nuclei that control lordotic behavior in the male rat. *Behav. Brain Res.* 196, 261–267.
- Shelley, D.N., Choleris, E., Kavaliers, M., Pfaff, D.W., 2006. Mechanisms underlying sexual and affiliative behaviors of mice: relation to generalized CNS arousal. *Soc. Cogn. Affect. Neurosci.* 1, 260–270.
- Siegel, A., Roeling, T.A., Gregg, T.R., Kruk, M.R., 1999. Neuropharmacology of brain-stimulation-evoked aggression. *Neurosci. Biobehav. Rev.* 23, 359–389.
- Silva, B.A., Mattucci, C., Krzykowski, P., Murana, E., Illarionova, A., Grinevich, V., Canteras, N.S., Ragozzino, D., Gross, C.T., 2013. Independent hypothalamic circuits for social and predator fear. *Nat. Neurosci.* 16, 1731–1733.
- Sirinathsinghi, D.J., 1985. Modulation of lordosis behaviour in the female rat by corticotropin releasing factor, beta-endorphin and gonadotropin releasing hormone in the mesencephalic central gray. *Brain Res.* 336, 45–55.
- Sirinathsinghi, D.J., 1986. Regulation of lordosis behaviour in the female rat by corticotropin-releasing factor, beta-endorphin/corticotropin and luteinizing hormone-releasing hormone neuronal systems in the medial preoptic area. *Brain Res.* 375, 49–56.
- Snoeren, E.M., Veening, J.G., Olivier, B., Oosting, R.S., 2013a. Serotonin 1A receptors and sexual behavior in female rats: a review. *Pharmacol., Biochem., Behav.*
- Snoeren, E.M., Veening, J.G., Olivier, B., Oosting, R.S., 2013b. Serotonin 1A receptors and sexual behavior in male rats: a review. *Pharmacol., Biochem., Behav.*
- Snowdon, C.T., Pieper, B.A., Boe, C.Y., Cronin, K.A., Kurian, A.V., Ziegler, T.E., 2010. Variation in oxytocin is related to variation in affiliative behavior in monogamous, pairbonded tamarins. *Horm. Behav.* 58, 614–618.
- Sobrinho, L.G., 1993. The psychogenic effects of prolactin. *Acta Endocrinol.* 129 (Suppl 1), S38–S40.
- Song, Z., Tai, F., Yu, C., Wu, R., Zhang, X., Broders, H., He, F., Guo, R., 2010. Sexual or paternal experiences alter alloparental behavior and the central expression of ERalpha and OT in male mandarin voles (*Microtus mandarinus*). *Behav. Brain Res.* 214, 290–300.
- Sosulski, D.L., Bloom, M.L., Cutforth, T., Axel, R., Datta, S.R., 2011. Distinct representations of olfactory information in different cortical centres. *Nature* 472, 213–216.
- Stern, J.M., Kolunje, J.M., 1993. Maternal aggression of rats is impaired by cutaneous anesthesia of the ventral trunk, but not by nipple removal. *Physiol. Behav.* 54, 861–868.
- Stoneham, M.D., Everitt, B.J., Hansen, S., Lightman, S.L., Todd, K., 1985. Oxytocin and sexual behaviour in the male rat and rabbit. *J. Endocrinol.* 107, 97–106.
- Striepens, N., Kendrick, K.M., Maier, W., Hurlmann, R., 2011. Prosocial effects of oxytocin and clinical evidence for its therapeutic potential. *Front. Neuroendocrinol.* 32, 426–450.
- Summerlee, A.J., 1981. Extracellular recordings from oxytocin neurones during the expulsive phase of birth in unanaesthetized rats. *J. Physiol.* 321, 1–9.
- Summerlee, A.J., Lincoln, D.W., 1981. Electrophysiological recordings from oxytocinergic neurones during suckling in the unanaesthetized lactating rat. *J. Endocrinol.* 90, 255–265.
- Svennersten-Sjaunja, K., Olsson, K., 2005. Endocrinology of milk production. *Domest. Anim. Endocrinol.* 29, 241–258.
- Swanson, L.W., 1987. The Hypothalamus. In: Björklund, A., Hökfelt, T., Swanson, L.W. (Eds.), *Integrated Systems of the CNS*, part 1. Elsevier, Amsterdam, pp. 1–124.
- Swanson, L.W., Kuypers, H.G., 1980. The paraventricular nucleus of the hypothalamus: cytoarchitectonic subdivisions and organization of projections to the pituitary, dorsal vagal complex, and spinal cord as demonstrated by retrograde fluorescence double-labeling methods. *J. Comp. Neurol.* 194, 555–570.
- Swanson, L.W., McKellar, S., 1979a. The distribution of oxytocin- and neurophysin-stained fibers in the spinal cord of the rat and monkey. *J. Comp. Neurol.* 188, 87–106.
- Swanson, L.W., McKellar, S., 1979b. The distribution of oxytocin- and neurophysin-stained fibers in the spinal cord of the rat and monkey. *J. Comp. Neurol.* 188, 87–106.
- Takahashi, L.K., Lore, R.K., 1982. Intermale and maternal aggression in adult rats tested at different ages. *Physiol. Behav.* 29, 1013–1018.
- Tallam, S.K., Walton, J.S., Johnson, W.H., 2000. Effects of oxytocin on follicular development and duration of the estrous cycle in heifers. *Theriogenology* 53, 951–962.
- Tang, Y., Rampin, O., Calas, A., Facchinetti, P., Giuliano, F., 1998. Oxytocinergic and serotonergic innervation of identified lumbosacral nuclei controlling penile erection in the male rat. *Neuroscience* 82, 241–254.
- Tang, Y., Rampin, O., Giuliano, F., Ugolini, G., 1999. Spinal and brain circuits to motoneurons of the bulbospongiosus muscle: retrograde transneuronal tracing with rabies virus. *J. Comp. Neurol.* 414, 167–192.
- Tetel, M.J., Celentano, D.C., Blaustein, J.D., 1994a. Intraneuronal convergence of tactile and hormonal stimuli associated with female reproduction in rats. *J. Neuroendocrinol.* 6, 211–216.
- Tetel, M.J., Getzinger, M.J., Blaustein, J.D., 1993. Fos expression in the rat brain following vaginal-cervical stimulation by mating and manual probing. *J. Neuroendocrinol.* 5, 397–404.
- Tetel, M.J., Getzinger, M.J., Blaustein, J.D., 1994b. Estradiol and progesterone influence the response of ventromedial hypothalamic neurons to tactile stimuli associated with female reproduction. *Brain Res.* 646, 267–272.

- Theodosios, D.T., 2002. Oxytocin-secreting neurons: a physiological model of morphological neuronal and glial plasticity in the adult hypothalamus. *Front. Neuroendocrinol.* 23, 101–135.
- Tsukahara, S., Ezawa, N., Yamanouchi, K., 2003. Neonatal estrogen decreases neural density of the septum-midbrain central gray connection underlying the lordosis-inhibiting system in female rats. *Neuroendocrinology* 78, 226–233.
- Tsukahara, S., Yamanouchi, K., 2001. Neurohistological and behavioral evidence for lordosis-inhibiting tract from lateral septum to periaqueductal gray in male rats. *J. Comp. Neurol.* 431, 293–310.
- Tsukahara, S., Yamanouchi, K., 2002. Sex difference in septal neurons projecting axons to midbrain central gray in rats: a combined double retrograde tracing and ER-immunohistochemical study. *Endocrinology* 143, 285–294.
- Tsuneoka, Y., Maruyama, T., Yoshida, S., Nishimori, K., Kato, T., Numan, M., Kuroda, K.O., 2013. Functional, anatomical, and neurochemical differentiation of medial preoptic area subregions in relation to maternal behavior in the mouse. *J. Comp. Neurol.* 521, 1633–1663.
- Uckert, S., Becker, A.J., Ness, B.O., Stief, C.G., Scheller, F., Knapp, W.H., Jonas, U., 2003. Oxytocin plasma levels in the systemic and cavernous blood of healthy males during different penile conditions. *World J. Urol.* 20, 323–326.
- Uvnas-Moberg, K., 1994. Role of efferent and afferent vagal nerve activity during reproduction: integrating function of oxytocin on metabolism and behaviour. *Psychoneuroendocrinology* 19, 687–695.
- Uvnas-Moberg, K., 1998. Oxytocin may mediate the benefits of positive social interaction and emotions. *Psychoneuroendocrinology* 23, 819–835.
- Vaccari, C., Lolait, S.J., Ostrowski, N.L., 1998. Comparative distribution of vasopressin V1b and oxytocin receptor messenger ribonucleic acids in brain. *Endocrinology* 139, 5015–5033.
- Van de Kar, L.D., Levy, A.D., Li, Q., Brownfield, M.S., 1998. A comparison of the oxytocin and vasopressin responses to the 5-HT_{1A} agonist and potential anxiolytic drug alnespirone (S-20499). *Pharmacol., Biochem., Behav.* 60, 677–683.
- Van Houten, M., Brawer, J.R., 1978a. Cytology of neurons of the hypothalamic ventromedial nucleus in the adult male rat. *J. Comp. Neurol.* 178, 89–116.
- Van Houten, M., Brawer, J.R., 1978b. Regional variations in glia and neuropil within the hypothalamic ventromedial nucleus. *J. Comp. Neurol.* 179, 719–737.
- Veenema, A.H., Neumann, I.D., 2008. Central vasopressin and oxytocin release: regulation of complex social behaviours. *Prog. Brain Res.* 170, 261–276.
- Veening, J.G., 1975. Behavioural functions of the VMH in the rat: an ethological approach. *Zoology, Groningen, The Netherlands.*
- Veening, J.G., 1992. Brain and behaviour: morphological and functional aspects of the hypothalamus in the rat. *Eur. J. Morphol.* 30, 53–66.
- Veening, J.G., Barendregt, H.P., 2010. The regulation of brain states by neuroactive substances distributed via the cerebrospinal fluid; a review. *Cereb. Fluid Res.* 7, 1.
- Veening, J.G., Coolen, L.M., 1998. Neural activation following sexual behavior in the male and female rat brain. *Behav. Brain Res.* 92, 181–193.
- Veening, J.G., Coolen, L.M., 2013. Neural mechanisms of sexual behavior in the male rat: emphasis on ejaculation-related circuits. *Pharmacol., Biochem., Behav.*
- Veening, J.G., Coolen, L.M., de Jong, T.R., Joosten, H.W., de Boer, S.F., Koolhaas, J.M., Olivier, B., 2005. Do similar neural systems subserve aggressive and sexual behaviour in male rats? Insights from c-Fos and pharmacological studies. *Eur. J. Pharmacol.* 526, 226–239.
- Veening, J.G., Coolen, L.M., Gerrits, P.O., 2013. Neural mechanisms of female sexual behavior in the rat; comparison with male ejaculatory control. *Pharmacol., Biochem., Behav.*
- Veening, J.G., de Jong, T., Barendregt, H.P., 2010. Oxytocin-messages via the cerebrospinal fluid: behavioral effects; a review. *Physiol. Behav.* 101, 193–210.
- Veening, J.G., Gerrits, P.O., Barendregt, H.P., 2012. Volume transmission of beta-endorphin via the cerebrospinal fluid; a review. *Fluids Barriers CNS* 9, 16.
- Veening, J.G., Olivier, B., 2013. Intranasal administration of oxytocin: behavioral and clinical effects, a review. *Neurosci. Biobehav. Rev.* 37, 1445–1465.
- Veening, J.G., Swanson, L.W., Cowan, W.M., Nieuwenhuys, R., Geeraedts, L.M., 1982. The medial forebrain bundle of the rat. II. An autoradiographic study of the topography of the major descending and ascending components. *J. Comp. Neurol.* 206, 82–108.
- Vega Matuszczyk, J., Fernandez-Guasti, A., Larsson, K., 1988. Sexual orientation, proceptivity, and receptivity in the male rat as a function of neonatal hormonal manipulation. *Horm. Behav.* 22, 362–378.
- Veronneau-Longueville, F., Rampin, O., Freund-Mercier, M.J., Tang, Y., Calas, A., Marson, L., McKenna, K.E., Stoeckel, M.E., Benoit, G., Giuliano, F., 1999. Oxytocinergic innervation of autonomic nuclei controlling penile erection in the rat. *Neuroscience* 93, 1437–1447.
- Vicentic, A., Li, Q., Battaglia, G., Van de Kar, L.D., 1998. WAY-100635 inhibits 8-OH-DPAT-stimulated oxytocin, ACTH and corticosterone, but not prolactin secretion. *Eur. J. Pharmacol.* 346, 261–266.
- Vilela, F.C., Giusti-Paiva, A., 2011. Glucocorticoids disrupt neuroendocrine and behavioral responses during lactation. *Endocrinology* 152, 4838–4845.
- Wagner, C.K., Clemens, L.G., 1993. Neurophysin-containing pathway from the paraventricular nucleus of the hypothalamus to a sexually dimorphic motor nucleus in lumbar spinal cord. *J. Comp. Neurol.* 336, 106–116.
- Waldherr, M., Neumann, I.D., 2007. Centrally released oxytocin mediates mating-induced anxiolysis in male rats. *Proc. Natl. Acad. Sci. USA* 104, 16681–16684.
- Waldinger, M.D., 2013. History of premature ejaculation. In: Jannini, E.A., McMahon, C.G., Ealdinger, M.D. (Eds.), *Premature Ejaculation. From Etiology to Diagnosis and Treatment*. Springer-Verlag, Italia, Italia, pp. 5–24.
- Waldinger, M.D., 2008. Premature ejaculation: different pathophysiologies and etiologies determine its treatment. *J. Sex Marital Ther.* 34, 1–13.
- Waldinger, M.D., 2014. Ejaculatio praecox, erectio praecox, and detumescencia praecox as symptoms of a hypertonic state in lifelong premature ejaculation: a new hypothesis. *Pharmacol. Biochem. Behav.* 121, 189–194. <http://dx.doi.org/10.1016/j.pbb.2013.12.004>.
- Wang, Z.X., Liu, Y., Young, L.J., Insel, T.R., 2000. Hypothalamic vasopressin gene expression increases in both males and females postpartum in a biparental rodent. *J. Neuroendocrinol.* 12, 111–120.
- Wang, Y.F., Hamilton, K., 2009. Chronic vs. acute interactions between supraoptic oxytocin neurons and astrocytes during lactation: role of glial fibrillary acidic protein plasticity. *Sci. World J.* 9, 1308–1320.
- Wang, Y.F., Hatton, G.I., 2009. Astrocytic plasticity and patterned oxytocin neuronal activity: dynamic interactions. *J. Neurosci.: Off. J. Soc. Neurosci.* 29, 1743–1754.
- Wang, Y.F., Negoro, H., Higuchi, T., 2013. Lesions of hypothalamic mammillary body desynchronise milk-ejection bursts of rat bilateral supraoptic oxytocin neurons. *J. Neuroendocrinol.* 25, 67–75.
- Ward, I.L., 1972. Prenatal stress feminizes and demasculinizes the behavior of males. *Science* 175, 82–84.
- Ward, I.L., 1977. Exogenous androgen activates female behavior in noncopulating, prenatally stressed male rats. *J. Comp. Physiol. Psychol.* 91, 465–471.
- Ward, I.L., Franck, J.E., Crowley, W.R., 1977. Central progesterone induces female sexual behavior in estrogen-primed intact males rats. *J. Comp. Physiol. Psychol.* 91, 1416–1423.
- Weisman, O., Zagoory-Sharon, O., Feldman, R., 2012. Oxytocin administration to parent enhances infant physiological and behavioral readiness for social engagement. *Biol. Psychiatry* 72, 982–989.
- Weisz, J., Ward, I.L., 1980. Plasma testosterone and progesterone titers of pregnant rats, their male and female fetuses, and neonatal offspring. *Endocrinology* 106, 306–316.
- Wersinger, S.R., Baum, M.J., Erskine, M.S., 1993. Mating-induced FOS-like immunoreactivity in the rat forebrain: a sex comparison and a dimorphic effect of pelvic nerve transection. *J. Neuroendocrinol.* 5, 557–568.
- Whalen, R.E., Gladue, B.A., Olsen, K.L., 1986. Lordotic behavior in male rats: genetic and hormonal regulation of sexual differentiation. *Horm. Behav.* 20, 73–82.
- Whipple, B., Komisaruk, B.R., 2002. Brain (PET) responses to vaginal-cervical self-stimulation in women with complete spinal cord injury: preliminary findings. *J. Sex Marital Ther.* 28, 79–86.
- Whitney, J.B., Herrenkohl, L.R., 1977. Effects of anterior hypothalamic lesions on the sexual behavior of prenatally-stressed male rats. *Physiol. Behavior* 19, 167–169.
- Whitney, J.F., 1986. Effect of medial preoptic lesions on sexual behavior of female rats is determined by test situation. *Behav. Neurosci.* 100, 230–235.
- Williams, J.R., Carter, C.S., Insel, T., 1992a. Partner preference development in female prairie voles is facilitated by mating or the central infusion of oxytocin. *Ann. N. Y. Acad. Sci.* 652, 487–489.
- Williams, J.R., Catania, K.C., Carter, C.S., 1992b. Development of partner preferences in female prairie voles (*Microtus ochrogaster*): the role of social and sexual experience. *Behav. Neurosci.* 26, 339–349.
- Wilson, L.A., Wayman, C.P., Jackson, V.M., 2009. Neuropeptide modulation of a lumbar spinal reflex: potential implications for female sexual function. *J. Sex. Med.* 6, 947–957.
- Winslow, J.T., Insel, T.R., 1991. Social status in pairs of male squirrel monkeys determines the behavioral response to central oxytocin administration. *J. Neurosci.: Off. J. Soc. Neurosci.* 11, 2032–2038.
- Witt, D.M., Carter, C.S., Walton, D.M., 1990. Central and peripheral effects of oxytocin administration in prairie voles (*Microtus ochrogaster*). *Pharmacol., Biochem. Behav.* 37, 63–69.
- Witt, D.M., Insel, T.R., 1991. A selective oxytocin antagonist attenuates progesterone facilitation of female sexual behavior. *Endocrinology* 128, 3269–3276.
- Witt, D.M., Insel, T.R., 1992. Central oxytocin antagonism decreases female reproductive behavior. *Ann. N. Y. Acad. Sci.* 652, 445–447.
- Witt, D.M., Insel, T.R., 1994. Increased Fos expression in oxytocin neurons following masculine sexual behavior. *J. Neuroendocrinol.* 6, 13–18.
- Woller, M.J., Sosa, M.E., Chiang, Y., Prudom, S.L., Keilty, P., Moore, J.E., Ziegler, T.E., 2012. Differential hypothalamic secretion of neurocrines in male common marmosets: parental experience effects? *J. Neuroendocrinol.* 24, 413–421.
- Woodside, B., Budin, R., Wellman, M.K., Abizaid, A., 2012. Many mouths to feed: the control of food intake during lactation. *Front. Neuroendocrinol.* 33, 301–314.
- Wu, T.J., Segal, A.Z., Miller, G.M., Gibson, M.J., Silverman, A.J., 1992. FOS expression in gonadotropin-releasing hormone neurons: enhancement by steroid treatment and mating. *Endocrinology* 131, 2045–2050.
- Yanagimoto, M., Honda, K., Goto, Y., Negoro, H., 1996. Afferents originating from the dorsal penile nerve excite oxytocin cells in the hypothalamic paraventricular nucleus of the rat. *Brain Res.* 733, 292–296.
- Yang, S.P., Voegt, J.L., 2002. Mating-activated nitric oxide-producing neurons in specific brain regions in the female rat. *Brain Res.* 950, 79–87.
- Young, L.J., Murphy Young, A.Z., Hammock, E.A., 2005. Anatomy and neurochemistry of the pair bond. *J. Comp. Neurol.* 493, 51–57.
- Young 3rd, W.S., Shepard, E., Amico, J., Hennighausen, L., Wagner, K.U., LaMarca, M. E., McKinney, C., Ginns, E.I., 1996. Deficiency in mouse oxytocin prevents milk ejection, but not fertility or parturition. *J. Neuroendocrinol.* 8, 847–853.
- Yu, G.Z., Kaba, H., Okutani, F., Takahashi, S., Higuchi, T., 1996a. The olfactory bulb: a critical site of action for oxytocin in the induction of maternal behaviour in the rat. *Neuroscience* 72, 1083–1088.

- Yu, G.Z., Kaba, H., Okutani, F., Takahashi, S., Higuchi, T., Seto, K., 1996b. The action of oxytocin originating in the hypothalamic paraventricular nucleus on mitral and granule cells in the rat main olfactory bulb. *Neuroscience* 72, 1073–1082.
- Zagoory-Sharon, O., Schroeder, M., Levine, A., Moran, T.H., Weller, A., 2008. Adaptation to lactation in OLETF rats lacking CCK-1 receptors: body weight, fat tissues, leptin and oxytocin. *Int. J. Obes.* 32, 1211–1221.
- Zellner, M.R., Watt, D.F., Solms, M., Panksepp, J., 2011. Affective neuroscientific and neuropsychanalytic approaches to two intractable psychiatric problems: why depression feels so bad and what addicts really want. *Neurosci. Biobehav. Rev.* 35, 2000–2008.
- Zhang, X., Leng, G., Feng, J., 2010. Coherent peptide-mediated activity in a neuronal network controlled by subcellular signaling pathway: experiments and modeling. *J. Biotechnol.* 149, 215–225.
- Zhang, X.H., Filippi, S., Vignozzi, L., Morelli, A., Mancina, R., Luconi, M., Donati, S., Marini, M., Vannelli, G.B., Forti, G., Maggi, M., 2005. Identification, localization and functional in vitro and in vivo activity of oxytocin receptor in the rat penis. *J. Endocrinol.* 184, 567–576.
- Zhang, Y., Gray, T.S., D'Souza, D.N., Carrasco, G.A., Damjanoska, K.J., Dudas, B., Garcia, F., Zainelli, G.M., Sullivan Hanley, N.R., Battaglia, G., Muma, N.A., Van de Kar, L.D., 2004. Desensitization of 5-HT_{1A} receptors by 5-HT_{2A} receptors in neuroendocrine neurons in vivo. *J. Pharmacol. Exp. Ther.* 310, 59–66.