


# Current Biology

## Pervasive gaps in Amazonian ecological research

### Graphical abstract



### Authors

Raquel L. Carvalho,  
 Angelica F. Resende, Jos Barlow, ...,  
 Jansen Zuanon, The Synergize  
 Consortium, Joice Ferreira

### Correspondence

raqueluly@gmail.com (R.L.C.),  
 gel.florestal@gmail.com (A.F.R.),  
 jos.barlow@lancaster.ac.uk (J.B.),  
 mariormoura@gmail.com (M.R.M.)

### In brief

Carvalho et al. map the locations and drivers of ecological research across the Brazilian Amazon. Research facilities and accessibility were strong predictors of research location. Areas with the lowest probability (<0.1%) of research covered about 27.3%, 17.3%, and 54.1% of aquatic, wetland, and upland habitats, respectively.

### Highlights

- Ecological metadata were compiled for 7,694 sites across the Brazilian Amazon
- Accessibility and proximity to research facilities influenced research probability
- Knowledge gaps are greater in uplands than in wetlands and aquatic habitats
- Undersampled areas overlap predicted hotspots of climate change and deforestation



## Report

## Pervasive gaps in Amazonian ecological research

Raquel L. Carvalho,<sup>1,2,26,27,\*</sup> Angelica F. Resende,<sup>1,3,26,\*</sup> Jos Barlow,<sup>4,\*</sup> Filipe M. França,<sup>5</sup> Mario R. Moura,<sup>6,7,\*</sup> Rafaella Maciel,<sup>8</sup> Fernanda Alves-Martins,<sup>9</sup> Jack Shutt,<sup>10</sup> Cassio A. Nunes,<sup>11</sup> Fernando Elias,<sup>12</sup> Juliana M. Silveira,<sup>4</sup> Lis Stegmann,<sup>1</sup> Fabricio B. Baccaro,<sup>13</sup> Leandro Juen,<sup>12</sup> Juliana Schietti,<sup>13,14</sup> Luiz Aragão,<sup>15</sup> Erika Berenguer,<sup>4,16</sup> Leandro Castello,<sup>17</sup> Flavia R.C. Costa,<sup>14</sup> Matheus L. Guedes,<sup>18</sup> Cecilia G. Leal,<sup>4</sup> Alexander C. Lees,<sup>10</sup> Victoria Isaac,<sup>12</sup> Rodrigo O. Nascimento,<sup>12</sup> Oliver L. Phillips,<sup>18,19</sup> Fernando Augusto Schmidt,<sup>20</sup> Hans ter Steege,<sup>21,22</sup> Fernando Vaz-de-Mello,<sup>23</sup> Eduardo M. Venticinque,<sup>24</sup> Ima Célia Guimarães Vieira,<sup>25</sup> Jansen Zuanon,<sup>14</sup> The Synergize Consortium, and Joice Ferreira<sup>1,12</sup>

<sup>1</sup>Empresa Brasileira de Pesquisa Agropecuária, Amazônia Oriental, Belém 66095-903, Brazil

<sup>2</sup>Universidade de São Paulo, São Paulo 05508-220, Brazil

<sup>3</sup>Universidade de São Paulo, Esalq, Piracicaba 13418-900, Brazil

<sup>4</sup>Lancaster University, LA1 4YQ Lancaster, UK

<sup>5</sup>University of Bristol, BS8 1QU Bristol, UK

<sup>6</sup>Universidade Estadual de Campinas, Campinas 13083-862, Brazil

<sup>7</sup>Universidade Federal da Paraíba, Areia 58397-000, Brazil

<sup>8</sup>Universidade de Brasília, Brasília 70910-900, Brazil

<sup>9</sup>Universidade do Porto, 4099-002 Porto, Portugal

<sup>10</sup>Manchester Metropolitan University, M15 6BH Manchester, UK

<sup>11</sup>Universidade Federal de Lavras, Lavras 37200-000, Brazil

<sup>12</sup>Universidade Federal do Pará, Belém 66075-119, Brazil

<sup>13</sup>Universidade Federal do Amazonas, Manaus 69067-005, Brazil

<sup>14</sup>Instituto Nacional de Pesquisas da Amazônia, Manaus 69067-375, Brazil

<sup>15</sup>Instituto Nacional de Pesquisas Espaciais, São José dos Campos 12227-010, Brazil

<sup>16</sup>University of Oxford, OX1 3QY Oxford, UK

<sup>17</sup>Virginia Tech, Blacksburg, VA 24061, USA

<sup>18</sup>Universidade Federal Rural da Amazônia, Belém 66077-830, Brazil

<sup>19</sup>University of Leeds, LS2 9JT Leeds, UK

<sup>20</sup>Universidade Federal do Acre, Rio Branco 69920-900, Brazil

<sup>21</sup>Naturalis Biodiversity Center, 2333 CR Leiden, the Netherlands

<sup>22</sup>Utrecht University, 3584 CS Utrecht, the Netherlands

<sup>23</sup>Universidade Federal de Mato Grosso, Cuiabá 78060-900, Brazil

<sup>24</sup>Universidade Federal do Rio Grande do Norte, Natal 59078-970, Brazil

<sup>25</sup>Museu Paraense Emílio Goeldi, Belém 66040-170, Brazil

<sup>26</sup>These authors contributed equally

<sup>27</sup>Lead contact

\*Correspondence: [raqueluly@gmail.com](mailto:raqueluly@gmail.com) (R.L.C.), [gel.florestal@gmail.com](mailto:gel.florestal@gmail.com) (A.F.R.), [jos.barlow@lancaster.ac.uk](mailto:jos.barlow@lancaster.ac.uk) (J.B.), [mariormoura@gmail.com](mailto:mariormoura@gmail.com) (M.R.M.)

<https://doi.org/10.1016/j.cub.2023.06.077>

## SUMMARY

Biodiversity loss is one of the main challenges of our time,<sup>1,2</sup> and attempts to address it require a clear understanding of how ecological communities respond to environmental change across time and space.<sup>3,4</sup> While the increasing availability of global databases on ecological communities has advanced our knowledge of biodiversity sensitivity to environmental changes,<sup>5–7</sup> vast areas of the tropics remain understudied.<sup>8–11</sup> In the American tropics, Amazonia stands out as the world's most diverse rainforest and the primary source of Neotropical biodiversity,<sup>12</sup> but it remains among the least known forests in America and is often underrepresented in biodiversity databases.<sup>13–15</sup> To worsen this situation, human-induced modifications<sup>16,17</sup> may eliminate pieces of the Amazon's biodiversity puzzle before we can use them to understand how ecological communities are responding. To increase generalization and applicability of biodiversity knowledge,<sup>18,19</sup> it is thus crucial to reduce biases in ecological research, particularly in regions projected to face the most pronounced environmental changes. We integrate ecological community metadata of 7,694 sampling sites for multiple organism groups in a machine learning model framework to map the research probability across the Brazilian Amazonia, while identifying the region's vulnerability to environmental change. 15%–18% of the most neglected areas in ecological research are expected to experience severe climate or land use changes by 2050. This means that unless we take immediate action, we will not be able to establish their current status, much less monitor how it is changing and what is being lost.



## RESULTS

Our detailed assessment of the ecological research in the Brazilian Amazon assessed how logistics and human influence on the forests explained research probability across 7,694 community ecology sites surveyed from 2010 to 2020. Across nine organism groups—benthic invertebrates, heteropterans, odonates, fishes, macrophytes, birds, woody vegetation, ants, and dung beetles—ecological research was unevenly distributed in all three major habitat types investigated with easily identifiable research gaps (research probability < 0.1) covering 54.1% of unflooded areas locally known as terra firme (uplands, hereafter), 27.3% of aquatic habitats, and 17.3% of wetlands (Figure 1). While ecological research effort differs across organism groups, our findings highlight very consistent spatial patterns of research probability across habitat types, even among groups assessed in multiple habitat types (plants and birds; Figure 2).


### Drivers of research biases

Overall, logistics and human influence factors explained 64% of the variation in research probability. Among the logistic-related factors, accessibility and distance to research facilities

consistently emerged as important predictors of research probability (Figure 3), highlighting the role of logistical constraints and ease of access. Research probability increased with closer proximity to transportation and research facilities for all upland organisms and most representatives of wetlands and aquatic habitats. In addition, dry season length mattered for ecological research on wetland birds but showed little contribution with respect to other organisms (Figure 3C). Dry season length also had a contrasting effect across habitat types, increasing the research probability in uplands and aquatic habitats, but decreasing it in wetlands. Although logistics influenced ecological research the most, forest degradation and land tenure also showed a modest but consistent importance across all organism groups. Both predictors affected ecological research in the same direction across organisms, with research probability slightly declining in more degraded areas and indigenous lands but increasing in protected areas (Figures 3D and 3E).


### Research biases and projected environmental changes

Unfortunately, about half of the Brazilian Amazon is either already deforested (23.50%) or projected (27.29%) to be by 2050,<sup>20</sup> with these regions showing contrasting chances of



**Figure 1. Research probability across the Brazilian Amazon**

The central map represents the average research probability across all organism groups and habitat types. The inset maps at the bottom illustrate the research probability for different organisms in aquatic (bluish maps), wetland (greenish maps), and upland (orangish maps) habitats. In all maps, the black crosses indicate the sampling points for the period between 2010 and 2020. The donut shows the percentage of samples belonging to each biological group across different habitats (beetles and ants from wetlands were not modeled due to low sample size). Research probability was accurately predicted across all organism groups (mean Sorensen 0.89, range = 0.84–0.93). See Table S3.



**Figure 2. Research probability for different organisms in aquatic, wetland, and upland habitats**

ecological research (Figures 4E and S2). For instance, research probability was higher among areas currently deforested than in areas projected to be deforested within the next three decades (Figure 4E), corroborating the finding that research has occurred mostly in human-modified landscapes. Our findings also indicate that 15%–18% of the most scientifically neglected areas (Figure S2), herein defined as those in the first quartile of research probability, show high susceptibility to climate changes by 2050 (Figures 4A and 4C) and habitat destruction (Figures 4B and 4D).

## DISCUSSION

We elucidate how logistics and human influence have affected ecological research coverage in Brazilian Amazonia. Our comprehensive regional synthesis improves the understanding of Amazonian ecological research and opens new avenues to redress the underrepresentation of tropical rainforests in biodiversity databases. Besides offering further support for the role of accessibility and research facilities in ecological studies,<sup>21</sup> our findings also highlight some of the challenges involved in expanding research to areas that have not been sampled before.<sup>22</sup> While increasing overall research effort is valuable for many reasons, our models show that this would mostly lead to more surveys in areas with a high probability of research, as the lack of accessibility and research infrastructure<sup>15</sup> promotes close distances between new and existing sampling sites.<sup>19</sup> Additional measures will be required to overcome these barriers and reach regions where the research probability is low. Here we explore two key challenges related to resolving spatial research gaps in the Brazilian Amazon.


## Remote regions

Accessibility was a key factor in our results. One strategy to reach undersampled regions would be to fund expedition-like programs such as the *Expedição Serra da Mocidade*, where Amazon-based researchers from different institutions surveyed remote mountains in northern Amazonia, where no previous studies had been conducted.<sup>23</sup> While the *Expedição Serra da Mocidade* focussed mainly on finding new species and understanding species distributions, an ecological expedition would require longer periods in the field to enable the use of the standardized sampling protocols required to assess biotic changes across space and time. For instance, regions with low research probability partially overlap with those projected to experience either high and low climate change, as well as with regions facing high risk of future deforestation and degradation (Figure 4). Hence, enhancing ecological research in these remote regions could be the sole chance to unveil pieces of the Amazon's biodiversity puzzle before they succumb to human-induced modifications, while also seizing one of our prime opportunities to comprehend climate change effects without the potential anthropogenic influence.<sup>24</sup>

Distance to research centers was also an important factor, and an alternative approach would be to fund new centers in cities that are within or close to the areas with the lowest probability of research. While this could be a more logistically demanding approach, it has four major longer-term benefits. First, it would encourage the training of local researchers and help science to endure beyond individual assessments. Second, it could build the base for more detailed scientific research, including the logistical support necessary to identify and prepare specimens and develop local collections. Third, it could enhance the capacity required to conduct the longer-term research required to understand global change. Finally, it may be a more sustainable strategy than investing in expeditions, as these are rarely repeated and risk being canceled under new governments. Although research centers have suffered under recent governments, being subjected to long-lasting neglect affecting funding, structural maintenance, and emptying of key scientific and technical staff,<sup>25</sup> they have also shown themselves to be resilient, and continue to lead and support long-term research across the Amazon. Whatever approach is taken, it is key that it does not undermine the current network of research and education facilities in the Brazilian Amazonia or remove funding from long-term monitoring programs on biodiversity baselines and ecological changes,<sup>24</sup> which often take place near existing research facilities.

## Indigenous lands

Results highlighted the limited research effort in indigenous lands when compared to strictly protected and sustainable use reserves. This reflects a major ecological knowledge gap, as indigenous lands cover about 23% of the Brazilian Amazon and play a fundamental role in preserving Amazonia's biocultural diversity.<sup>26,27</sup> Over the last years, Brazil's indigenous lands have come under increasing threat from illegal activities, such as logging, invasion by squatters, and gold mining,<sup>28</sup> with this latter activity also strongly impacting the health of riverside peoples.<sup>29</sup> The lack of government support has forced traditional and




**Figure 3. Magnitude and direction of logistics and human influence effects on research probability**

Partial dependence plots illustrate the research probability across increasing values of (A) accessibility, (B) distance to research facilities, (C) dry season length, (D) degradation, and (E) categories of land tenure in the Brazilian Amazonia. Continuous predictors (A–D) were standardized between 0 and 1 to facilitate comparisons, whereas the land tenure predictor (E) is represented by mean values of each category. Circle size is proportional to the variable importance measured as the increase in mean squared error (MSE) after randomizing values of each predictor.

indigenous peoples to defend their territories on their own.<sup>30</sup> Indigenous lands such as the Kaiapó and Araribóia already represent some of the last areas of extensive forest in the south

and eastern Amazon, and the role of indigenous lands is expected to become more pronounced under business-as-usual deforestation practices (Figure 4E).



**Figure 4. Research probability in relation to anthropogenic disturbances**

(A and B) Maps illustrate the overlap between research probability and areas subject to future (A) climate change and (B) deforestation-degradation across the Brazilian Amazon. (B) also indicates areas currently deforested (gray pixels) or expected to face deforestation by 2050 (black pixels). Currently, deforested areas may be reforested in case of land abandonment or political incentives.

(C–E) Bottom plots show changes in research probability in relation to climate change (C), degradation (D), and deforestation (E). Each disturbance is standardized within the range of 0 to 1.

See also [Figure S2](#).

While bureaucratic requirements and limitations in local communicability may reduce the research propensity in such areas,<sup>19</sup> coordinated actions between the environment ministry (a specific organization within the Brazilian government) research centers and indigenous peoples have high potential to minimize knowledge gaps within indigenous lands. Any such knowledge co-production needs to be equitable and decolonial, and to recognize and respect the diverse knowledge systems in place.<sup>31,32</sup> For instance, traditional practices of indigenous people should be valued and incorporated into research methodologies, ensuring their active participation and ownership in the process. The co-creation of research with local communities will likely require additional support and training for both the scientific community and for local communities and their organizations.<sup>31</sup> Such a program could have many additional benefits, as enhancing the involvement of local communities could support more inclusive science as well as better resource management and livelihoods.<sup>33</sup>

#### Differences between organisms and habitats

We find remarkably consistent drivers of research effort across different organisms and habitats. But one noticeable exception is the disproportionate importance that a small number of local experts have in the spatial sampling of underrepresented taxa.

For instance, most sampling (95%) of aquatic invertebrates (heteropterans and odonates) are distributed in eastern Amazonia (Pará state) and come from a single research group composed of local experts based in the Amazonian city of Belém. Other aquatic invertebrate specialists have collected in different regions of Amazonia, but their focus on taxonomy means they rarely use the standardized sampling required for ecological data. Considering the distinct objectives of taxonomic research and ecological sampling, prior planning and greater collaboration will be required to achieve the potential benefits that could be accrued from integrating Amazonian ecology and taxonomy. We also demonstrate that research gaps are higher for uplands than wetlands and aquatic habitats, which likely reflects the role that the broad network of navigable waterways has in facilitating access to wetland and aquatic areas.

#### Can regional curation help reduce global biases?

The metadata we based this research on describe a large number of datasets with substantial coverage across multiple organisms in the Brazilian Amazon (see the Synergize project<sup>34</sup>). To date, few of these datasets are integrated into global databases. For example, the BioTIME,<sup>35</sup> BIOFRAG,<sup>6</sup> FragSAD,<sup>7</sup> and Predicts<sup>5</sup> databases collectively include only 222 datasets for the

Brazilian Amazon, representing less than 3% of the Synergize effort. In contrast, more than 40% of Synergize datasets ( $n = 3,281$ ) can potentially contribute to global assessments, adding 1,103 time series datasets for BioTIME, 506 datasets under the requirements of BIOFRAG/FragSAD, and 1,672 datasets on land use comparisons that meet the objectives of Predicts. Although these are upper bound estimates that will fall due to the additional requirements of specific global networks (e.g., pre-defined taxa and habitats), the differences highlight the value of carefully produced regional datasets to mitigate data biases in collaborative networks.<sup>36</sup> To secure the engagement of tropical research communities, it is crucial to implement an inclusive code of conduct that values data ownership in resulting products.<sup>36–38</sup>

### Conclusion

Our large-scale assessment of ecological research across the Brazilian Amazon not only highlights the extent of research gaps and biases in tropical systems, but also provides valuable insights into potential solutions to improve conservation planning for the world's most diverse rainforest. We show the importance of going beyond areas that are accessible and close to research bases, and expanding research into regions that will likely be affected by climate change or deforestation. Doing so will not be easy, and ecology alone will not help resolve the environmental crises facing the world. But understanding the responses of biodiversity and ecosystems forms a key part of keeping society informed about its impacts and supporting the implementation of evidence-based policies and practices that can help mitigate the worst outcomes.

### STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **RESOURCE AVAILABILITY**
  - Lead contact
  - Materials availability
  - Data and code availability
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
  - Data collection
- **METHOD DETAILS**
  - Predictors of research probability
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
  - Research probability modelling
  - Overlap between research and anthropogenic disturbances
- **ADDITIONAL RESOURCES**

### SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.cub.2023.06.077>.

### CONSORTIA

The members of The Synergize Consortium are Raquel L. Carvalho, Angelica F. Resende, Jos Barlow, Filipe França, Mario R. Moura, Rafaella Maciel,

Fernanda Alves-Martins, Jack Shutt, Cassio A. Nunes, Fernando Elias, Juliana M. Silveira, Lis Stegmann, Fabricio B. Baccaro, Leandro Juen, Juliana Schietti, Luiz Aragão, Erika Berenguer, Leandro Castello, Flavia R.C. Costa, Matheus L. Guedes, Cecilia G. Leal, Alexander C. Lees, Victoria Isaac, Rodrigo O. Nascimento, Oliver L. Phillips, Fernando Augusto Schmidt, Hans ter Steege, Fernando Vaz-de-Mello, Eduardo M. Venticinque, Ima Célia Guimarães Vieira, Jansen Zuanon, Joice Ferreira, Adem Nagibe dos Santos Geber Filho, Ademir Ruschel, Adolfo Ricardo Calor, Adriana de Lima Alves, Adriane Esquivel Muelbert, Adriano Quaresma, Alberto Vicentini, Alexandra Rocha da Piedade, Alexandre Adalardo de Oliveira, Alexandre Aleixo, Alexandre Casadei-Ferreira, Alexandre Gontijo, Alexandre Hercos, Aline Andriolo, Aline Lopes, Aline Pontes-Lopes, Allan Paulo Moreira dos Santos, Amanda Batista da Silva de Oliveira, Amanda Frederico Mortati, Ana Karina Moreyra Salcedo, Ana Luisa Albernaz, Ana Luisa Fares, Ana Luiza-Andrade, Ana Maria Oliveira Pes, Ana Paula Justino Faria, Anderson Pedro Bernadina Batista, Anderson Puker, Anderson S. Bueno, André Braga Junqueira, André Luiz Ramos Holanda de Andrade, André Ricardo Ghidini, André V. Galuch, Addressa Silvana Oliveira de Menezes, Angelo Gilberto Manzatto, Anne Sthephane A.S. Correa, Antonio C.M. Queiroz, Antonio Carlos da Silva Zanzini, Antonio Miguel Olivo Neto, Antonio Willian Flores de Melo, Aretha Franklin Guimaraes, Arilson Bezerra Castro, Augusto Borges, Aurélia Bentes Ferreira, Beatriz S. Marimon, Ben Hur Marimon-Junior, Bernardo M. Flores, Bethânia Oliveira de Resende, Bianca Weiss Albuquerque, Boris Villa, Bradley Davis, Bruce Nelson, Bruce Williamson, Bruna Santos Bitencourt de Melo, Bruno B.L. Cintra, Bruno Borges Santos, Bruno da Silveira Prudente, Bruno Garcia Luize, Bruno Spacek Godoy, Cameron L. Rutt, Camila Duarte Ritter, Camila V.J. Silva, Carla Rodrigues Ribas, Carlos A. Peres, Carlos Augusto Silva de Azevêdo, Carlos Freitas, Carlos Leandro Cordeiro, Carlos Rodrigo Brocardo, Carolina Castilho, Carolina Levis, Carolina Rodrigues da Costa Doria, Caroline C. Arantes, Cássia Anicá dos Santos, Catarina C. Jakovac, Celice Alexandre Silva, Cesar João Benetti, Chaim Lasmar, Charles J. Marsh, Christian Borges Andretti, Cinthia Pereira de Oliveira, Cintia Cornelius, Clarissa Alves da Rosa, Cláudia Baider, Cláudia G. Gualberto, Claudia Pereira de Deus, Cláudio da Silva Monteiro Jr., Cláudio Rabelo dos Santos Neto, Cleonice Maria Cardoso Lobato, Cleverson Rannieri Meira dos Santos, Cristian Camilo Mendoza Penagos, Daniel da Silva Costa, Daniel Luis Mascia Vieira, Daniel Praia Portela de Aguiar, Daniel Silas Veras, Daniela Pauletto, Danielle de Lima Braga, Danielle Storck-Tonon, Daniely da Frota Almeida, Danyhelton Douglas, Dário Dantas do Amaral, Darlene Gris, David Luther, David P. Edwards, David Pedroza Guimarães, Deane Cabral dos Santos, Débora Rodrigues de Souza Campana, Denis Silva Nogueira, Dennis Rodrigues da Silva, Dhâmyla Bruna de Souza Dutra, Dian Carlos Pinheiro Rosa, Diego Armando Silva da Silva, Diego Pedroza, Diego V. Anjos, Diego Viana Melo Lima, Divino V. Silvério, Domingos de Jesus Rodrigues, Douglas Bastos, Douglas Daly, Edelcilio Marques Barbosa, Edith Rosario Clemente Arenas, Edmar Almeida de Oliveira, Elnaira Alencar dos Santos, Edrielly Caroline Carvalho de Santana, Edson Guilherme, Edson Vidal, Eduardo Malta Campos-Filho, Eduardo van den Berg, Elder Ferreira Morato, Eliodomar R. da Silva, Elineide E. Marques, Elizabeth G. Pringle, Elizabeth Nichols, Ellen Andresen, Emanuelle de Sousa Farias, Emely Laiara Silva de Siqueira, Emília Zoppas de Albuquerque, Eric Bastos Görgens, Erlane José Rodrigues da Cunha, Ethan Householder, Evlyn Márcia Moraes de Leão Novo, Fabiana Ferreira de Oliveira, Fabio de Oliveira Roque, Fabricio Coletti, Fagno Reis, Felipe F.F. Moreira, Felipe Todeschini, Fernanda Antunes Carvalho, Fernanda Coelho de Souza, Fernando Augusto Barbosa Silva, Fernando Geraldo Carvalho, Fernando Gonçalves Cabeceira, Fernando Mendonça d'Horta, Fernando P. Mendonça, Fernando Prado Florêncio, Fernando Rogério de Carvalho, Filipe Viegas de Arruda, Flávia Alessandra da Silva Nonato, Flávia Delgado Santana, Flavia Durgante, Flávia Kelly Siqueira de Souza, Flávio Amorim Obermuller, Flávio Siqueira de Castro, Florian Wittmann, Francisco Matheus da Silva Sales, Francisco Valente-Neto, Frederico Falcão Salles, Gabriel Costa Borba, Gabriel Damasco, Gabriel Gazzana Barros, Gabriel Lourenço Bregião, Gabriela Abrantes Jardim, Ghilleen T. Prance, Gisiane Rodrigues Lima, Gleison Robson Desidério, Gracilene da Costa de Melo, Guilherme Henrique Pompiano do Carmo, Guilherme Sampaio Cabral, Guillaume Xavier Rousseau, Gustavo Cardoso da Silva, Gustavo Schwartz, Hannah Griffiths, Helder Lima de Queiroz, Helder M.V. Espírito-Santo, Helena Soares Ramos Cabette, Henrique Eduardo Mendonça Nascimento, Heraldo L. Vasconcelos, Herison Medeiros, Hilton Jeferson Alves Cardoso de Aguiar, Hingara Leão, Icaro Wilker,

Inês Correa Gonçalves, Inocência de Sousa Gorayeb, Ires Paula de Andrade Miranda, Irving Foster Brown, Isis Caroline Siqueira Santos, Itanna Oliveira Fernandes, Izaias Fernandes, Jacques Hubert Charles Delabie, Jadson Coelho de Abreu, Jaime de Liege Gama Neto, Janaina Barbosa Pedrosa Costa, Janaína Costa Noronha, Janaina Gomes de Brito, Jared Wolfe, Jean Carlos Santos, Jefferson Ferreira-Ferreira, Jerriane Oliveira Gomes, Jesse R. Lasky, Jéssica Caroline de Faria Falcão, Jessica Gomes Costa, Jessica Soares Cravo, Jesús Enrique Burgos Guerrero, Jhonatan Andrés Muñoz Gutiérrez, João Carreiras, João Lanna, Joás Silva Brito, Jochen Schöngart, Jonas José Mendes Aguiar, Jônatas Lima, Jorcely G. Barroso, Jorge Ari Noriega, Jorge Luiz da Silva Pereira, Jorge Luiz Nessimian, Jorge Luiz Pereira de Souza, José Julio de Toledo, José Leonardo Lima Magalhães, José Luís Camargo, José Max B. Oliveira Jr., José Moacir Ferreira Ribeiro, José Orlando de Almeida Silva, José Renan da Silva Guimarães, Joseph E. Hawes, Joudellys Andrade-Silva, Juan David Cardenas Revilla, Júlia Santana da Silva, Juliana da Silva Menger, Juliana Rechetelo, Juliana Stropp, Julianna Freires Barbosa, Julio Daniel do Vale, Julio Louzada, Július César Cerqueira Silva, Karina Dias da Silva, Karina Melgaço, Karine Santana Carvalho, Kedma Cristine Yamamoto, Keila Rêgo Mendes, Kevina Vulínec, Laís Ferreira Maia, Larissa Cavalheiro, Laura Barbosa Vedovato, Layon Oreste Demarchi, Leandro Giacomini, Leandro Lourenço Dumas, Leandro Maracahipes, Leandro Schlemmer Brasil, Leandro Valle Ferreira, Lenize Batista Calvão, Leonardo Maracahipes-Santos, Leonardo Pequeno Reis, Letícia Fernandes da Silva, Lia de Oliveira Melo, Lidiany Camila da Silva Carvalho, Lílian Casatti, Lílian Lund Amado, Líliane Steidle de Matos, Lisandro Vieira, Livia Pires do Prado, Luana Alencar, Luane Fontenele, Lucas Mazzei, Lucas Navarro Paolucci, Lucas Pereira Zanzini, Lucélia Nobre Carvalho, Luciana Carvalho Crema, Luciane Ferreira Barbosa Brulinger, Luciano Fogaça de Assis Montag, Luciano Nicolas Naka, Ludson Azara, Luis Fábio Silveira, Luis Gabriel de Oliveira Nunes, Luís Miguel do Carmo Rosalino, Luiz A.M. Mestre, Luiz Carlos de Matos Bonates, Luiz de Souza Coelho, Luiz Henrique Medeiros Borges, Luzia da Silva Lourenço, Madson Antonio Benjamin Freitas, Maiara Tábatha da Silva Brito, Maihyra Marina Pombo, Maíra da Rocha, Maira Rodrigues Cardoso, Marcelino Carneiro Guedes, Marcelo Bassols Raseira, Marcelo Brilhante de Medeiros, Marcelo de Jesus Veiga Carim, Marcelo Fragomeni Simon, Marcelo Petratti Pansonato, Marcelo Rodrigues dos Anjos, Marcelo Trindade Nascimento, Márcia Regina de Souza, Marcília Gabriella Tavares Monteiro, Márcio Joaquim da Silva, Marcio Uehara-Prado, Marco Antonio de Oliveira, Marcos Callisto, Marcos José Salgado Vital, Marcos Pêrsio Dantas Santos, Marcos Silveira, Marcus Vinício Neves D. Oliveira, Maria Angélica Pérez-Mayorga, Maria Antonia Carniello, Maria Aparecida Lopes, Maria Aurea Pinheiro de Almeida Silveira, Maria Cristina Espósito, Maria Eduarda Maldaner, Maria Inês S. Passos, Maria José Pinheiro Anacleto, Maria Katiane Sousa Costa, Maria Pires Martins, Maria Teresa Fernandez Piedade, Mariana Victória Irumé, Marília Maria Silva da Costa, Marina Franco de Almeida Maximiano, Marina Guimarães Freitas, Mark A. Cochrane, Markus Gastauer, Marllus Rafael Negreiros Almeida, Mateus Fernando de Souza, Michel Catarino, Michela Costa Batista, Mike R. Massam, Mila Ferraz de Oliveira Martins, Milena Holmgren, Morgana Almeida, Murilo S. Dias, Nádia Barbosa Espírito Santo, Naraiana Loureiro Benone, Natalia Macedo Ivanauskas, Natália Medeiros, Natalia Targhetta, Nathalia Silva Félix, Nelson Ferreira Jr., Neusa Hamada, Nubia Campos, Nubia França da Silva Giehl, Oliver Charles Metcalf, Otávio Guilherme Morais da Silva, Pablo Vieira Cerqueira, Pamela Moser, Patrícia Nakayama Miranda, Patricia Santos Ferreira Perucetti, Paula Palhares de Polari Alverga, Paula Prist, Paula Souto, Paulo Brando, Paulo dos Santos Pompeu, Paulo Eduardo Barni, Paulo Maurício de Alencastro Graça, Paulo S. Morandi, Paulo Vilela Cruz, Pedro Giovâni da Silva, Pitágoras C. Bispo, Plínio Barbosa de Camargo, Priscila S. de M. Sarmiento, Priscila Souza, Rafael Barreto de Andrade, Rafael Benzi Braga, Rafael Boldrini, Rafael Costa Bastos, Rafael Leandro de Assis, Rafael P. Salomão, Rafael Pereira Leitão, Raimundo N.G. Mendes, Jr., Rainiellen de Sá Carpanedo, Ramiro Dário Melinski, Raphael Ligeiro, Raúl Enrique Pirela Pérez, Reinaldo Imbrozio Barbosa, Reinaldo Lucas Cajaiba, Renato Azevedo Matias Silvano, Renato Portela Salomão, Renato Richard Hilário, Renato Tavares Martins, Ricardo de Oliveira Perdiz, Ricardo Eduardo Vicente, Ricardo José da Silva, Ricardo Koroiva, Ricardo Solar, Richarly da Costa Silva, Robson Borges de Lima, Robson dos Santos Alves da Silva, Rodolfo Mariano, Rodrigo Arison Barbosa Ribeiro, Rodrigo Ferreira Fadini, Rodrigo Leonardo Costa de Oliveira, Rodrigo Machado Feitosa, Rodrigo Matavelli, Roger Paulo Mormul,

Rogério Rosa da Silva, Ronald Zanetti, Ronaldo Barthem, Rony Peterson Santos Almeida, Sabina Cerruto Ribeiro, Salustiano Vilar da Costa Neto, Samuel Nienow, Sérgio Augusto Vidal de Oliveira, Sérgio Henrique Borges, Sérgio Milheiras, Sérvio Pontes Ribeiro, Sheyla Regina Marques Couceiro, Sidney Araújo de Sousa, Sílvia Barbosa Rodrigues, Sílvia Leitão Dutra, Simon Mahood, Simone Aparecida Vieira, Solange Arrolho, Sonaira Souza da Silva, Stefania Pinzón Triana, Susan Laurance, Sustanis Horn Kunz, Swanni T. Alvarado, Tais Helena Araujo Rodrigues, Talitha Ferreira dos Santos, Tatiana Lemos da Silva Machado, Ted R. Feldpausch, Thaiane Sousa, Thaisa Sala Michelan, Thaise Emilio, Thaline de Freitas Brito, Thiago André, Thiago Augusto Pedroso Barbosa, Thiago Barros Miguel, Thiago Junqueira Izzo, Thiago Orsi Laranjeiras, Thiago Pereira Mendes, Thiago Sanna Freire Silva, Tiago Kütter Krolow, Tiago Octavio Begot, Timothy R. Baker, Tomas F. Domingues, Tommaso Giarrizzo, Tony Vizcarra Bentos, Torbjorn Haugaasen, Ualerson Peixoto, Uly Matilde Pozzobom, Vanessa Korasaki, Vanessa Soares Ribeiro, Veridiana Vizoni Scudeller, Victor Hugo Fonseca Oliveira, Victor Lemes Landeiro, Victor Renan Santos Ferreira, Victória de Nazaré Gama Silva, Vitor Hugo Freitas Gomes, Vivian Campos de Oliveira, Viviane Firmino, Wagner Tadeu Vieira Santiago, Wallace Beiroz, Wanessa Rejane de Almeida, Washington Luis de Oliveira, Wegiane Campelo da Silva, Wendeson Castro, Wesley Dáttilo, Wesley Jonatar Alves da Cruz, Wheriton Fernando Moreira da Silva, William E. Magnusson, William Laurance, William Milliken, William Sousa de Paula, Yadvinder Malhi, Yosio Edemir Shimabukuro, Ysadhora Gomes de Lima, Yulie Shimano, and Yuri Feitosa. A full list of affiliations for The Synergize Consortium members can be found in [Table S5](#).

#### ACKNOWLEDGMENTS

The Synergize Network was developed through funding from the Brazilian Research Council (CNPq) under the Brazilian Synthesis Centre on Biodiversity and Ecosystem Services (Sinbiose) (grant CNPq/MCTIC 442354/2019-3). We also recognize support from the Conselho Nacional de Desenvolvimento Científico (CNPq) for research grants in support of R.L.C. (151221/2021-9) and A.F.R. (150196/2020-2) and São Paulo Research Foundation (FAPESP) for grants in support of R.L.C. (2022/07381-9), A.F.R. (2019/24049-5), and M.R.M. (2021/11840-6 and 2022/12231-6). J.B. thanks the Natural Environment Research Council ARBOLES project NE/S011811/1. F.M.F. thanks the University of Bristol (PolicyBristol, 1989427; Climate and Net Zero Impact Awards, 170839; and Elizabeth Blackwell Institute Rapid Research Funding, 2208557). E.B. thanks the Natural Environment Research Council (NE/S01084X/1) for supporting her work and part of that of the data curation team. F.A.-M. acknowledges funding from the European Union's Horizon 2020 research and innovation programme under grant agreement 854248. [ForestPlots.net](#) communication and data management support in Brazil since 2020 was made possible by ERC Starter Grant 758873 TreeMort and the Norwegian Research Council grant "Cascading consequences of hunting & fishing for ecosystem services in Amazonian Forest," which supported O.L.P.'s contribution. Work to establish and sustain RAINFOR plots in Brazil has been supported by the European Research Council (advanced grant 291585 – T-FORCES), the Gordon and Betty Moore Foundation (1656 RAINFOR), the European Union's Fifth, Sixth, and Seventh Framework Programme (EVK2-CT-1999-00023 – CARBONSINK-LBA, 283080 – GEOCARBON, and 282664 – AMAZALERT), the Natural Environment Research Council (NE/D005590/1 – TROBIT, NE/F005806/1 – AMAZONICA, and PPFOR E/M0022021/1), and the NERC/State of São Paulo Research Foundation (FAPESP) consortium grants BIO-RED (NE/N012542/1 and 2012/51872-5), ECOFOR (NE/K016431/1 and 2012/51509-8), ARBOLES (NE/S011811/1), and the Royal Society (Global Challenges Award FORAMA, ICA/R1/180100). Synergize Consortium members thank the following funders: CAPES-PROCAD 88887.200472/2018-00, FAPESPA ICAAF 070/2020, and FAPESP 2019/25478-7 (BEX5528/13-5, 383744/2015-6, and 403811/2012-0). We acknowledge all the researchers and field assistants who helped with the data collection but are not within the authors. They are Ana Segalin de Andrade, Arlete Almeida, Atila Alves de Oliveira, Carlos Alberto Quesada, Carolina Toledo Garib Soriano, Charles Eugene Zartman, Cid Ferreira, Cintia Rodrigues De Souza, Claudinei Oliveira-Santos, Daniel Nepstad, David J. Mellow, Diego de Alcântara Pires, Dilce de Fátima Rossetti, Diogenes de Andrade Lima Filho, Eduardo Hase, Everton José Almeida, Flávia Rodrigues Barbosa,



Francisca Dionizia de Almeida Matos, Francisco de Assis Esteves, Gabriel de Oliveira de Almeida, Gustavo Paiva Evangelista da Rocha, Haron Xaud, Iêda Leão do Amaral, Ileana A Acosta, Jennifer Balch, João Batista da Silva, Jocielei de Oliveira, Jorge Oliviera, José Ferreira Ramos, Juan Carlos Montero, Kallene Kharla Gonçalves da Silva, Kelly Liane da Silva Sampaio, Kelsy Ellen Vale da Silva, Lola da Costa, Lorena Rincón, Luana Fidelis da Silva, Lucas Lima da Silva, Luis Gustavo Canesi Ferreira, Márcia Cléia Vilela dos Santos, Márcia Patrícia Nascimento Cidade, Márcio Hofmann, Margarita Maria Jaramillo, Maria Carmozina Araújo, Maria do Socorro, Mario Cohn-Haft, Mateus Junior Garcia de Oliveira, Michael Coe, Natalino Silva, Nelson Silva Pinto, Pedro Lisboa, Perseu da Silva Aparício, Priscila Madoka Ito, Raimunda Oliveira de Araújo, Ricardo Barbieri, Richard Bardgett, Richard Lucas, Rita Mesquita, Rodrigo Gravina Prates Junqueira, Rogerio Gribel, Terezinha Maria de Castro Della Lucia, Vilany Matilla Colares Carneiro, William Balee, and Wully Barreto da Silva.

#### AUTHOR CONTRIBUTIONS

R.L.C. wrote the first draft; A.F.R., M.R.M., and J.B. edited the manuscript; J.B. supervised the work; R.L.C., F.M.F., F.A.-M., and L.S. performed the literature review; R.L.C., A.F.R., C.A.N., F.M.F., J.M. S., R.M., F.A.-M., and L.S. compiled the data; R.L.C., A.F.R., M.R.M., and J.B. designed the methodology; R.L.C., A.F.R., and M.R.M. performed the analysis and data visualization; R.L.C., A.F.R., J.B., F.M.F., M.R.M., R.M., F.A.-M., J. Shutt, C.A.N., F.E., J.M.S., L.S., F.B.B., L.J., J. Schietti, L.A., E.B., L.C., F.R.C.C., M.L.G., C.G.L., A.C.L., V.I., R.O.N., O.L.P., F.A.S., H.t.S., F.V.-d.-M., E.M.V., I.C.G.V., J.Z., and J.F. conceived the ideas; and J.F., F.M.F., F.V.-d.-M., A.C.L., C.G.L., E.B., J.B., L.C., F.R.C.C., I.C.G.V., O.L.P., L.A., J.Z., H.t.S., V.I., L.J., E.M.V., F.B.B., and F.A.S. obtained project funding. All authors contributed in the form of discussions, revisions, and suggestions and approved the final manuscript.

#### DECLARATION OF INTERESTS

The authors declare no competing interests.

#### INCLUSION AND DIVERSITY

We support inclusive, diverse, and equitable conduct of research.

Received: January 13, 2023

Revised: May 19, 2023

Accepted: June 28, 2023

Published: July 19, 2023; corrected online: July 25, 2023

#### REFERENCES

- Dirzo, R., and Raven, P.H. (2003). Global state of biodiversity and loss. *Annu. Rev. Environ. Resour.* 28, 137–167. <https://doi.org/10.1146/annurev.energy.28.050302.105532>.
- Barlow, J., França, F., Gardner, T.A., Hicks, C.C., Lennox, G.D., Berenguer, E., Castello, L., Economo, E.P., Ferreira, J., Guénard, B., et al. (2018). The future of hyperdiverse tropical ecosystems. *Nature* 559, 517–526. <https://doi.org/10.1038/s41586-018-0301-1>.
- Mokany, K., and Ferrier, S. (2011). Predicting impacts of climate change on biodiversity: A role for semi-mechanistic community-level modelling. *Divers. Distrib.* 17, 374–380. <https://doi.org/10.1111/j.1472-4642.2010.00735.x>.
- Ribeiro, G.V.T., Teixeira, A.L., Barbosa, N.P.U., and Silveira, F.A.O. (2016). Assessing bias and knowledge gaps on seed ecology research: implications for conservation agenda and policy. *Ecol. Appl.* 26, 2033–2043. <https://doi.org/10.1890/15-1852.1>.
- Newbold, T., Hudson, L.N., Hill, S.L.L., Contu, S., Lysenko, I., Senior, R.A., Börger, L., Bennett, D.J., Choimes, A., Collen, B., et al. (2015). Global effects of land use on local terrestrial biodiversity. *Nature* 520, 45–50. <https://doi.org/10.1038/nature14324>.
- Pfeifer, M., Lefebvre, V., Peres, C.A., Banks-Leite, C., Wearn, O.R., Marsh, C.J., Butchart, S.H.M., Arroyo-Rodríguez, V., Barlow, J., Cerezo, A., et al. (2017). Creation of forest edges has a global impact on forest vertebrates. *Nature* 551, 187–191. <https://doi.org/10.1038/nature24457>.
- Chase, J.M., McGill, B.J., Thompson, P.L., Antão, L.H., Bates, A.E., Blowes, S.A., Dornelas, M., Gonzalez, A., Magurran, A.E., Supp, S.R., et al. (2019). Species richness change across spatial scales. *Oikos* 128, 1079–1091. <https://doi.org/10.1111/oik.05968>.
- Mora, C., Tittensor, D.P., and Myers, R.A. (2008). The completeness of taxonomic inventories for describing the global diversity and distribution of marine fishes. *Proc. Biol. Sci.* 275, 149–155. <https://doi.org/10.1098/rspb.2007.1315>.
- Sano, E.E., Rosa, R., Luís, J., Laerte, S.B., and Ferreira, G. (2007). Documentos 190 Mapeamento de Cobertura Vegetal do Bioma Cerrado: Estratégias e Resultados Embrapa Cerrados Planaltina, DF 2007 (Empresa Brasileira de Pesquisa Agropecuária Embrapa Cerrados Ministério da Agricultura, Pecuária e Abastecimento).
- Ter Steege, H., Pitman, N.C.A., Killeen, T.J., Laurance, W.F., Peres, C.A., Guevara, J.E., Salomão, R.P., Castilho, C.v., Amaral, I.L., de Almeida Matos, F.D., et al. (2015). Estimating the global conservation status of more than 15,000 Amazonian tree species. *Sci. Adv.* 1, e1500936. <https://doi.org/10.1126/sciadv.1500936>.
- Moura, M.R., and Jetz, W. (2021). Shortfalls and opportunities in terrestrial vertebrate species discovery. *Nat. Ecol. Evol.* 5, 631–639. <https://doi.org/10.1038/s41559-021-014115>.
- Antonelli, A., Zizka, A., Carvalho, F.A., Scharn, R., Bacon, C.D., Silvestro, D., and Condamine, F.L. (2018). Amazonia is the primary source of Neotropical biodiversity. *Proc. Natl. Acad. Sci. USA* 115, 6034–6039. <https://doi.org/10.1073/pnas.1713819115>.
- Oliveira, U., Paglia, A.P., Brescovit, A.D., Carvalho, C.J.B., Silva, D.P., Rezende, D.T., Leite, F.S.F., Batista, J.A.N., Barbosa, J.P.P.P., Stehmann, J.R., et al. (2016). The strong influence of collection bias on biodiversity knowledge shortfalls of Brazilian terrestrial biodiversity. *Divers. Distrib.* 22, 1232–1244. <https://doi.org/10.1111/ddi.12489>.
- Stropp, J., Umbelino, B., Correia, R.A., Campos-Silva, J.V., Ladle, R.J., and Malhado, A.C.M. (2020). The ghosts of forests past and future: deforestation and botanical sampling in the Brazilian Amazon. *Ecography* 43, 979–989. <https://doi.org/10.1111/ecog.05026>.
- Moura, M.R., Costa, H.C., Peixoto, M.A., Carvalho, A.L., Santana, D.J., and Vasconcelos, H.L. (2018). Geographical and socioeconomic determinants of species discovery trends in a biodiversity hotspot. *Biol. Conserv.* 220, 237–244. <https://doi.org/10.1016/j.biocon.2018.01.024>.
- Pendrill, F., Gardner, T.A., Meyfroidt, P., Persson, U.M., Adams, J., Azevedo, T., Bastos Lima, M.G., Baumann, M., Curtis, P.G., De Sy, V., Garrett, R., et al. (2022). Disentangling the numbers behind agriculture-driven tropical deforestation. *Science* 377, eabm9267.
- Withey, K., Berenguer, E., Palmeira, A.F., Espírito-Santo, F.D.B., Lennox, G.D., Silva, C.V.J., Aragão, L.E.O.C., Ferreira, J., França, F., Malhi, Y., et al. (2018). Quantifying immediate carbon emissions from El Niño-mediated wildfires in humid tropical forests. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 373, 20170312.
- Martin, L.J., Blossey, B., and Ellis, E. (2012). Mapping where ecologists work: biases in the global distribution of terrestrial ecological observations. *Front. Ecol. Environ.* 10, 195–201. <https://doi.org/10.1890/110154>.
- dos Santos, J.G., Malhado, A.C.M., Ladle, R.J., Correia, R.A., and Costa, M.H. (2015). Geographic trends and information deficits in Amazonian conservation research. *Biodivers. Conserv.* 24, 2853–2863. <https://doi.org/10.1007/s10531-015-0981-x>.
- Lapola, D.M., Pinho, P., Barlow, J., Aragão, L.E.O.C., Berenguer, E., Carmentra, R., Liddy, H.M., Seixas, H., Silva, C.V.J., Silva-Junior, C.H.L., et al. (2023). The drivers and impacts of Amazon forest degradation. *Science* 379, eabp8622.
- Magnusson, W., Braga-Neto, R., Pezzini, F., Baccaro, F., Bergallo, H., Penha, J., Rodrigues, D., Verdade, L.M., Albertina, L., Albernaz, A.L., et al. (2014). Biodiversidade e Monitoramento Ambiental Integrado Biodiversidade and Integrated Environmental Monitoring (Instituto Nacional de Pesquisas da Amazônia).

22. Nóbrega, C.C., and de Marco, P. (2011). Unprotecting the rare species: a niche-based gap analysis for odonates in a core Cerrado area. *Divers. Distrib.* *17*, 491–505. <https://doi.org/10.1111/j.1472-4642.2011.00749.x>.
23. Laranjeiras, T.O., Melinski, R.D., Naka, L.N., Leite, G.A., Lima, G.R., d’Affonseca-Neto, J.A., and Cohn-Haft, M. (2019). Three bird species new to Brazil from the Serra da Mocidade, a remote mountain in Roraima. *Rev. Bras. Ornitol.* *27*, 275–283. <https://doi.org/10.1007/BF03546073>.
24. Berenguer, E., Lennox, G.D., Ferreira, J., Malhi, Y., Aragão, L.E.O.C., Barreto, J.R., Del Bon Espírito-Santo, F., Figueiredo, A.E.S., França, F., Gardner, T.A., et al. (2021). Tracking the impacts of El Niño drought and fire in human-modified Amazonian forests. *Proc. Natl. Acad. Sci. USA* *118*, e2019377118, <https://doi.org/10.1073/pnas.2019377118>.
25. Fernandes, G.W., Vale, M.M., Overbeck, G.E., Bustamante, M.M., Grelle, C.E., Bergallo, H.G., Magnusson, W.E., Akama, A., Alves, S.S., Amorim, A., et al. (2017). Dismantling Brazil’s science threatens global biodiversity heritage. *Perspect. Ecol. Conserv.* *15*, 239–243. <https://doi.org/10.1016/j.pecon.2017.07.004>.
26. Barretto Filho, H.T., Ramos, A., Sobral Barra, C., Barroso, M., Caron, P., Grupioni, B., Hildebrand, M.V., Jarretti, C., Junior, D.P., Moutinho, P., et al. (2021). Strengthening land and natural resource governance and management: protected areas, Indigenous lands, and local communities’ territories. In *The Amazon We Want*. <https://www.theamazonwewant.org/spa-reports/>.
27. Josse, C., Futaba, S. de M., von Hildebrand, M., de los Ríos, M.M., Oliveira-Miranda, M.A., Tenório, E.A.M., and Tuesta, E. (2021). The state of conservation policies, protected areas, and Indigenous territories, from the past to the present. In *The Amazon We Want*. <https://www.theamazonwewant.org/spa-reports/>.
28. Begotti, R.A., and Peres, C.A. (2019). Brazil’s indigenous lands under threat. *Science* *363*, 592. <https://doi.org/10.1126/science.aaw3864>.
29. Pestana, I.A., de Rezende, C.E., Almeida, R., de Lacerda, L.D., and Bastos, W.R. (2022). Let’s talk about mercury contamination in the Amazon (again): the case of the floating gold miners’ village on the Madeira River. *Extr. Ind. Soc.* *11*, 101122, <https://doi.org/10.1016/j.exis.2022.101122>.
30. Peres, C.A., Campos-Silva, J., and Ritter, C.D. (2023). Environmental policy at a critical junction in the Brazilian Amazon. *Trends Ecol. Evol.* *38*, 113–116. <https://doi.org/10.1016/j.tree.2022.11.011>.
31. Malmer, P., Vanessa, M., Austin, B., and Tengö, M. (2020). Mobilisation of indigenous and local knowledge as a source of useable evidence for conservation partnerships. In *Conservation Research, Policy and Practice* (Cambridge University Press), pp. 82–113. <https://doi.org/10.1017/9781108638210.006>.
32. Guidelines for Ethical Research in Indigenous Studies (2000). Guidelines for Ethical Research in Indigenous Studies (The Australian Institute of Aboriginal and Torres Strait Islander Studies). [https://www.wipo.int/export/sites/www/tk/en/databases/creative\\_heritage/docs/aiatsis\\_ethical\\_research.pdf](https://www.wipo.int/export/sites/www/tk/en/databases/creative_heritage/docs/aiatsis_ethical_research.pdf).
33. Campos-Silva, J.V., and Peres, C.A. (2016). Community-based management induces rapid recovery of a high-value tropical freshwater fishery. *Sci. Rep.* *6*, 34745, <https://doi.org/10.1038/srep34745>.
34. The Synergize Network (2023). Synergize. <https://synergize.xibe.org/>.
35. Dornelas, M., Antão, L.H., Moyes, F., Bates, A.E., Magurran, A.E., Adam, D., Akhmetzhanova, A.A., Appeltans, W., Arcos, J.M., Arnold, H., et al. (2018). BioTIME: a database of biodiversity time series for the Anthropocene. *Glob. Ecol. Biogeogr.* *27*, 760–786. <https://doi.org/10.1111/geb.12729>.
36. ForestPlots.net, Blundo, C., Carilla, J., Grau, R., Malizia, A., Malizia, L., Osinaga-Acosta, O., Bird, M., Bradford, M., Catchpole, D., et al. (2021). Taking the pulse of Earth’s tropical forests using networks of highly distributed plots. *Biol. Conserv.* *260*, 108849, <https://doi.org/10.1016/j.biocon.2020.108849>.
37. Grattarola, F., Botto, G., da Rosa, I., Gobel, N., González, E.M., González, J., Hernández, D., Laufer, G., Maneyro, R., Martínez-Lanfranco, J.A., et al. (2019). Biodiversidata: an open-access biodiversity database for Uruguay. *Biodivers. Data J.* *7*, e36226. <https://doi.org/10.3897/BDJ.7.e36226>.
38. Hazlett, M.A., Henderson, K.M., Zeitzer, I.F., and Drew, J.A. (2020). The geography of publishing in the Anthropocene. *Conserv. Sci. Pract.* *2*, <https://doi.org/10.1111/csp2.270>.
39. Carvalho, R., Resende, A., Barlow, J., Moura, M.R. Pervasive gaps in Amazonian ecological research. Zenodo. 2023; <https://doi.org/10.5281/zenodo.7951033>.
40. School of Geography, University of Leeds (2014). ForestPlots.NET. <https://forestplots.net/>.
41. University of Leeds (2023). Rainfor – Amazon Forest Inventory Network. <https://rainfor.org/en/>.
42. Ter Steege, H., Pitman, N.C.A., Sabatier, D., Baraloto, C., Salomão, R.P., Guevara, J.E., Phillips, O.L., Castilho, C.V., Magnusson, W.E., Molino, J.F., et al. (2013). Hyperdominance in the Amazonian tree flora. *Science* *342*, 1243092.
43. 2ndFOR. <https://sites.google.com/view/2ndfor/home>.
44. Taoca. <https://www.taoca.net/>.
45. Amazon Fish Project. <https://www.amazon-fish.com/pt>.
46. labecoufpa. <https://www.labecoufpa.com.br/contatos>.
47. de Oliveira-Junior, J.M.B., Shimano, Y., Gardner, T.A., Hughes, R.M., de Marco Júnior, P., and Juen, L. (2015). Neotropical dragonflies (Insecta: Odonata) as indicators of ecological condition of small streams in the eastern Amazon. *Austral Ecol.* *40*, 733–744.
48. Huete, A.R., Didan, K., Shimabukuro, Y.E., Ratana, P., Saleska, S.R., Hutyra, L.R., Yang, W., Nemani, R.R., and Myneni, R. (2006). Amazon rain-forests green-up with sunlight in dry season. *Geophys. Res. Lett.* *33*, L06405, <https://doi.org/10.1029/2005GL025583>.
49. Weiss, D.J., Nelson, A., Gibson, H.S., Temperley, W., Peedell, S., Lieber, A., Hancher, M., Poyart, E., Belchior, S., Fullman, N., et al. (2018). A global map of travel time to cities to assess inequalities in accessibility in 2015. *Nature* *553*, 333–336. <https://doi.org/10.1038/nature25181>.
50. e-MEC - Ministério da Educação. <https://emec.mec.gov.br/>.
51. GEOCAPES. <https://geocapes.capes.gov.br/geocapes/>.
52. International - Portal Embrapa. <https://www.embrapa.br/en/international>.
53. ICMBio - Instituto Chico Mendes de Conservação da Biodiversidade. <https://www.gov.br/icmbio/pt-br>.
54. Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A., and Michaelsen, J. (2015). The climate hazards infrared precipitation with stations - A new environmental record for monitoring extremes. *Sci. Data* *2*, 150066. <https://doi.org/10.1038/sdata.2015.66>.
55. da Rocha, H.R., Goulden, M.L., Miller, S.D., Menton, M.C., Pinto, L.D.V.O., de Freitas, H.C., and e Silva Figueira, A.M. (2004). Seasonality of water and heat fluxes over a tropical forest in eastern Amazonia. *Ecol. Appl.* *14*, 22–32. <https://doi.org/10.1890/02-6001>.
56. Bullock, E.L., Woodcock, C.E., Souza, C., and Olofsson, P. (2020). Satellite-based estimates reveal widespread forest degradation in the Amazon. *Glob. Chang. Biol.* *26*, 2956–2969. <https://doi.org/10.1111/gcb.15029>.
57. Matricardi, E.A.T., Skole, D.L., Costa, O.B., Pedlowski, M.A., Samek, J.H., and Miguel, E.P. (2020). Long-term forest degradation surpasses deforestation in the Brazilian Amazon. *Science* *369*, 1378–1382. <https://doi.org/10.1126/science.abb3021>.
58. Atlas Agropecuário <http://atlasagropecuario.imaflora.org/mapa>.
59. Shimada, M., Itoh, T., Motooka, T., Watanabe, M., Shiraishi, T., Thapa, R., and Lucas, R. (2014). New global forest/non-forest maps from ALOS PALSAR data (2007–2010). *Remote Sens. Environ.* *155*, 13–31. <https://doi.org/10.1016/j.rse.2014.04.014>.
60. Hess, L.L., Melack, J.M., Affonso, A.G., Barbosa, C., Gastil-Buhl, M., and Novo, E.M.L.M. (2015). Wetlands of the Lowland Amazon Basin: extent, vegetative cover, and dual-season inundated area as mapped with

- JERS-1 synthetic aperture radar. *Wetlands* 35, 745–756. <https://doi.org/10.1007/s13157-015-0666-y>.
61. Gumbrecht, T., Roman-Cuesta, R.M., Verchot, L., Herold, M., Wittmann, F., Householder, E., Herold, N., and Murdiyarto, D. (2017). An expert system model for mapping tropical wetlands and peatlands reveals South America as the largest contributor. *Glob. Chang. Biol.* 23, 3581–3599. <https://doi.org/10.1111/gcb.13689>.
  62. Venticinque, E., Forsberg, B., Barthem, R., Petry, P., Hess, L., Mercado, A., Cañas, C., Montoya, M., Durigan, C., and Goulding, M. (2016). An explicit GIS-based river basin framework for aquatic ecosystem conservation in the Amazon. *Earth Syst. Sci. Data* 8, 651–661. <https://doi.org/10.5194/essd-8-651-2016>.
  63. Breiman, L. (2001). *Random Forests*. *Mach. Learn.* 45, 5–32.
  64. Mi, C., Huettmann, F., Guo, Y., Han, X., and Wen, L. (2017). Why choose Random Forest to predict rare species distribution with few samples in large undersampled areas? Three Asian crane species models provide supporting evidence. *PeerJ* 5, e2849. <https://doi.org/10.7717/peerj.2849>.
  65. Kutner, M.H., Nachtsheim, C.J., and Neter, J. (2004). *Applied Linear Regression Models*, 4th ed. (McGraw-Hill).
  66. Naimi, B. (2013). usdm: Uncertainty analysis for species distribution models. Version 1.1-12. <http://cran.r-project.org/web/packages/usdm/index.html>.
  67. Barbet-Massin, M., Jiguet, F., Albert, C.H., and Thuiller, W. (2012). Selecting pseudo-absences for species distribution models: how, where and how many? *Methods Ecol. Evol.* 3, 327–338. <https://doi.org/10.1111/j.2041-210X.2011.00172.x>.
  68. Valavi, R., Elith, J., Lahoz-Monfort, J.J., and Guillera-Aroita, G. (2021). Modelling species presence-only data with random forests. *Ecography* 44, 1731–1742. <https://doi.org/10.1111/ecog.05615>.
  69. Kuhn, M. (2008). Building predictive models in R using the caret package. *J. Stat. Softw.* 28, 1–26. <https://doi.org/10.18637/JSS.V028.I05>.
  70. Liaw, A., and Wiener, M. (2002). Classification and regression by randomForest. *R. News* 2, 18–22.
  71. Proosdij, A.S.J., Sosef, M.S.M., Wieringa, J.J., and Raes, N. (2016). Minimum required number of specimen records to develop accurate species distribution models. *Ecography* 39, 542–552. <https://doi.org/10.1111/ecog.01509>.
  72. Leroy, B., Delsol, R., Hugueny, B., Meynard, C.N., Barhoumi, C., Barbet-Massin, M., and Bellard, C. (2018). Without quality presence–absence data, discrimination metrics such as TSS can be misleading measures of model performance. *J. Biogeogr.* 45, 1994–2002. <https://doi.org/10.1111/jbi.13402>.
  73. Gutiérrez, J.M., Jones, R.G., Narisma, G.T., Alves, L.M., Amjad, M., Gorodetskaya, I.V., Grose, M., Klutse, N.A.B., Krakovska, S., Li, J., Martínez-Castro, D., et al. (2021). Atlas supplementary material. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, V. Masson-Delmotte, P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, and M.I. Gomis, et al., eds. (The Intergovernmental Panel on Climate Change). <https://www.ipcc.ch/>.
  74. Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R. (2017). Google Earth Engine: planetary-scale geospatial analysis for everyone. *Remote Sens. Environ.* 202, 18–27. <https://doi.org/10.1016/j.rse.2017.06.031>.
  75. R Development Core Team (2021). *R: A language and environment for statistical computing* (R Foundation for Statistical Computing).

## STAR★METHODS

### KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
<b>Deposited data</b>		
Raw data and code for the analyses on research probability across the Brazilian Amazoni	This paper	<a href="https://zenodo.org/record/7951033">https://zenodo.org/record/7951033</a>
<b>Software and algorithms</b>		
R software	The R Foundation / R Development Core Team	<a href="https://www.r-project.org">https://www.r-project.org</a>
Rstudio IDE	AGPL v3 / Posit-PBC	<a href="https://posit.co/products/open-source/rstudio/">https://posit.co/products/open-source/rstudio/</a>
Google Earth Engine	Google LLC	<a href="https://earthengine.google.com/">https://earthengine.google.com/</a>
QGIS	QGIS Development Team	<a href="https://www.qgis.org/en/site/">https://www.qgis.org/en/site/</a>
Inkscape	GNU GPL-2.0-or-later	<a href="https://inkscape.org/">https://inkscape.org/</a>

### RESOURCE AVAILABILITY

#### Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the Lead Contact, Raquel L. Carvalho ([raqueluly@gmail.com](mailto:raqueluly@gmail.com)).

#### Materials availability

This study did not generate new reagents, sequences or eventuate in the archiving of specimens.

#### Data and code availability

- Metadata have been deposited at Zenodo and are publicly available as of the date of publication. DOIs are listed in the [key resources table](#).
- All original code has been deposited at Zenodo and is publicly available as of the date of publication. DOIs are listed in the [key resources table](#).
- Any additional information required to reanalyse the data reported in this paper is available from the lead contact upon request.

### EXPERIMENTAL MODEL AND SUBJECT DETAILS

#### Data collection

All data collection took place within the SYNERGIZE (*Synthesising Ecological Responses to Degradation in Amazonian Environments*) project, a collaborative effort of scientists from institutions in Brazil and other countries that integrate and synthesise data on Amazonian biodiversity studies. More information about the project and study region can be found elsewhere ([key resources table](#)).<sup>34,39</sup> We gathered ecological community datasets for nine organism groups (benthic invertebrates, heteropterans, odonates, fishes, macrophytes, birds, woody vegetation, ants, and dung beetles) between 2010 and 2020 in the Brazilian Amazonia. We focused on this period to keep results relevant to the recent potential drivers of research effort.

To be included in our database, any published research needed to be considered as an ecological community dataset derived from quantitative and repeatable sampling protocols, with sampled taxa identified at a minimum of family level. For each selected study (see [Table S1](#)), we contacted the first and/or corresponding author. Because work published in English may be a biased research subset, we extended our search by contacting experts known to work in the Brazilian Amazonia. In the following, we have summarized the information derived from the compiled data (but see [Table S1](#)).

#### Woody vegetation

We first compiled the metadata available in three consolidated databases: [ForestPlots.net](#),<sup>36,40</sup> Amazon Tree Diversity Network (Rainfor,<sup>41</sup> ATDN<sup>42</sup>), and Secondary Forests Research Network (2ndFOR<sup>43</sup>) for Brazil. These databases have deployed huge efforts to acquire data and metadata across the Brazilian Amazonia. For the ForestPlots database, we contacted 79 people to request data owners' permission to use the metadata. In 1% of the number of people contacted, we had no answers from any team member and, consequently, no permission to use metadata. In parallel, we listed all the universities and research institutes with graduate courses in ecology or related fields in all states covering the Brazilian Amazonia, and invited additional

90 data owners to collaborate with their metadata (Table S1). Overall, 47 authors returned our contact and shared metadata on 2,597 inventories meeting our criteria.

### Terrestrial animals

We focused on published studies on ants, birds, dung beetles in Amazonia using the Web of Science platform (Table S1 for string words used and search dates). Searches were conducted in English and with no restriction on year. We did not focus on a search period between 2010 and 2020 for these groups, as the Synergize project is developing a database that goes beyond the period considered here TAOCA.<sup>44</sup> From the initial total of 2,244 published manuscripts obtained through the platform, only 225 were ecological studies. Among these 225 studies, we contacted 99 authors, many of whom were corresponding authors of more than one article. Ninety one percent of these researchers returned our contact and shared metadata of their studies (Table S1).

### Aquatic groups

We focused on published studies of Amazon fishes using the Web of Science platform (Table S4 for string words used and search dates). Since fishes are also research subjects in many areas of knowledge (such as social food sciences), we selected only articles related to biodiversity and conservation areas. For other aquatic groups (benthic, heteropterans, odonates and macrophytes), we focused on a period between 2010 and 2020. We divided aquatic invertebrates into benthic, heteropterans, odonates, because most datasets on heteropterans and odonates referred to sampling of adults near streams, while for benthic invertebrates (Ephemeroptera, Plecoptera, Trichoptera, Diptera and Coleoptera), the datasets corresponded to the sampling of larvae in streams. From the initial total of 1,073 published manuscripts obtained through the platform, only 523 were ecological studies. Among these 523 studies, we contacted 69 authors, many of whom were corresponding authors of more than one article. Eighty-one percent of these researchers returned our contact and shared metadata of their studies (Table S1).

All sampling sites – informed within the ecological studies herein considered – had their geographic coordinates verified and cross-checked with databases of the political and environmental limits of the Brazilian Amazon. Studies were included if data collection utilized repeatable methods, which varied among different biological groups (see details in Table S1). Organism groups were sampled along transects, grids, or plots, which exhibited significant variations in size.

Our original dataset comprised 3,917 sampling sites in uplands (635 for ants, 742 for birds, 687 for dung beetles, and 1,853 for woody vegetation), 923 in wetlands (18 for ants, 136 for birds, 25 for dung beetles, and 44 for woody vegetation) and 2,897 in aquatic habitats (536 for benthic invertebrates, 1710 for fishes, 153 for heteropterans, 159 for macrophytes and 339 for odonates), totalling 7,737 sampling sites.

Metadata used in this study, including habitat, year of study, organism groups, and geographical coordinates (longitude and latitude) can be found in the STAR Methods (see data and code availability). Other metadata and community data can be found in the original sources, which varied between groups. For woody vegetation, data is available in three consolidated databases: ForestPlots,<sup>36,40</sup> Amazon Tree Diversity Network (Rainfor,<sup>41</sup> ATDN<sup>42</sup>), and Secondary Forests Research Network (2ndFOR<sup>43</sup>). For terrestrial fauna, the data are deposited at the TAOCA platform.<sup>44</sup> Fish group data is available in the AmazonFish,<sup>45</sup> aquatic invertebrates and macrophytes are available, under request, with Leandro Juen and Thaise Michelan, respectively, both from the Laboratory of Ecology and Conservation (UFPA, Brazil<sup>46</sup>).

## METHOD DETAILS

### Predictors of research probability

To model research probability for each organism group, we used predictors related to logistics (accessibility, research facilities, and dry season length), and human influence (land tenure, degradation, and dry season length). We define human influence as those variables used to assess the ecological condition of a particular habitat or ecosystem.<sup>47</sup> We used logistics-related variables as certain areas are easier to work in due to accessibility, proximity to research centres, or location within public protected areas.<sup>19</sup> As the dry season length affords extended accessibility to many of the drier regions in the Brazilian Amazon each year, we deemed it a crucial logistical factor.<sup>48</sup> Although predictor variables change over time, we assumed that these changes in a short time window are negligible relative to the variation across space and adopted a static layer to represent the Brazilian Amazonia over the ten years from which data were collected.

### Accessibility

We used travel time from each surveyed site to the nearest village or city, considering a population between five thousand to five million people in 2015. The travel time to the nearest major city indicates the fastest travel speed considering all transportation facilities (road, rail, river, canal, and sea lane) as well as characteristics of land cover, slope, and elevation of the areas. We used the dataset available in Weiss and collaborators,<sup>49</sup> downloaded at the spatial resolution of 30 arc-sec (Figure S1A).

### Research facilities

We used the geographical location of institutes and universities that offer undergraduate and graduate courses related to biology, ecology, and forestry, as well as experimental farms, forest reserves, and other institutions that support these courses. We obtained the geographical distribution of undergraduate courses from the Brazilian Education Ministry,<sup>50</sup> and we extracted information on graduate courses from the Coordination for the Improvement of Higher Education Personnel (*Coordenação de Aperfeiçoamento de Pessoal de Nível Superior* – CAPES<sup>51</sup>). We also gathered data on institutes from the Ministry of Agriculture, Livestock and Supply (*Ministério da Agricultura, Pecuária e Abastecimento* – MAPA<sup>52</sup>) and Ministry of the Environment and Climate Change (*Ministério do Meio Ambiente* – MMA<sup>53</sup>). In addition, we gathered information on experimental farms and forest reserves from the website of each

University or Research Institution, covering all Brazilian states that spatially overlap with Amazonia. Then, we rasterised this predictor at 8 arc-sec (~ 0.25 km) resolution. Finally, we extracted the distance from each surveyed site to the nearest university or infrastructure containing research facilities (research facility hereafter) (Figure S1B).

#### Dry season length

We used the average dry season length in consecutive months, extracted from the Climate Hazards Group Infrared Precipitation with Stations (CHIRPS) dataset.<sup>54</sup> The CHIRPS dataset was created from interpolation techniques of satellite data and rainfall station data from 1981 to the present.<sup>54</sup> We considered dry months to be those with less than 100 mm of precipitation.<sup>55</sup> This variable ranged from zero to eight months and originally presented a 2.5 arc-min (~5 km) resolution (Figure S1D).

#### Degradation

We extracted the accumulated degradation from 1995 to 2017 using a 1 arc-sec (~0.03-km) resolution layer developed by Bullock and collaborators.<sup>56</sup> This is the most suitable and recent dataset for the period considered in this study (but see Matricardi et al.<sup>57</sup>). However, it does not cover a small area of our study – the Amazon Network of Georeferenced Socio-Environmental Information-RAISG – which led us to redefine our study area limits according to Bullock and collaborators.<sup>56</sup> Therefore, 3.2% of sampling sites in our original database were removed from analyses described below as they fell outside the RAISG-defined Amazonia biome (see Figure S1F). Degradation was considered as any natural or anthropogenic disturbance (e.g., fire, windthrow, selective logging, and fuelwood) that does not alter a pixel's original land cover forest category. To assess the forest degradation in the vicinity of the sampling area, we counted the proportion of degraded forest in a circle with a radius of 1000 m surrounding each plot. We used 1 km as a compromise between capturing enough forests to get a reasonable degradation sample while minimising the overlap between the buffers of neighbouring plots. The mean geographic distance between plots was 926.9 km (range: 0–2536.3 km, Figure S1E).

#### Land tenure

We obtained a land tenure map at 1 arc-sec (~0.03-km) of spatial resolution for the year 2017.<sup>58</sup> We classified the surveyed sites in nine categories: (1) Indigenous lands, both unofficial and official land by the National Foundation for Indigenous People; (2) Unassigned public land that has not been registered in National Institute of Colonisation and Agrarian Reform (INCRA) database and/or incorporated into other public land categories; (3) Private land recognised by public government databases [e.g. INCRA and Environmental Rural Registry]; (4) Protected area, with sustainable use of natural resources (i.e. low-level non-commercial use of resources allowed [category VI, IUCN]); (5) Strict Reserve area, where the use and management of natural resources are strictly controlled and limited to science or wilderness preservation [category IA, IUCN]; (6) Rural settlements registered by INCRA; (7) Water body, including continental and/or coastal waters; (8) Quilombola land, afro-Brazilian settlements established by escaped or enslaved people up to the 19<sup>th</sup> century; and (9) Other, including railway, road, military and urban areas (Figure S1C).

## QUANTIFICATION AND STATISTICAL ANALYSIS

### Research probability modelling

Before running the analyses, we defined the habitats (upland, wetland and aquatic) using existing forest/non-forest and wetland layers. We removed the non-forested areas from our study areas using a 25 m resolution ALOS PALSAR 2 Forest/Non-Forest map for 2015.<sup>59</sup> Next, we obtained the wetland delimitation by complementing the Hess and collaborators<sup>60</sup> layer, which covers the lowland Amazonia Basin wetlands (below 500 m asl), with the wetlands layer from the Center for International Forestry Research (CIFOR<sup>61</sup>) to cover higher altitudes and eastern Amazon Forest areas outside the basin (Figure S1F). For the wetland habitat, we removed upland and water surface, keeping only forested areas. Lastly, we used the same wetland delimitation for aquatic habitat but included open water bodies by joining a river level layer.<sup>62</sup> We removed only the areas that did not meet the inclusion criteria (being the habitat where each taxon occurs and being forest for upland and wetland habitats; Figure S1F). We considered a 1 km spatial resolution and the extension of the degradation layer to run all analyses. In addition, we used equal-area projections for all our layers (CRS WGS 84 – EPSG: 4326).

We used occurrence data of community ecological studies (i.e., surveyed sites) in a Random Forest (RF) framework<sup>63,64</sup> to model the ecological probability of sampling research for each biological group in each pixel (1 km) across the Brazilian Amazon. Before building RF models, we tested for multicollinearity among predictors (accessibility, research facility, land tenure, dry season length, and degradation layers) using the Variance Inflation Factor (VIF). Computations were performed separately for each combination of biological group and habitat type. Since all predictors showed VIF < 2, we retained each variable in the analyses<sup>65</sup> (Table S2). We undertook VIF analyses using 'usdm' package.<sup>66</sup>

We built RF models of ecological research probability (hereafter, research probability) separately for each organism and habitat. Like many species distribution modelling techniques, RF requires absence or pseudoabsence data, and shows better performance when used with balanced classes (i.e., pseudoabsence ratio of 1:1,<sup>67,68</sup>). Therefore, we computed pseudoabsences using the same number of observed presences in each organism group. We validated the RF models using a 5-fold cross-validation approach. The presence-pseudoabsence data were randomly partitioned into five equal parts, with four of those used for training and the fifth as validation-fold.<sup>69</sup> For each RF model, we initially used the *tuneRF* function in the randomForest package<sup>70</sup> to identify the number of predictors (*mtry* argument) and trees (*ntree* argument) returning the most accurate RF, that is, the model with the lowest mean-squared error (MSE) and highest explained deviance ( $R^2$ ). RF computations were then repeated for each training-fold using the optimal parameters identified by *tuneRF*. We excluded ants and dung beetles from the RF models for wetland analyses, since they had fewer than 25 sampling sites.<sup>71</sup>

To evaluate model performance, we used the Sørensen similarity index to measure the similarity between predictions and observations. This index is independent of the prevalence of sampled sites (ratio of observed presences to all sites) and less sensitive to under/overprediction issues than metrics based on specificity, such as the True Skill Statistic – TSS.<sup>72</sup> We measured variable importance using the proportional increase in MSE, which measures the relative decrease in model accuracy by shuffling variable values. We also built partial dependence plots to represent the marginal effects of predictor values on research probability across taxa and habitats.

### Overlap between research and anthropogenic disturbances

We intersected research probability with susceptibility to current and future anthropogenic disturbances to identify areas with ecological knowledge most at risk. We used three indicators of susceptibility to anthropogenic disturbances that reflect continuous trends of long-term demographic growth and economic development: climate change,<sup>73</sup> deforestation, and degradation.<sup>20</sup>

To indicate climate change, we computed the difference between current and future projections of climate represented through 13 bioclimatic variables (e.g.,  $\Delta\text{Bio1} = \text{Bio1}_{\text{future}} - \text{Bio1}_{\text{current}}$ ) obtained from the Intergovernmental Panel on Climate Change (IPCC) Interactive Atlas<sup>73</sup> (Table S4). For each bioclimatic variable, the  $\Delta\text{climatic}$  values were measured between projections for 2041–2060 and 1981–2010 under the SSP585 scenario. To provide a consensus metric of future climate change across different generalised circulation models (GCM), we average  $\Delta\text{climatic}$  values across all GCM available at the IPCC Interactive Atlas (Table S4). We rescaled the  $\Delta\text{climatic}$  values in the interval between -1 and 1 and passed their absolute values ( $|\Delta\text{climate}|$ ) through a Principal Component Analysis (PCA) to remove multicollinearity. The magnitude of climate change for each pixel was calculated as the Euclidean distance between its position in the two first axes PCA space and the origin; which represents a reference point of no climate change (where  $|\Delta\text{climate}| = 0$  along all axes).

To identify areas most threatened by deforestation and degradation, we used projections made recently available indicating trends under a business-as-usual scenario for 2050.<sup>20</sup> The term deforestation refers to the complete removal of canopy cover, whereas degradation is the term used to describe a natural or human-induced disturbance that does not alter the land cover category assigned to a pixel.<sup>56,57</sup> Currently deforested areas are also shown since they may be reforested in case of land abandonment or political incentives. We prepared the spatial layers using Google Earth Engine (GEE<sup>74</sup>) and carried out all statistical analyses in R 4.0.5.<sup>75</sup>

Next, both measures of anthropogenic disturbances, (i) magnitude of climate change and (ii) combined deforestation and degradation, as well as research probability, were split into equal-sized quantiles holding 0–25, 25–50, 50–75, and 75–100% of samples (pixels). We used the first quartile of research probability (0–25%) to identify the most neglected areas in ecological research, and the last quartile (75–100%) of climate change and deforestation-degradation to identify areas most susceptible to anthropogenic disturbances.

### ADDITIONAL RESOURCES

For terrestrial fauna, the metadata used in this study were stored in the TAOCA database. This database emerged in response to the demand for organizing, standardizing, and securely storing a large amount of data received by the Synergize project (<https://www.taoca.net/>).