

Geophysical Research Letters®

RESEARCH LETTER






10.1029/2023GL106132

Flawed Emergency Intervention: Slow Ocean Response to Abrupt Stratospheric Aerosol Injection



Key Points:

- Efficacy of stratospheric aerosol injection (SAI) impaired by anthropogenic ocean heating
- Deep ocean heating, weakened Atlantic Meridional Overturning Circulation (AMOC) and collapsed North Atlantic deep convection only partially addressed by late SAI
- SAI decouples AMOC and global mean surface temperature, thereby inducing climate states not seen in purely greenhouse gas-forced scenarios

Daniel Pflüger¹ , Claudia E. Wieners¹ , Leo van Kampenhout¹ , René R. Wijngaard¹ , and Henk A. Dijkstra¹ 

¹Institute Marine and Atmospheric Research Utrecht, Utrecht, The Netherlands

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

D. Pflüger,
d.pflug@uu.nl

Citation:

Pflüger, D., Wieners, C. E., van Kampenhout, L., Wijngaard, R. R., & Dijkstra, H. A. (2024). Flawed emergency intervention: Slow ocean response to abrupt stratospheric aerosol injection. *Geophysical Research Letters*, 51, e2023GL106132. <https://doi.org/10.1029/2023GL106132>

Received 29 AUG 2023

Accepted 17 JAN 2024

Author Contributions:

Conceptualization: Daniel Pflüger, Claudia E. Wieners
Data curation: Leo van Kampenhout, René R. Wijngaard, Henk A. Dijkstra
Formal analysis: Daniel Pflüger
Funding acquisition: Claudia E. Wieners
Investigation: Daniel Pflüger, Leo van Kampenhout, René R. Wijngaard, Henk A. Dijkstra
Methodology: Daniel Pflüger, Claudia E. Wieners, Leo van Kampenhout, René R. Wijngaard, Henk A. Dijkstra

© 2024. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs License](#), which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Abstract Given the possibility of irreversible, anthropogenic changes in the climate system, technologies such as solar radiation management (SRM) are sometimes framed as possible emergency interventions. However, little knowledge exists on the efficacy of such deployments. To fill in this gap, we perform Community Earth System Model 2 simulations of an intense warming scenario on which we impose gradual early-century SRM or rapid late-century cooling (an emergency intervention), both realized via stratospheric aerosol injection (SAI). While both scenarios cool Earth's surface, ocean responses differ drastically. Rapid cooling fails to release deep ocean heat content or restore an ailing North Atlantic deep convection but partially stabilizes the Atlantic meridional overturning circulation. In contrast, the early intervention effectively mitigates changes in all of these features. Our results suggest that slow ocean timescales impair the efficacy of some SAI emergency interventions.

Plain Language Summary Stratospheric aerosol injection (SAI) is a promising, yet controversial proposal to mask the effects of anthropogenic climate change by releasing sunlight-reflecting particles into the atmosphere. Currently, many studies are focusing on the benefits of near future SAI deployments. We, however, investigate SAI as a late *emergency intervention*. To what extent can SAI still help if we continue to heat and destabilize the climate? In this study, we simulate the impacts of an abrupt, SAI cooling intervention deployed against the backdrop of a climate much hotter than today's. While SAI readily cools Earth's surface, it is challenged by a slow ocean response. Heat trapped below the ocean surface remains a contributor to sea-level rise and important currents weakened by climate change linger in ailing condition. In contrast, an earlier SAI intervention effectively mitigates changes in these features. Our findings re-emphasize the urgent need for climate action. If anthropogenic heating continues, even an intervention as powerful as SAI will encounter its limits.

1. Introduction

While global heating puts increasing pressure on societies and ecosystems (IPCC, 2022a), current policies are insufficient to prevent 1.5°C or even 2°C of warming (IPCC, 2022b). To mitigate the associated risks, interventions that cool Earth by reflecting sunlight—*Solar Radiation Management* (SRM)—are being explored as complementary measures to emission reductions (National Academies of Sciences, Engineering, and Medicine, 2021). Among several potential schemes, *Stratospheric Aerosol Injection* (SAI) received considerable attention due to its low perceived technical barriers (Smith, 2020), plausible physical effectiveness (Kleinschmitt et al., 2018; Plazzotta et al., 2018). While model studies demonstrate its benefits (Tilmes et al., 2018, 2020; Visioni et al., 2021), including its ability to control global mean surface temperature (GMST), SAI can not address all consequences of rising greenhouse gas (GHG) concentrations and may induce side-effects of its own (Irvine et al., 2016; Zarnetske et al., 2021). Ethical concerns (Oomen, 2021; Svoboda, 2017) lead some to suggest a ban on its research and deployment (Biermann et al., 2022) whereas others suggest further research (Wieners et al., 2023).

It is not enough to ask *whether* SRM should be deployed. Multiple degrees of freedom in SRM deployments implore us to ask *how and to what end* may be SRM used. Currently popular frameworks include *peak-shaving* (Long & Shepherd, 2014; Reynolds, 2019), in which SRM stabilizes GMST, while other measures gradually tackle GHG concentrations. However, there is no guarantee SRM would be deployed in such a well-optimized and *proactive* fashion. In this study, we examine as SRM as an *emergency* intervention instead, to be deployed only after prolonged heating. This notion, adapted from Lockley et al. (2022), naturally arises when SRM deployments are restricted to particularly extreme situations such as rapid climate tipping. To what extent

Project administration: Claudia E. Wieners
Software: Daniel Pflüger, Leo van Kampenhout, René R. Wijngaard, Henk A. Dijkstra
Supervision: Claudia E. Wieners, Henk A. Dijkstra
Visualization: Daniel Pflüger
Writing – original draft: Daniel Pflüger, Claudia E. Wieners
Writing – review & editing: Daniel Pflüger, Claudia E. Wieners, Henk A. Dijkstra

can later deployments reverse the impacts of heating? How would they compare to earlier, proactive interventions?

In this study, we focus on SRM's physical impact on the ocean. There, long response timescales hamper prospects of reversibility under an emergency intervention. Many ocean features are subject to anthropogenic climate change and have profound impacts on humans and ecosystems which elevates the study of them above a purely academic exercise. We are interested in

- ocean heat content (OHC) change, a major contributor to sea-level rise (Church et al., 2013).
- the Atlantic Meridional Overturning Circulation (AMOC) which may weaken (or even collapse) in the future (Weijer et al., 2020), thereby reducing meridional heat transport (MHT) and modulating regional sea level rise.
- North Atlantic deep convection which may shut down in the future, leading to abrupt cooling and shifts in the jet-stream (Sgubin et al., 2017; Swingedouw et al., 2021).

We consider only SRM scenarios with extreme levels of GHG and aerosol forcing, including abrupt changes thereof. They should not be seen as desirable or politically realistic futures but rather as physical edge cases that provide valuable intuitions and constraints for more cautious scenarios: if an abrupt cooling struggles to reverse certain ocean changes, a slower deployment would likely do so, too. Furthermore, we restrict ourselves to a single SRM implementation: planetary-scale SAI.

2. Methods

Our scenarios are simulated in CESM2 (Danabasoglu et al., 2020) with atmospheric component CAM6 at $1^\circ \times 1^\circ$ horizontal resolution and ocean model POP2 at similar resolution. Ice sheets are non-evolving, which also prohibits calving, but the land model CLM provides glacial run-off fluxes.

SAI is implemented via prescribed aerosol fields: a compromise between physical realism and computational cost. We favored this approach as it may enable computationally cheap simulations capturing longer ocean timescales in the future. Other groups have used similar scaling-based implementations (Visioni et al., 2021).

Schematically, the protocol works as follows:

- Every year, observe the deviation of GMST from the target
- Based on past GMST deviations, infer the level of SAI—expressed in terms of global mean aerosol optical depth (AOD)—which is necessary to achieve the desired target.
- Use the AOD to scale all SAI-related aerosol fields appropriately.
- Feed the scaled fields into CAM6.

The first two steps are implemented via an established feedforward-feedback control algorithm (Kravitz et al., 2016, 2017). Our specific implementation stabilizes GMST as its sole objective, whereas interactive aerosol simulations (MacMartin et al., 2017; Tilmes et al., 2020) can also control other features such as the inter-hemispheric temperature contrast.

The input aerosol fields derive from an interactive aerosol simulation performed by Tilmes et al. (2020) in CESM2-WACCM, more specifically their Geo SSP5 8.5 1.5 scenario. In contrast to CAM6, the improved, albeit more costly, atmospheric component WACCM allows for detailed chemical aerosol dynamics (Danabasoglu et al., 2020). Our prescribed aerosol fields are averaged versions of the WACCM aerosol fields as described in Supporting Information S1.

We simulate three scenarios based on SSP5-8.5 background concentrations:

- Control (2015–2100): historical spin-up continued by SSP5-8.5
- SAI 2020 (gradual SAI): branch off Control in 2020; stabilize GMST at 1.5°C above pre-industrial conditions; analogous to Geo SSP5-8.5 1.5 by Tilmes et al. (2020).
- SAI 2080 (emergency intervention): branch off Control in 2080, deploy SAI to restore GMST to 1.5°C .

Note that SAI 2080 involves some adjustments to the control algorithm, described in Supporting Information S1. Otherwise, the initially high deviation from the target GMST can overcharge the feedback controller and risk excessive cooling.




Figure 1. (a) Annual mean global mean surface temperature above pre-industrial reference temperature (b) Change in annual mean total depth ocean heat content (OHC) relative to 2020–2030 conditions in Control. (c) Difference in vertical OHC between end-of-simulation (2090–2100) conditions and present-day conditions in Control.

3. Results

3.1. Temperature Response

In Figure 1a, we see that the gradual SAI strategy (SAI 2020) indeed stabilizes GMST at target level. By contrast, SAI 2080 experiences rapid cooling and even shoots past the target. This undercooling is an artifact of the feedback controller and can be removed by fine-tuning the cooling process (Figure S2 in Supporting Information S1).

Even though GMST is stabilized, total depth OHC accumulates continuously in SAI 2020 (Figure 1b) in agreement with past studies (Fasullo et al., 2018; MacMartin et al., 2022). The warming takes place below the surface and likely stems from deep ocean response timescales (Cheng et al., 2022) combined with the goal of maintaining GMST. As sub-surface layers have not yet adapted to increased surface temperatures, they act as a heat sink for the ocean surface. The induced downward heat flux is then compensated by the feedback controller that allows for a residual top-of-atmosphere radiative imbalance in order to stabilize GMST.

SAI 2080 accumulates more total depth OHC than SAI 2020. The deep tail of OHC in SAI 2080 (Figure 1c) matches that of Control while the near-surface layers are cooled to SAI 2020 levels. Given the short time-frame of SAI 2080, it is not clear whether the vertical OHC distribution has reached equilibrium or deeper layers are simply cooling very slowly.

On the surface, however, both SAI scenarios have comparable OHC anomalies. This suggests that while abrupt SAI readily cools the ocean surface, heat anomalies trapped in deeper layers are more persistent.

Surface temperature responses to SAI are spatially inhomogeneous (Figure 2). Most strikingly, the subpolar North Atlantic is significantly undercooled in both SAI scenarios. This pattern resembles a *North Atlantic warming hole*




Figure 2. (a) Reference (2020–2030) annual mean near-surface air temperatures in Control (b–d) Late-century (2090–2100) temperature changes with respect to the reference for Control, stratospheric aerosol injection (SAI) 2020 and SAI 2080 respectively. (e) Difference between SAI scenarios (d) minus (c).




Figure 3. (a) Annual mean Atlantic northwards heat transport at 26° N where we apply a rolling average over 5 year periods with backward window (b) Atlantic Meridional Overturning Circulation (AMOC) index defined as the maximum of the annual mean meridional overturning streamfunction at 26° N below 200 m—Partially transparent uncertainty bands depict three CESM2 CMIP6 (Coupled Model Intercomparison Project Phase 6) ensemble members (Danabasoglu, 2019a, 2019b) per greenhouse gas concentration pathway. The uncertainty is the ensemble standard deviation. Again, we apply rolling averages over 5 year periods. (c) Annual mean global mean surface temperature versus AMOC index—The marker saturation denotes the year: light (2020) to dark (2100).

known from purely GHG-forced simulations (Drijfhout et al., 2012; Menary & Wood, 2018), which to some extent is also visible in Control. SAI may have merely unmasked this feature rather than induce it. The warming hole is expanded and colder in SAI 2080, while the Southern Hemisphere is warm compared to SAI 2020.

Multi-objective feedback procedures (Kravitz et al., 2017; MacMartin et al., 2017) allow for a more elaborate control of the global temperature pattern including the interhemispheric temperature gradient. Therefore, the asymmetric response of SAI 2080 (Figure 2e) may be mitigated in a refined control scheme. In our study, however, both SAI scenarios use spatially identical aerosol patterns which rules out a control of the asymmetry.

3.2. AMOC Response

The AMOC index and meridional heat transport (MHT) roughly halve in Control (Figures 3a and 3b). Even the low-emission SSP1-2.6 scenario is projected to lead to similar AMOC index changes. SAI 2020 drastically mitigates but does not halt the AMOC and MHT decline, similar to other studies (MacMartin et al., 2022; Xie et al., 2022). SAI 2080 stabilizes the AMOC index but only has an inconclusive impact on the MHT.

SAI effectively decouples the GMST and the AMOC index (Figure 3c). This could explain the interhemispheric temperature contrast featured in SAI 2080: a weak AMOC impedes northward heat transport leading to a see-saw temperature pattern (Liu et al., 2020; Stocker, 1998) not masked by heat otherwise present in Control.

To study the spatial pattern of the AMOC, we plot meridional streamfunction changes under all scenarios from 2070 to 2080 to 2090–2100 (Figure 4). This choice of time intervals helps to reveal the immediate AMOC response to SAI 2080. Additionally, we subtract the changes in Control from the ones in the SAI scenarios in an attempt to disentangle GHG from SAI-related impacts.

Figure 4d reveals a potential feedback in the AMOC stabilization under SAI 2080. Following the deployment, the pattern of relative AMOC strengthening closely mirrors the pre-deployment streamfunction, albeit mostly near the surface and in the northern hemisphere. This suggests that the AMOC response to abrupt SAI is dependent on the AMOC state itself. While a similar observation can be made for SAI 2020 (Figure 4c), disentangling the forced response from internal feedback is not obvious during the gradual change in aerosol forcing. SAI 2080 gives a much better indication that it is indeed the state of the AMOC which steers its response to SAI.

3.3. North Atlantic Deep Convection

We now focus on deep convection processes in the North Atlantic. Using mixed layer depth as a proxy for deep convection, we identify two regions, *East* and *West*, where the mixed layer depth in April (the month with the




Figure 4. (a) Atlantic Meridional Overturning Circulation (AMOC) streamfunction in Control averaged over 2020–2030. In panels (b)–(d), for any simulation X, ΔX is the mean over 2090–2100 minus the mean over 2070–2080. (b) Change in AMOC streamfunction under Control—Black contour lines show the mean streamfunction over 2070–2080 for Control while the shading indicates Δ Control. (c) Change in AMOC streamfunction in stratospheric aerosol injection (SAI) 2020 relative to Control—Black contour lines show the mean streamfunction over 2070–2080 for SAI 2020 while the shading indicates Δ SAI 2020— Δ Control. (d) Analogous to (c) but for SAI 2080.

deepest mixed layer) exceeds 550 m (Figure 5a). This threshold depth was chosen as it is sufficiently large to distinguish deep convection from regular mixed-layer conditions and small enough to provide a good signal-to-noise ratio. The regions are separated longitudinally by the southern tip of Greenland.

In Control, deep convection in *West* ceases around 2050, followed by a shutdown in *East* around 2060. SAI 2020 prevents the shutdown in *East*, but only postpones the shutdown in *West* by about a decade. The *West* shutdown is not as definite as in the case of Control and isolated years with deep convection still occur. For SAI 2080, deep convection remains absent in both regions with the exception of a single outlier year for *East*.

Why does cooling in SAI 2080 not revive deep convection before 2100? We address this question by studying the ocean stratification over both deep convection regions. Deep convection in April is inhibited if the surface density in the previous September has been too low, that is, the water column was too stratified (Figure S4 in Supporting Information S1). Thus, surface density serves as a proxy for favorable convection conditions.

The sea surface density is determined by temperature and salinity, seen in Figures 5d–5f. In all scenarios, final salinities are well below reference conditions. SAI 2020 roughly halves the decline with respect to Control. This difference becomes very noticeable mid-century simultaneously with the *East* and *West* shutdown in Control. SAI 2080 does not fundamentally alter the trajectory of Control apart from a transient increase in salinity that correlates with an isolated year of deep convection. Therefore, freshening contributes to density loss in all scenarios.

In the case of Control, temperature trends are non-monotonous (Figure 5d) and do not lead to a denser surface. What appears to be a weak cooling trend is mostly masked by inter-annual variability and eventually superseded by intense heating. Typically, deep convection shutdown induces a rapid cooling (Sgubin et al., 2017; Swingedouw et al., 2021) which is not obvious from Figure 5d. It can, however, be detected by using a CESM2 SSP5-8.5 ensemble and switching to an annual-mean rather than a single-month perspective (Figure S6 in Supporting Information S1).




Figure 5. (a) North Atlantic April mixed layer depths in CESM2 (2020–2030)—*East* and *West* are enclosed by solid and dashed lines respectively. Shutdown dates are denoted with a cross and colored according to scenario (blue: Control, green: SAI 2020). (b–c) April mixed layer depths in *West* and *East* respectively—Solid lines are 5 year rolling means (with backward window) applied to the data shown by transparent lines. (d–f) September mean sea surface density, temperature and salinity over the total *East* and *West* region.

SAI 2020 shows an overall cooling trend dominated by a quick decline at time of *West* shutdown. Former observation could indicate AMOC weakening whereas latter phenomenon is again consistent with abrupt cooling during deep convection collapse (Sgubin et al., 2017; Swingedouw et al., 2021). In SAI 2080, the cooling is more drastic (Figure 5d), perhaps a result of full deep convection shutdown and a weakened AMOC. These temperature drops have a positive effect on density and thereby convection, albeit not sufficient to bring SAI 2080 densities to SAI 2020 levels (Figure 5f). Therefore, the salinity deficit of SAI 2080 with respect to SAI 2020 (Figure 5e) presents a clear obstacle to restarting deep convection.

Recognizing the importance of salinity changes, we sketch a possible mechanism behind SAI 2080's failure to spur convection. First, all scenarios see an increase in surface freshwater forcing (Figure S3 in Supporting Information S1) which contributes to a gradual salinity loss. This weakens convection and consequently the AMOC. Subsequently, weak AMOC and convection reduce salt transport into the subpolar gyre reinforcing the salinity decline (Kuhlbrodt et al., 2007). While SAI 2020 mitigates these feedbacks early on, SAI 2080 arrives only after substantial freshening. Closing the density gap via cooling then runs into 'diminishing returns': density gains are less than proportional to temperature decreases (Figure S5 in Supporting Information S1).

Another potentially important factor not included in this analysis is Greenland run-off. It likely contributes to fresher subpolar gyre conditions in SAI 2080. Additionally, Arctic outflows also supply freshwater to the deep convection regions and could vary depending on the scenario (Li et al., 2021).

4. Discussion

In our simulations, the quick drop in GMST due to abrupt SAI is contrasted by a slow ocean response. Gradual early-century SAI, on the other hand, retains an ocean state much closer to the present-day reference. Elevated OHC, weak AMOC and absent deep convection coupled with a lower GMST presents a (transient) climate state unknown from purely GHG-forced scenarios.

Note that our scenarios are extreme cases with a high signal-to-noise ratio, rather than desirable or plausible futures. More cautious protocols typically deploy SAI in tandem with emission mitigation to limit a temporary temperature overshoot (National Academies of Sciences, Engineering, and Medicine, 2021). If a cooling scenario were actually considered, a ramp-up of SAI would be more sensible than the sudden deployment in SAI 2080. Such a gradual approach would enable a fine-tuning of the injection scheme based on observations.

Besides the high forcings, our scenarios also involve a limited SAI scheme. As our implementation relies on a single degree of freedom, we can only meet a GMST target but not control other aspects of the temperature pattern. More control parameters, on the other hand, may be beneficial to prevent a interhemispheric temperature asymmetry which risks a displacement of the ITCZ (Bischoff & Schneider, 2016; Broccoli et al., 2006). Still, restoring the meridional temperature pattern in SAI 2080 would come with problems of its own: less cooling over the North Atlantic further endangers deep convection.

As for our results, a mitigating effect of SAI on AMOC decline was already known in multiple models and scenarios (MacMartin et al., 2022; Tilmes et al., 2018, 2020; Xie et al., 2022) but not in the case of late-century abrupt deployment. To our knowledge, no studies have been performed on the effect of SAI on deep convection shutdown either. Model dependencies are certain as deep convection shutdown is not a universal phenomenon in CMIP6 (Swingedouw et al., 2021). In fact, the absence of a warming hole in another SAI study using the UKESM1 model (Henry et al., 2023) could indicate a deep convection more stable than that of CESM2.

It is worth pointing out similarities between our abrupt SAI case and rapid negative emission scenarios (Schwinger et al., 2022). Removal of GHG after prolonged heating can lead to an interhemispheric temperature asymmetry if the timescale of extraction is shorter than that of the AMOC recovery. Therefore, the possibility of SAI to manage the interhemispheric temperature gradient is an advantage compared to GHG removal.

A major questions remains open: do the climates of both SAI scenarios eventually converge? This question cannot be answered without extending the simulations, which is outside the scope of this study. When extrapolating our results, the OHC difference is expected to lessen due to residual ocean warming in SAI 2020. Whether the gap fully closes may also depend on the AMOC and deep convection because of their impact on ocean heat uptake (Marshall & Zanna, 2014). As for deep convection, the aforementioned salinity deficit in SAI 2080 inhibits convergence of the SAI scenarios. Nevertheless, should some years of deep convection arise in SAI 2080 (e.g., as a result of natural variability), salt import would be strengthened, thereby improving long-term prospects of a revival.

5. Summary

In this study, we presented model results of a late-century SAI emergency intervention that aims to restore surface temperatures under simultaneous GHG forcing. By comparing our findings with a gradual early-century SAI scenario, we show that abrupt late-century SAI is less effective at mitigating changes in OHC, the AMOC and North Atlantic deep convection.

First, abrupt SAI failed to release heat trapped in deeper ocean layers. Even an early onset of SAI only mitigates but does not halt OHC accumulation. Both results are linked to slow ocean equilibration times and the target of GMST stabilization.

Second, abrupt SAI partially stabilized a weakened AMOC, albeit not halting the decline of northward heat transport. Under earlier SAI, the AMOC decline is mitigated in both, volume and heat transport. As a result, the scenarios reach drastically different AMOC states despite comparable GMST. A weaker AMOC may contribute to the observed undercooling of the northern hemisphere in the emergency intervention scenario. This, in turn, may be relevant for the choice of injection pattern.

Third, a shutdown of North Atlantic deep convection could not be reversed with rapid, SAI-induced cooling. We suspect that a weakened AMOC, absence of convective feedback, fresher surface conditions and a sub-proportional density response of water to cooling pose an obstacle for restarting deep convection. An early intervention, on the other hand, retains more salt in the North Atlantic, hence the partial stabilization of deep convection.

Our findings reveal limitations of an SAI emergency deployment: reversing ocean changes after they occur is less feasible than preventing them in the first place. In this context, proactive SAI deployments may be beneficial. Delaying climate action—this includes emission mitigation—in the hope of a later rescue through SAI will come at a price.

Data Availability Statement

Our CESM2-CAM6 SAI implementation (Pflüger, 2023b), including the input aerosol fields we used, analysis tools (Pflüger, 2023a) and the notebooks used to generate figures (Pflüger, 2023c) can be found on public GitHub

repositories. The simulation output required to create all figures is stored in a Zenodo repository (Pflüger et al., 2024). More simulation data can be made available upon reasonable request. The CMIP6 data used for comparison in Figure 3 is publicly available (Danabasoglu, 2019a, 2019b).

Acknowledgments

We thank Daniele Visioni, Doug MacMartin and Ben Kravitz for sharing their feedback controller code, our colleague Michael Kliphuis for providing his AMOC streamfunction tools and Simone Tilmes for providing CESM2-WACCM data. Furthermore, we thank Jasper de Jong and Michiel Baatsen for fruitful discussions on the feedback control of SAI 2080. Daniel Pflüger and Claudia Wieners are supported by the Dutch Ministry for Education, Culture and Science via the Sectorplan Science and Technology.

References

- Biermann, F., Oomen, J., Gupta, A., Ali, S. H., Conca, K., Hajer, M. A., et al. (2022). Solar geoengineering: The case for an international non-use agreement. *WIREs Climate Change*, 13(3), e754. <https://doi.org/10.1002/wcc.754>
- Bischoff, T., & Schneider, T. (2016). The equatorial energy balance, ITCZ position, and double-ITCZ bifurcations. *Journal of Climate*, 29(8), 2997–3013. <https://doi.org/10.1175/JCLI-D-15-0328.1>
- Broccoli, A. J., Dahl, K. A., & Stouffer, R. J. (2006). Response of the ITCZ to northern hemisphere cooling. *Geophysical Research Letters*, 33(1), L01702. <https://doi.org/10.1029/2005GL024546>
- Cheng, L., Von Schuckmann, K., Abraham, J. P., Trenberth, K. E., Mann, M. E., Zanna, L., et al. (2022). Past and future ocean warming. *Nature Reviews Earth & Environment*, 3(11), 776–794. <https://doi.org/10.1038/s43017-022-00345-1>
- Church, J. A., White, N. J., Domingues, C. M., Monselesan, D. P., & Miles, E. R. (2013). Sea-level and ocean heat-content change. In *International geophysics* (Vol. 103, pp. 697–725). Elsevier. <https://doi.org/10.1016/B978-0-12-391851-2.00027-1>
- Danabasoglu, G. (2019a). NCAR CESM2 model output prepared for CMIP6 ScenarioMIP ssp370 [Dataset]. Earth System Grid Federation. <https://doi.org/10.22033/ESGF/CMIP6.7753>
- Danabasoglu, G. (2019b). NCAR CESM2 model output prepared for CMIP6 ScenarioMIP ssp585 [Dataset]. Earth System Grid Federation. <https://doi.org/10.22033/ESGF/CMIP6.7768>
- Danabasoglu, G., Lamarque, J., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., et al. (2020). The community earth system model version 2 (CESM2). *Journal of Advances in Modeling Earth Systems*, 12, 35. <https://doi.org/10.1029/2019MS001916>
- Drijfhout, S., van Oldenborgh, G. J., & Cimatoribus, A. (2012). Is a decline of AMOC causing the warming hole above the North Atlantic in observed and modeled warming patterns? *Journal of Climate*, 25(24), 8373–8379. <https://doi.org/10.1175/JCLI-D-12-00490.1>
- Fasullo, J. T., Tilmes, S., Richter, J. H., Kravitz, B., MacMartin, D. G., Mills, M. J., & Simpson, I. R. (2018). Persistent polar ocean warming in a strategically geoengineered climate. *Nature Geoscience*, 11(12), 910–914. <https://doi.org/10.1038/s41561-018-0249-7>
- Henry, M., Haywood, J., Jones, A., Dalvi, M., Wells, A., Visioni, D., et al. (2023). Comparison of UKESM1 and CESM2 simulations using the same multi-target stratospheric aerosol injection strategy. *Atmospheric Chemistry and Physics*, 23(20), 13369–13385. <https://doi.org/10.5194/acp-23-13369-2023>
- IPCC. (2022a). *Climate change 2022: Impacts, adaptation and vulnerability*. Cambridge University Press. <https://doi.org/10.1017/9781009325844>
- IPCC. (2022b). *Climate change 2022: Mitigation of climate change*. Cambridge University Press.
- Irvine, P. J., Kravitz, B., Lawrence, M. G., & Muri, H. (2016). An overview of the earth system science of solar geoengineering. *WIREs Climate Change*, 7(6), 815–833. <https://doi.org/10.1002/wcc.423>
- Kleinschmitt, C., Boucher, O., & Platt, U. (2018). Sensitivity of the radiative forcing by stratospheric sulfur geoengineering to the amount and strategy of the SO₂ injection studied with the LMDZ-S3A model. *Atmospheric Chemistry and Physics*, 18(4), 2769–2786. <https://doi.org/10.5194/acp-18-2769-2018>
- Kravitz, B., MacMartin, D. G., Mills, M. J., Richter, J. H., Tilmes, S., Lamarque, J., et al. (2017). First simulations of designing stratospheric sulfate aerosol geoengineering to meet multiple simultaneous climate objectives. *Journal of Geophysical Research: Atmospheres*, 122(23), 12616. <https://doi.org/10.1002/2017JD026874>
- Kravitz, B., MacMartin, D. G., Wang, H., & Rasch, P. J. (2016). Geoengineering as a design problem. *Earth System Dynamics*, 7(2), 469–497. <https://doi.org/10.5194/esd-7-469-2016>
- Kuhlbrodt, T., Griesel, A., Montoya, M., Levermann, A., Hofmann, M., & Rahmstorf, S. (2007). On the driving processes of the Atlantic meridional overturning circulation. *Reviews of Geophysics*, 45(2), RG2001. <https://doi.org/10.1029/2004rg000166>
- Li, F., Lozier, M. S., Holliday, N. P., Johns, W. E., Le Bras, I. A., Moat, B. I., et al. (2021). Observation-based estimates of heat and freshwater exchanges from the subtropical North Atlantic to the arctic. *Progress in Oceanography*, 197, 102640. <https://doi.org/10.1016/j.poccean.2021.102640>
- Liu, W., Fedorov, A. V., Xie, S.-P., & Hu, S. (2020). Climate impacts of a weakened Atlantic meridional overturning circulation in a warming climate. *Science Advances*, 6(26), eaaz4876. <https://doi.org/10.1126/sciadv.aaz4876>
- Lockley, A., Xu, Y., Tilmes, S., Sugiyama, M., Rothman, D., & Hinds, A. (2022). 18 politically relevant solar geoengineering scenarios. *Socio-Environmental Systems Modelling*, 4, 18127. <https://doi.org/10.18174/sesmo.18127>
- Long, J. C. S., & Shepherd, J. G. (2014). The strategic value of geoengineering research. In B. Freedman (Ed.), *Global environmental change* (pp. 757–770). Springer Netherlands. https://doi.org/10.1007/978-94-007-5784-4_24
- MacMartin, D. G., Kravitz, B., Tilmes, S., Richter, J. H., Mills, M. J., Lamarque, J., et al. (2017). The climate response to stratospheric aerosol geoengineering can be tailored using multiple injection locations. *Journal of Geophysical Research: Atmospheres*, 122(23), 12616. <https://doi.org/10.1002/2017JD026868>
- MacMartin, D. G., Visioni, D., Kravitz, B., Richter, J., Felgenhauer, T., Lee, W. R., et al. (2022). Scenarios for modeling solar radiation modification. *Proceedings of the National Academy of Sciences*, 119(33), e2202230119. <https://doi.org/10.1073/pnas.2202230119>
- Marshall, D. P., & Zanna, L. (2014). A conceptual model of ocean heat uptake under climate change. *Journal of Climate*, 27(22), 8444–8465. <https://doi.org/10.1175/JCLI-D-13-00344.1>
- Menary, M. B., & Wood, R. A. (2018). An anatomy of the projected North Atlantic warming hole in CMIP5 models. *Climate Dynamics*, 50(7–8), 3063–3080. <https://doi.org/10.1007/s00382-017-3793-8>
- National Academies of Sciences, Engineering, and Medicine. (2021). *Reflecting sunlight: Recommendations for solar geoengineering research and research governance*. National Academies Press. <https://doi.org/10.17226/25762>
- Oomen, J. (2021). *Imagining climate engineering: Dreaming of the designer climate*. Routledge, Taylor & Francis Group.
- Pflüger, D. (2023a). *cesm_tools* [Software]. Github. Retrieved from https://github.com/daniel-pflueger/cesm_tools
- Pflüger, D. (2023b). *imau_cesm2_prescribed_aerosols* [Software]. Github. Retrieved from https://github.com/daniel-pflueger/imau_cesm2_prescribed_aerosols
- Pflüger, D. (2023c). *sai_ocean_figures_2023* (Notebook). Retrieved from https://github.com/daniel-pflueger/sai_ocean_figures_2023

- Pflüger, D., Wieners, C., van Kampenhout, L., Wijngaard, R., & Dijkstra, H. (2024). Flawed emergency intervention: Slow ocean response to abrupt stratospheric aerosol injection (Dataset) [Dataset]. Zenodo. <https://doi.org/10.5281/zenodo.10606102>
- Plazzotta, M., Séférian, R., Douville, H., Kravitz, B., & Tjiputra, J. (2018). Land surface cooling induced by sulfate geoengineering constrained by major volcanic eruptions. *Geophysical Research Letters*, *45*(11), 5663–5671. <https://doi.org/10.1029/2018GL077583>
- Reynolds, J. L. (2019). Solar geoengineering to reduce climate change: A review of governance proposals. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, *475*(2229), 20190255. <https://doi.org/10.1098/rspa.2019.0255>
- Schwinger, J., Asaadi, A., Goris, N., & Lee, H. (2022). Possibility for strong northern hemisphere high-latitude cooling under negative emissions. *Nature Communications*, *13*(1), 1095. <https://doi.org/10.1038/s41467-022-28573-5>
- Sgubin, G., Swingedouw, D., Drijfhout, S., Mary, Y., & Bennabi, A. (2017). Abrupt cooling over the North Atlantic in modern climate models. *Nature Communications*, *8*(1), 14375. <https://doi.org/10.1038/ncomms14375>
- Smith, W. (2020). The cost of stratospheric aerosol injection through 2100. *Environmental Research Letters*, *15*(11), 114004. <https://doi.org/10.1088/1748-9326/aba7e7>
- Stocker, T. F. (1998). The seesaw effect. *Science*, *282*(5386), 61–62. <https://doi.org/10.1126/science.282.5386.61>
- Svoboda, T. (2017). *The ethics of climate engineering: Solar radiation management and non-ideal justice*. Routledge, Taylor & Francis Group.
- Swingedouw, D., Bily, A., Esquerdo, C., Borchert, L. F., Sgubin, G., Mignot, J., & Menary, M. (2021). On the risk of abrupt changes in the North Atlantic subpolar gyre in CMIP6 models. *Annals of the New York Academy of Sciences*, *1504*(1), 187–201. <https://doi.org/10.1111/nyas.14659>
- Tilmes, S., MacMartin, D. E., Lenaerts, J. T. M., Van Kampenhout, L., Muntjewerf, L., Xia, L., et al. (2020). Reaching 1.5°C and 2.0°C global surface temperature targets using stratospheric aerosol geoengineering. *Earth System Dynamics*, *11*(3), 579–601. <https://doi.org/10.5194/esd-11-579-2020>
- Tilmes, S., Richter, J. H., Kravitz, B., MacMartin, D. G., Mills, M. J., Simpson, I. R., et al. (2018). CESM1(WACCM) stratospheric aerosol geoengineering large ensemble project. *Bulletin of the American Meteorological Society*, *99*(11), 2361–2371. <https://doi.org/10.1175/BAMS-D-17-0267.1>
- Visioni, D., MacMartin, D. G., Kravitz, B., Boucher, O., Jones, A., Lurton, T., et al. (2021). Identifying the sources of uncertainty in climate model simulations of solar radiation modification with the G6sulfur and G6solar geoengineering model intercomparison project (GeoMIP) simulations (p. 21). <https://doi.org/10.5194/acp-21-10039-2021>
- Weijer, W., Cheng, W., Garuba, O. A., Hu, A., & Nadiga, B. T. (2020). CMIP6 models predict significant 21st century decline of the Atlantic meridional overturning circulation. *Geophysical Research Letters*, *47*(12), 502. <https://doi.org/10.1029/2019GL086075>
- Wieners, C. E., Hofbauer, B. P., De Vries, I. E., Honegger, M., Visioni, D., Russchenberg, H. W. J., & Felgenhauer, T. (2023). Solar radiation modification is risky, but so is rejecting it: A call for balanced research. *Oxford Open Climate Change*, *3*(1), kgad002. <https://doi.org/10.1093/oxfclm/kgad002>
- Xie, M., Moore, J. C., Zhao, L., Wolovick, M., & Muri, H. (2022). Impacts of three types of solar geoengineering on the Atlantic meridional overturning circulation. *Atmospheric Chemistry and Physics*, *22*(7), 4581–4597. <https://doi.org/10.5194/acp-22-4581-2022>
- Zarnetske, P. L., Gurevitch, J., Franklin, J., Groffman, P. M., Harrison, C. S., Hellmann, J. J., et al. (2021). Potential ecological impacts of climate intervention by reflecting sunlight to cool earth. *Proceedings of the National Academy of Sciences*, *118*(15), e1921854118. <https://doi.org/10.1073/pnas.1921854118>

References From the Supporting Information

- Astrom, K. J., & Rundqwist, L. (1989). Integrator windup and how to avoid it. In *1989 American control conference*.
- Hansen, J., Sato, M., Ruedy, R., Nazarenko, L., Lacis, A., Schmidt, G., et al. (2005). Efficacy of climate forcings. *Journal of Geophysical Research*, *110*(D18), D18104. <https://doi.org/10.1029/2005jd005776>
- Lawrence, D., Fisher, R., Koven, C., Oleson, K., Swenson, S., Vertenstein, M., et al. (2018). *Technical description of version 5.0 of the community land model (CLM) (Technical Report)*. National Center for Atmospheric Research. Retrieved from <https://www2.cesm.ucar.edu/models/cesm2/land/CLM50TechNote.pdf>
- Visioni, D., MacMartin, D. G., Kravitz, B., Lee, W., Simpson, I. R., & Richter, J. H. (2020). Reduced poleward transport due to stratospheric heating under stratospheric aerosols geoengineering. *Geophysical Research Letters*, *47*(17), 5553. <https://doi.org/10.1029/2020GL089470>