



© The Author(s), 2020. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike licence (<http://creativecommons.org/licenses/by-nc-sa/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the same Creative Commons licence is included and the original work is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use.  
First published online 28 September 2020

*Nutrition Society Live 2020 was held virtually on 14–15 July 2020*

## Symposium two: Novel methods for assessing protein metabolism

### Monitoring food digestion with magnetic resonance techniques

Paul A. M. Smeets<sup>1,2\*</sup>, Ruoxuan Deng<sup>1,3</sup>, Elise J. M. van Eijnatten<sup>1</sup> and Morwarid Mayar<sup>1,4</sup>

<sup>1</sup>Division of Human Nutrition and Health, Wageningen University, Stippeneng 4, 6708 WE, Wageningen, The Netherlands

<sup>2</sup>Image Sciences Institute, University Medical Center Utrecht, Utrecht University, Heidelberglaan 100, 3584 CX, Utrecht, The Netherlands

<sup>3</sup>Laboratory of Food Process Engineering, Wageningen University, Bornse Weiland 9, 6708 WE, Wageningen, The Netherlands

<sup>4</sup>Laboratory of Biophysics, Wageningen University, Stippeneng 4, 6708 WE, Wageningen, The Netherlands

This review outlines the current use of magnetic resonance (MR) techniques to study digestion and highlights their potential for providing markers of digestive processes such as texture changes and nutrient breakdown. *In vivo* digestion research can be challenging due to practical constraints and biological complexity. Therefore, digestion is primarily studied using *in vitro* models. These would benefit from further *in vivo* validation. NMR is widely used to characterise food systems. MRI is a related technique that can be used to study both *in vitro* model systems and *in vivo* gastro-intestinal processes. MRI allows visualisation and quantification of gastric processes such as gastric emptying and coagulation. Both MRI and NMR scan sequences can be configured to be sensitive to different aspects of gastric or intestinal contents. For example, magnetisation transfer and chemical exchange saturation transfer can detect proton (<sup>1</sup>H) exchange between water and proteins. MRI techniques have the potential to provide molecular-level and quantitative information on *in vivo* gastric (protein) digestion. This requires careful validation in order to understand what these MR markers of digestion mean in a specific digestion context. Combined with other measures they can be used to validate and inform *in vitro* digestion models. This may bridge the gap between *in vitro* and *in vivo* digestion research and can aid the optimisation of food properties for different applications in health and disease.

#### Digestion: Gastric emptying: MRI: Protein

The process of digestion is necessary for acquiring nutrients from the foods we ingest. Digestion encompasses a series of complex physiological, mechanical and biochemical processing steps that lead to the mechanical and biochemical breakdown of food structures which ultimately allows absorption and utilisation of nutrients<sup>(1)</sup>. Both independent and interrelated processes at

multiple length scales are involved in food breakdown, mixing and absorption<sup>(2)</sup>. Briefly, anticipation of food intake triggers several anticipatory physiological responses such as increased salivation and production of gastric juice that prepare the body for the influx of nutrients<sup>(3)</sup>. The first phase of digestion is the oral phase (ingestion). During oral processing, mastication

**Abbreviations:** CEST, chemical exchange saturation transfer; GE, gastric emptying; MR, magnetic resonance; MT, magnetisation transfer; RF, radiofrequency.

\*Corresponding author: Paul A. M. Smeets, email [paul.smeets@wur.nl](mailto:paul.smeets@wur.nl)



and secretion of saliva lead to the formation of a food bolus that can be swallowed safely<sup>(4)</sup>. These oral processes are not only important for digestion, but also for sensory perception of foods, eating enjoyment and satiation<sup>(5)</sup>. The second phase is gastric digestion, during which the food mass in the stomach is prepared for further digestion and absorption in the intestines. Gastric digestion involves mixing and addition of hydrochloric acid along with pepsin (a protease) and gastric lipase<sup>(6)</sup>. The third phase is intestinal digestion; from the stomach food passes through the pyloric valve into the small intestine where pancreatic proteases, lipases and amylase are added along with bile and the resulting chyme is mixed. The chyme is passed along the small intestine where most of the nutrients are absorbed. The small intestine comprises the duodenum, jejunum and ileum and chyme proceeds from one to the other until it passes through the ileocaecal valve into the large intestine. There, some of the remaining undigested food such as dietary fibre is fermented by bacteria into absorbable compounds such as SCFA, and most of the remaining water is removed before defecation<sup>(1)</sup>. Across and within these different processing stages there are numerous physiological signals (neural and hormonal) that feed forward and backward, presumably to optimise digestion.

*In vivo* digestion research can be challenging due to practical constraints, biological complexity and ethical obstacles. Notably, classic techniques to study digestion *in vivo* are mostly invasive and involve for example taking gastric aspirates through a nasogastric tube or monitoring gastric pressure or pH with sensors. Therefore, *in vitro* model systems are widely used to study digestive processes under controlled and simplified conditions<sup>(7)</sup>. This provides detailed information on the effects of enzymatic processes on the physical and chemical characteristics of food structures during digestion<sup>(8,9)</sup>. Although validation and refinement of *in vitro* models using *in vivo* data is an ongoing collaborative effort in the field (see <https://www.cost-infogest.eu/> and Bohn *et al.*<sup>(10)</sup>), bridging the significant gap between *in vitro* model systems and the complexity of *in vivo* digestion remains a challenge. An emerging approach that could help address this in part involves the use of ingestible devices that can take samples or measurements as they pass through the gastrointestinal tract<sup>(11,12)</sup>. The core idea that will be explored in the present paper is that magnetic resonance (MR) techniques may be used to bridge this gap because they can be used to monitor relevant digestive processes both *in vitro* and *in vivo*.


NMR provides information on the state of water protons in foods and has been widely used as a characterisation and process quality control tool in different food systems<sup>(13,14)</sup>. It can be performed at relatively low magnetic field strengths (about 0.5 T) and is used for measuring *in vitro* samples of digesta or gastric aspirates. It has the advantage of low cost and ease of operation. MRI is a commonly used related technique that, among numerous other applications e.g. in medicine, can be used to perform both *in vitro* and *in vivo* imaging measurements<sup>(13)</sup> non-invasively. It is most commonly performed at 1.5 or 3 T and is more expensive than NMR. NMR and

MRI share the same underlying principles and use magnetisation in combination with radiofrequency (RF) pulses to obtain RF signals from nuclei of interest, usually water protons (<sup>1</sup>H) due to their natural abundance and sensitivity. Briefly, protons spinning in a magnetic field are 'excited' with a targeted RF pulse. During their subsequent 'relaxation' back to their equilibrium state they emit RF, which is measured with a coil (antenna). MRI can provide information not only on the volume of gastric content fractions, but also on intra-gastric processes such as phase separation and clot formation, on gallbladder responses<sup>(15)</sup>, and on intestinal parameters such as intestinal motility and small bowel water content (for overviews see Marciani *et al.*<sup>(16)</sup> and Spiller and Marciani<sup>(17)</sup>). We argue that by virtue of this common ground NMR measurements of *in vitro* samples or gastric aspirates can be used to aid the interpretation of substance-specific MR characteristics such as signal relaxation rates in a digestion context. This may serve to inform and validate MRI measurements of the same *in vitro* system as well as an *in vivo* equivalent, which in turn can validate and inform *in vitro* digestion models (Fig. 1). Thus, this review outlines the current use of MR techniques to study digestion and highlights their potential for providing markers of digestive processes such as gastric coagulation and nutrient breakdown and how MR techniques in combination with other measures may bridge the gap between *in vitro* and *in vivo* research. Collectively, insights from such interdisciplinary studies can foster the optimisation of food properties for different applications in health and disease.

### *In vitro* digestion models

To study digestion, various *in vitro* digestion models that can be used to mimic one or more digestion phases, such as gastric or intestinal digestion, have been developed<sup>(18)</sup>. These models can also be applied to study absorption of the digested material, by incorporating intestinal cell cultures<sup>(19)</sup>. They vary from simple static models to highly sophisticated dynamic, computer-controlled gastrointestinal models<sup>(20)</sup>. In static models, the digestive fluid and food materials are constant, and hence they are convenient for investigating the mechanisms of mass transport and structure breakdown<sup>(7,21,22)</sup>. Dynamic models, such as the Tim models<sup>(23)</sup> and the SHIME model<sup>(24)</sup>, include factors such as gastric emptying<sup>(25)</sup> (GE) and inflow of gastric and intestinal juice. Therefore, such models are more physiologically accurate than static models.

With the use of *in vitro* models, multiple approaches can be combined to determine the progression of digestion by e.g. (bio-)chemical and physical analysis of digesta samples. Several chemical analysis approaches have been applied to measure food hydrolysis during digestion. For example, examining changes in the size of peptides or amount of free amino groups for protein digestion<sup>(9,26)</sup>, glucose for starch digestion<sup>(27)</sup> and NEFA for fat digestion<sup>(28)</sup>. From a physical perspective, rheology or texture analysis, sometimes combined with microscopy, are used to measure changes in physical



**Fig. 1.** (Colour online) Overview of the proposed interdisciplinary approach to study digestion by employing magnetic resonance (MR) techniques in combination with a variety of other measurements.

properties such as viscosity, or in the structure of food particles (from macro to micro) during digestion<sup>(29–31)</sup>. Recently, several other approaches have been used to study digestion, e.g. hyperspectral imaging to monitor the mass transfer between digestive fluid and food particles<sup>(32)</sup> and MR techniques to monitor the hydrolysis of nutrients and changes in food composition (Bordoni *et al.*<sup>(33)</sup>; Deng *et al.*<sup>(26)</sup>).

The advantages of using *in vitro* models include easy sampling, well controlled and reproducible conditions, ability to assess chemical processes in detail, and the absence of ethical restrictions. In addition, such simplified systems also make interpretation easier (within model boundaries) and multiple follow-up experiments can be readily done to further unravel observed phenomena. However, the validation of *in vitro* models remains a big challenge due to the inherent simplifications such as the absence of feedback mechanisms. For instance, secretion of digestive juices in response to a meal *in vivo* is regulated by the autonomic nervous system and several hormones, which is extremely challenging to replicate within *in vitro* models<sup>(8)</sup>. To aid the validation of *in vitro* models, it is of interest to investigate the potential of non-invasive approaches for *in vivo* monitoring of digestion. Several promising MR techniques are described in the following sections.

### NMR spectroscopy

NMR spectroscopy is a non-destructive technique often used in the fields of biology, chemistry and food

technology to determine the molecular structure and to quantify the concentration of molecules in a sample. With NMR the interactions between an external magnetic field and atomic nuclei that have a magnetic property can be observed. The main nucleus of interest in NMR is a proton ( $^1\text{H}$ ). Examples of other commonly utilised nuclei in NMR are  $^{13}\text{C}$ ,  $^{31}\text{P}$  and  $^{15}\text{N}$ . In a typical NMR measurement, an NMR spectrum is recorded in which the position of the peaks can provide information on the molecular structure of the compounds present in the sample. Furthermore, the area under the curve of NMR peaks is proportional to the number of nuclei giving rise to the peak<sup>(14,34)</sup>. Therefore, it is possible to estimate (changes in) the molar percentage or concentration of several components. The combination of molecular level and quantitative information that can be obtained with NMR makes it a promising technique for examining food digestion *in vitro*. Specifically, it may be used to monitor macronutrient hydrolysis. For example, NMR has been used to quantify the products of lipid hydrolysis, such as diglycerides and fatty acids, in complex lipid mixtures<sup>(35)</sup> and in foods, such as fish<sup>(36,37)</sup> and sunflower oil<sup>(38)</sup> during *in vitro* digestion. In addition to lipid hydrolysis studies, NMR has also been applied to study protein hydrolysis *in vitro*. Sundekilde *et al.*<sup>(39)</sup> used NMR for monitoring enzyme-assisted hydrolysis of animal proteins under real-time conditions directly in the NMR spectrometer. This approach enabled the monitoring of free amino acids produced during enzymatic hydrolysis. Bordoni *et al.*<sup>(33)</sup> used NMR to monitor digestion of cheese in an *in vitro* digestion model simulating digestion in the mouth, stomach and small intestine.

Also, NMR spectroscopy is capable of providing information on the kinetics of carbohydrate hydrolysis under acidic conditions<sup>(40,41)</sup>. While these applications show that NMR spectroscopy is a powerful tool for monitoring food digestion *in vitro*, it is less suitable for *in vivo* studies. NMR spectra from *in vivo* samples will be more complex and more difficult to interpret. Moreover, NMR only allows the detection of molecules that are present in sufficiently high concentrations, which may not be the case *in vivo*.

### NMR relaxometry

In addition to NMR spectroscopy, there are several other MR techniques that could be used for both *in vitro* and *in vivo* monitoring of different aspects of food digestion. These include measurement of  $T_1$  and  $T_2$  relaxation times and chemical exchange markers with NMR or MRI (Fig. 1).


#### $T_1$ and $T_2$ relaxation times

$T_1$  and  $T_2$  relaxation times reflect how protons in a magnetic field relax back to their equilibrium position after excitation by an RF pulse. The main applications of  $T_1$  and  $T_2$  are based on the investigation of the relaxation behaviour of water protons in different environments<sup>(42)</sup>. Water proton relaxation is mainly determined by their mobility and is affected by macromolecular composition and structure.  $T_1$  and  $T_2$  measurements have been used to study various food properties such as moisture content, food structure and macromolecule concentration<sup>(43)</sup>. For instance, Ziegler *et al.*<sup>(44)</sup> used  $T_1$  measurements to predict water migration in starch-pectin gels during drying since  $T_1$  decreases with the decrease of their moisture content.  $T_2$  was used to predict water-holding capacity of whey protein particles; a higher water-holding capacity is associated with a longer  $T_2$ . Similarly,  $T_2$  has been used to study the swelling of hydrogels<sup>(45,46)</sup>.  $T_2$  has also been used to study the local structure of cheese; due to the inhomogeneity of the cheese, three distinct  $T_2$  relaxation components could be identified reflecting serum water (the water accumulated in the protein network), the water inside meshes of the casein gel-like network, and the water trapped within the casein matrix<sup>(47)</sup>. In addition,  $T_2$  can be used to determine the protein concentration in casein solutions; with increasing concentration the  $T_2$  decreases<sup>(48)</sup>. These examples show that  $T_1$  and  $T_2$  can be used to monitor changes in water migration, food structure and the composition of food and digestive juice that take place during digestion<sup>(6)</sup>. Despite this,  $T_1$  and  $T_2$  measurements have only been applied in a limited number of digestion studies. For instance,  $T_2$  has been shown to be useful in detecting penetration of digestion fluid into the food matrix during *in vitro* digestion<sup>(33,49)</sup>. Another study showed a linear association between viscosity of locust bean gum meal and  $T_2$  *in vitro*, and highlighted the possible application of  $T_2$  to monitor changes in meal viscosity in the gastric lumen *in vivo* with the use of MRI<sup>(50)</sup>. Moreover, in

our recent study we show that the hydrolysis of protein during *in vitro* gastric digestion can be monitored by  $T_2$ ;  $T_2$  was associated with protein released from food particles into the surrounding liquid<sup>(51)</sup>. However, *in vivo* gastric digestion is more complicated than the static *in vitro* model used here, which e.g. does not take into account dynamic processes such as the production of gastric juice and GE. These processes will introduce changes in the system that have multiple effects such as pH changes and dilution, in addition to the hydrolysis of nutrients. Because  $T_1$  and  $T_2$  are affected by many such factors, careful validation is needed to be able to interpret changes in  $T_1$  and  $T_2$  in different digestion contexts. This requires further investigation under dynamic circumstances, both *in vitro* and *in vivo* and the combination of NMR and MRI  $T_1$  and  $T_2$  measurements. For example, as shown in Fig. 2, the NMR  $T_2$  spectrum shows separate peaks that represent the protein gel and the simulated gastric fluid around it. This information contributes to the interpretation of the MRI  $T_2$  maps.

#### $T_{1\rho}$

Another relaxation time of interest in MR is the  $T_{1\rho}$  or the  $T_1$  relaxation time in the rotating frame.  $T_{1\rho}$  is useful to study low-frequency motion processes and chemical exchange in biological tissues. It is measured by applying an additional RF pulse after the excitation pulse to lock the magnetisation in the rotating frame. The time it takes for the locked magnetisation to decay to zero is the  $T_{1\rho}$  relaxation time<sup>(52)</sup>. In addition to conventional relaxation time measurements,  $T_2$  and  $T_{1\rho}$  relaxation time dispersion measurements<sup>(53)</sup> are promising MR markers for monitoring digestion because they can be used to examine macromolecules in solution or the interaction between bulk water protons and exchangeable macromolecule protons in semi-solids. In relaxation time dispersion measurements, the relaxation time is measured under varying measurement conditions. In the presence of chemical exchange, for example proton exchange between a macromolecule and bulk water, a dispersion of the relaxation time under those varying conditions can be observed. The extent of dispersion depends on the rate of chemical exchange. Compared to conventional relaxation time measurements, relaxation dispersion is more quantitative, since the experimental data can be fitted with theoretical models of two- or three-site exchange from which the exchange rate can be extracted<sup>(52,54,55)</sup>. This exchange rate depends on the state of a macromolecule, e.g. intact or digested, and hence, can potentially be related to the kinetics of digestion. However, to date the application of  $T_2$  dispersion has been limited to investigating molecular dynamics of proteins *in vitro*<sup>(56)</sup>.  $T_{1\rho}$  dispersion, in contrast, has been more commonly applied in *in vivo* MRI studies, but not yet in the domain of digestion. Duvvuri *et al.*<sup>(57)</sup> suggested that in cartilage proton exchange between protons from NH and OH groups in the proteoglycans and water dominate the  $T_{1\rho}$  dispersion of water. They showed that the exchange rates increase with proteoglycan breakdown. This suggests that  $T_{1\rho}$



**Fig. 2.** (Colour online) Illustration of how NMR and MRI of the same *in vitro* model can be used to study the dependence of magnetic resonance parameters on nutrient breakdown. Shown here is the increase in  $T_2$  relaxation rate ( $R_2 = 1/T_2$ ), measured with either technique, and the protein fraction in simulated gastric juice (SGF) during digestion of whey protein gel pieces. Adapted from Deng *et al.*<sup>(51)</sup>.

dispersion has potential as a marker for *in vivo* monitoring of protein digestion.

In conclusion, MR relaxation parameters and their meaning in a digestion context need to be further elucidated. NMR can aid the interpretation of *in vitro* and *in vivo* relaxometry measurements with MRI. In turn, *in vivo* MRI can serve to validate and inform *in vitro* models.

### NMR cross-relaxation

#### Magnetisation transfer

Magnetisation transfer (MT) is an MR technique that is used to create a contrast between tissues in which protons are present in three different states: (i) in free water, (ii) bound to semi-solid macromolecules and (iii) as water in the hydration layer between the macromolecules and free water. In an MT measurement, the magnetisation of macromolecular protons is saturated by application of an RF pulse. The saturation is then transferred to protons in water through proton exchange resulting in a decrease in the water signal intensity. The MT rate is sensitive to the formation of a semi-solid structure<sup>(58)</sup> and hence could serve as a marker for the degree of coagulation of proteins during gastric digestion. The degree of coagulation can affect digestion and GE rate<sup>(59)</sup>.

#### Chemical exchange saturation transfer

Chemical exchange saturation transfer (CEST) is a relatively novel MR technique. Its principle is similar to that

of MT where saturated protons are exchanged with non-saturated protons in water. However, the main difference is that in CEST, saturation transfer takes place between a (macro)molecule in solution and water and that the saturation is frequency-selective<sup>(60)</sup>. Dona *et al.*<sup>(41)</sup> showed that CEST can be used to monitor the *in vitro* enzymatic degradation of macromolecular starch granules. Moreover, from their CEST measurement, the kinetics of glucose release during the enzymatic hydrolysis of cooked starch was successfully monitored, demonstrating the potential of CEST for obtaining quantitative information on food digestion<sup>(41)</sup>. Longo *et al.*<sup>(61)</sup> used CEST to monitor the aggregation of bovine serum albumin during heat treatment and its subsequent hydrolysis by a protease *in vitro*. During heat treatment the proteins aggregate, thereby decreasing the accessibility of the protein protons for exchange with water protons. However, after digestion of the heat-treated protein, the protons become accessible for exchange again, resulting in an increase in saturation transfer. This suggests that CEST can serve as a marker for monitoring protein digestion and how this is affected by heat treatments. Another interesting application of CEST is pH imaging in which the endogenous amide proton transfer rate is related to the pH in the tissue of interest<sup>(62)</sup>. Most applications of CEST pH imaging are done in the brain<sup>(63–65)</sup> where the pH is between 6.5 and 8.5. It would be of great utility if CEST could be used to make 3-D stomach pH maps since pH is an important factor that among others influences pepsin activity. However, stomach pH is

between 1.5 and 3.5 in children or adults and between 3.5 and 5.5 in infants. This renders stomach pH imaging challenging due to the slower amide proton transfer rate at low pH.

#### *Magnetic resonance relaxometry and cross-relaxation outlook*

While NMR/MRI relaxometry and cross-relaxation techniques have not been applied much in food digestion research, the applications to date suggest that they have potential for monitoring digestion. For *in vitro* studies, it is possible to obtain quantitative information, such as the proton exchange rate from relaxation time dispersion, MT or CEST data by fitting with biophysical models of two- or three-site chemical exchange. The chemical exchange rate is expected to depend on the pH, macromolecular concentration and semi-solid fraction, which are all factors that change during digestion. However, reliable modelling of such data can only be done on sufficiently large data sets. The collection of such data sets is time-consuming and limited to *in vitro* studies. The main challenge lies in optimising experimental conditions such as the measurement time, size of datasets and sub-set of fitting parameters to enable quantitative application of the same measurements *in vivo*. Moreover, *in vivo* data acquisition has practical constraints and interpretation is harder. There are limits to the time that volunteers can spend in the scanner and scans have to be made during breath hold or with respiratory triggering to minimise image artefacts caused by breathing movements. Physiological noise and constraints on acquisition times will result in poorer data quality compared to *in vitro* data. However, these are common challenges for any *in vivo* application of MRI, and there is continuing technical development aimed at ameliorating these issues. Moreover, as pointed out before, careful *in vitro* validation experiments can be used in the development and validation of these promising MR markers of digestion.

### MRI

MRI is a popular medical imaging technique because it does not use ionising radiation and is extremely versatile. Similar to in NMR, in MRI a strong magnetic field (1.5–7 T) is used to line up water protons ( $^1\text{H}$ ) in the body. These protons are energised by exposing them to specific RF pulses. When they relax back into their lower-energy state RF is emitted. This RF signal (echo) is measured with a coil (antenna). By varying the local magnetic field, RF pulse characteristics and the timing of RF measurement it is possible to reconstruct different types of images from the measured RF signals<sup>(66)</sup>. Among other factors, the contrast of these images depends on the local  $T_1$  and  $T_2$  relaxation rates, which vary between tissue types. The anatomical detail provided by MRI allows accurate visualisation of the stomach and its contents<sup>(67)</sup>. Also, because MRI scan sequences can be configured to be sensitive to different aspects of gastric or intestinal contents, MRI is suitable for investigating

complex meals and intragastric processes such as gastric sieving, phase separation and coagulation (see later). Moreover, MRI can be used to examine model systems, such as *in vitro* digestion models, as well as human subjects using the same scan sequences (Fig. 1). This makes it an excellent technique to bridge *in vitro* and *in vivo* research.

#### ***In vivo* MRI: gastric contents**

Since the 1990s, gastric digestion research has embraced MRI as a method to measure GE, i.e. the change in gastric content volume over time in human subjects<sup>(68,69)</sup>. Understanding GE is important since it plays a major role in digestion, satiety and nutrient absorption. GE is a rate-limiting step in the delivery of nutrients to the small intestine for further break-down, and one of the factors influencing the susceptibility to maldigestion<sup>(22)</sup>. GE rate is largely determined by the chemical characteristics of food, mainly the macronutrient content, but also physical characteristics, such as the viscosity<sup>(70,71)</sup>. There are various other techniques to assess GE rate, but the most common approaches are either indirect (C-isotope breath analysis and paracetamol absorption test), or involve the use of ionising radiation (gamma scintigraphy<sup>(72)</sup>). Advantages of MRI are that it is well suited for individual GE assessment and less dependent on the food matrix than indirect tracer-based methods such as C-isotope breath analysis<sup>(73)</sup>. Quantification of postprandial volume changes with MRI has been shown to have low inter-observer variability, unless the stomach is nearly empty<sup>(74)</sup>.

GE can be different for different food fractions. So-called, ‘gastric sieving’ happens when a meal consists of multiple textures, or when these arise as part of digestion; the (more) liquid component has been shown to quickly ‘sieve’ through the stomach while the more solid part of the meal is retained<sup>(75–77)</sup>. When the two components are blended into a (nutrient-rich) liquid, satiety is enhanced because the fast entry of energy into the duodenum slows down GE<sup>(77)</sup>. Similar sieving has been shown to occur for water and a meal shake; water can drain from the stomach while a layer of energetic liquid is retained<sup>(78)</sup>. For such foods and mechanisms tracer-based methods are not very suitable, since different tracers with different kinetics would be required to show the transit of watery and fatty components separately (see e.g. Collins *et al.*<sup>(75)</sup> for a double isotope approach), depending on their solubility.

MRI can also show air or gas volumes in the stomach, which can significantly affect perceived fullness and appetite through providing gastric distention. For example, aerated milk-based drinks (foams) increased gastric volume and reduced hunger more than an iso-energetic liquid control drink<sup>(79)</sup>. This underscores the notion that stretching of the stomach wall contributes to satiation and perceived fullness. In a well-controlled study which used MRI in combination with measurement of gastric pressure Kwiatek *et al.*<sup>(80)</sup> elegantly show that there is a distinct early phase of GE with relatively rapid, uncontrolled passage of nutrients into the

duodenum, which is faster for larger meal volumes and unaffected by energetic load. However, subsequently the delivery of nutrients to the duodenum is related to the overall energy load of the meal, i.e. GE is inhibited more for greater energetic loads<sup>(80)</sup>. Subsequent work of our group with 500-ml meal shakes confirmed that energy density is the main driver of GE, but that greater viscosity additionally slows GE, and is more important for perceived fullness than gastric content volume<sup>(71)</sup>. Similarly, a semi-solid liquid meal had lower GE rate and resulted in greater suppression of appetite over 3 h than an iso-energetic liquid meal, despite lower plasma cholecystokinin (a satiety-related hormone) release in the first hour<sup>(81)</sup>. Appetite was correlated with gastric content volume rather than GE rate or plasma cholecystokinin, which suggests that the longer gastric retention was driving the decrease in appetite<sup>(81)</sup>. These studies illustrate the usefulness of MRI to study the gastric behaviour of different foods and drinks at a macroscopic level, especially in combination with other physiological measures and subjective ratings.

### ***In vivo* MRI: intra-gastric processes**

In addition to the processes discussed earlier, MRI is very suitable to investigate other intragastric processes such as layering (phase separation), mixing and macroscopic changes in the texture of the chyme, such as gelling or coagulation, because these will cause changes in the  $T_1$  and  $T_2$  relaxation rates and can thus be visualised. These processes depend on multiple food properties which are collectively referred to as the food matrix, which has been framed as 'a physical domain that contains and/or interacts with specific constituents of a food (e.g. a nutrient) providing functionalities and behaviours which are different from those exhibited by the components in isolation or a free state'<sup>(82)</sup>. The food matrix plays a key role in the kinetics of transit and hydrolysis of macronutrients<sup>(82)</sup>.

#### *Layering and fat quantification*


In the case of layer formation (phase separation), the different layers can simply be quantified on an MRI image. For example, when a more fatty layer forms this appears darker on a  $T_2$ -weighted gastric MRI image than a more watery layer, see e.g.<sup>(78)</sup>. Similarly, in the first hour after consumption breakfast porridges showed clear phase separation, with a brighter layer on top, consistent with a more liquid phase in the type of moderately  $T_2$ -weighted MRI images made, and a darker layer at the bottom, consistent with thicker or more particulate material<sup>(83)</sup>.

So far, we have mainly discussed volume measurements of different food fractions. However, after careful *in vitro* validation quantitative MRI measurements can be performed *in vivo*. For example, Marciani *et al.* investigated the dilution of polysaccharide test meals by gastric secretions<sup>(70)</sup> by exploiting the association between the  $T_2$  relaxation time ( $T_2$ ) and polysaccharide concentration<sup>(50)</sup>. MRI is also well-suited to distinguish water and fat. Kunz *et al.*<sup>(84)</sup> specifically measured the fat component of a pasta

meal (mayonnaise). By first assessing *in vitro* samples in which the fat concentration was varied they were able to calibrate their *in vivo* MRI measurements. Liu *et al.*<sup>(85)</sup> studied gastric and duodenal fat emptying and emulsion processing (creaming and phase separation) using fat emulsions that were administered through a nasogastric tube. They not only calibrated their fat quantification approach with *in vitro* MRI, but they also took gastric aspirate samples for further validation *in vivo*. The resulting fat fraction maps and intragastric emulsion profiles showed details of intraluminal phase separation and creaming that were not (well) visible on the conventional MRI images<sup>(85)</sup>. This approach was taken further by Scheuble *et al.*<sup>(86)</sup> who studied the gastric behaviour of fat emulsions stabilised by three different biopolymers *in vitro* as well as *in vivo* with MRI, combined with blood sampling for measurement of TAG and cholecystokinin concentrations. These studies on fat digestion show-case that MRI can bridge the gap between *in vitro* digestion models and *in vivo* behaviour by carefully combining different types of measurements.

### *Coagulation*

While eventual breakdown of structure is necessary to allow for GE, gastric conditions can also induce changes in the texture of the chyme such as gelling and coagulation. Since this involves the transformation from liquid to (semi-)solid it could slow GE. While coagulation will be readily visible on conventional stomach MRI images (see Fig. 3) it has hardly been systematically quantified. Coletta *et al.*<sup>(87)</sup> looked at GE of breads with different gluten contents and additionally visually categorised the degree of heterogeneity of the food bolus in the stomach on MRI scans. They found that gluten did not change GE, although it made the chyme more heterogeneous. Also, there were no differences between the breads in gastrointestinal symptoms, postprandial small bowel water content, colonic volume and gas content measured with MRI. Another example is milk protein coagulation; protein digestion is strongly affected by pH changes in the stomach and the associated activity of pepsin. Digestion by pepsin as well as the pH decline over time cause the caseins in milk to coagulate as demonstrated *in vitro*<sup>(88–90)</sup>, while the whey protein remains soluble. Casein coagulation is believed to slow down GE; the stomach empties only particles into the small intestine if they have a size of 1–2 mm<sup>(91)</sup>. This notion is supported by *in vivo* studies showing that amino acids from whey protein appear faster in the blood than those from casein<sup>(92,93)</sup>. *In vitro* data also show that casein coagulation is affected by several factors such as processing-induced protein modifications, product composition such as mineral composition, and variations in gastric acidification and protease secretion<sup>(94,95)</sup>. In addition, the source of the protein may influence gastric coagulation<sup>(96)</sup>. However, these findings require verification *in vivo*. This would benefit from accurate quantification of chyme structure changes. Although visual grading is a useful and relatively simple approach, this could be taken further by validating the use of image texture metrics that can capture the observed



**Fig. 3.** Examples of  $T_2$ -weighted magnetic resonance images showing cross-sections through an empty stomach after an overnight fast (baseline) and after 250 ml milk consumption. At  $t = 3$  min the gallbladder (gb) is clearly visible, and the gastric contents visible at baseline can be seen on top of the milk. At  $t = 30$  and 90 min milk protein coagulation can be observed.

heterogeneity; this may provide a more sensitive and objective assessment and information on what a certain degree of heterogeneity in an MRI image reflects in terms of particle sizes and texture attributes of coagulates. Such food matrix/chyme characteristics may influence gastric digestion and subsequent intestinal digestion and bioavailability (see e.g. Fardet *et al.*<sup>(97)</sup>).

#### Blood parameters

Although it is quite well possible to obtain blood samples from participants lying in an MRI scanner, not many studies have combined MRI of the digestive tract with blood sampling to assess hormone responses related to digestion and nutrient bioavailability (see e.g. Alyami *et al.*<sup>(83)</sup> and Mackie *et al.*<sup>(81)</sup>). GE and more detailed MRI markers of digestion could be linked to *in vivo* nutrient bioavailability measures such as blood glucose, fatty acid and amino acid profiles. For example, *in vitro* work showing that GE rate affects protein digestion, amino acid absorption and subsequent whole body protein anabolism after a meal<sup>(98)</sup> could be validated

with such an approach. One current candidate MRI marker for measuring the breakdown of protein foods is the  $T_2$  of the surrounding gastric fluid<sup>(51)</sup>, but as discussed earlier other MRI measures may also be of interest. Such studies would provide unprecedented detail on how food characteristics affect digestion and bioavailability and can inform optimised food design. For example, the effects of different protein sources and processing-induced protein modification on GE, protein digestion and amino acid absorption measured by MRI and amino acid absorption in the blood could be explored. Better understanding of the determinants of protein digestion can inform choices for or design of products that are easier digestible, which is beneficial for people who have trouble ingesting enough protein, such as older adults, athletes and critically ill.

#### Conclusion

In addition to more macroscopic structural information, NMR and MRI have the potential to provide molecular-



level and quantitative information on *in vivo* gastric (protein) digestion. This requires careful validation in order to understand what a specific MR parameter, or set of parameters, means in a specific digestion context. The resulting MR markers of digestion can be used to validate and inform *in vitro* digestion models and may bridge the gap between *in vitro* and *in vivo* studies. This can aid the optimisation of food properties for different applications in health and disease.

### Acknowledgements

We acknowledge the discussions with and continuing support of Monica Mars, Guido Camps, Kees de Graaf, Anja Janssen, Kasper Hettinga, Camilla Terenzi, John van Duynhoven and other collaborators.

### Financial Support

This work received no specific financial support from any funding agency, commercial or not-for-profit party. R. D. is funded by the China Scholarship Council. M. M. is financially supported by the Dutch Ministry of Economic Affairs; TKI-AF grant no. AF-18012 (Digestion & immunogenicity of proteins in infant nutrition).

### Conflict of Interest

None.

### Authorship

The authors jointly wrote and approved the paper. P. A. M. S. had final responsibility for the content.

### References

- Mackie A (2019) The digestive tract: a complex system. In *Interdisciplinary Approaches to Food Digestion*, pp. 11–27 [O Gouseti, G Bornhorst, S Bakalis and A Mackie, editors]. Cham: Springer International Publishing.
- Bornhorst GM, Gouseti O, Wickham MSJ *et al.* (2016) Engineering digestion: multiscale processes of food digestion. *J Food Sci* **81**, R534–R543.
- Smeets PAM, Erchner A & De Graaf C (2010) Cephalic phase responses and appetite. *Nutr Rev* **68**, 643–655.
- Hutchings JB & Lillford PJ (1988) The perception of food texture – the philosophy of the breakdown path. *J Texture Stud* **19**, 103–115.
- Krop EM, Hetherington MM, Nekitsing C *et al.* (2018) Influence of oral processing on appetite and food intake – a systematic review and meta-analysis. *Appetite* **125**, 253–269.
- Bornhorst GM (2017) Gastric mixing during food digestion: mechanisms and applications. *Annu Rev Food Sci Technol* **8**, 523–542.
- Brodkorb A, Egger L, Alminger M *et al.* (2019) INFOGEST static *in vitro* simulation of gastrointestinal food digestion. *Nat Protoc* **14**, 991–1014.
- Bornhorst GM & Singh RP (2014) Gastric digestion *in vivo* and *in vitro*: how the structural aspects of food influence the digestion process. *Annu Rev Food Sci Technol* **5**, 111–132.
- Luo Q, Boom RM & Janssen AEM (2015) Digestion of protein and protein gels in simulated gastric environment. *LWT – Food Sci Technol* **63**, 161–168.
- Bohn T, Carriere F, Day L *et al.* (2018) Correlation between *in vitro* and *in vivo* data on food digestion. What can we predict with static *in vitro* digestion models? *Crit Rev Food Sci Nutr* **85**, 2239–2261.
- Koziolek M, Grimm M, Becker D *et al.* (2015) Investigation of pH and temperature profiles in the GI tract of fasted human subjects using the Intellicap® System. *J Pharm Sci* **104**, 2855–2863.
- Kalantar-Zadeh K, Ha N, Ou JZ *et al.* (2017) Ingestible sensors. *ACS Sensors* **2**, 468–483.
- van Duynhoven J, Voda A, Witek M *et al.* (2010) *Time-domain NMR Applied to Food Products. Annual Reports NMR Spectroscopy*, vol. **69**, 1st ed., pp. 145–197 [G Webb, editor]. Oxford: Elsevier Ltd.
- Hatzakis E (2019) Nuclear magnetic resonance (NMR) spectroscopy in food science: a comprehensive review. *Compr Rev Food Sci Food Saf* **18**, 189–220.
- Marciani L, Cox EF, Hoad CL *et al.* (2013) Effects of various food ingredients on gall bladder emptying. *Eur J Clin Nutr* **67**, 2013/09/21, 1182–1187.
- Marciani L (2011) Assessment of gastrointestinal motor functions by MRI: a comprehensive review. *Neurogastroenterol Motil* **23**, 399–407.
- Spiller R & Marciani L (2019) Intraluminal impact of food: new insights from MRI. *Nutrients* **11**, 1147.
- Muttakin S, Moxon TE, Gouseti O (2019) *In vivo*, *in vitro*, and *in silico* studies of the GI tract. In *Interdisciplinary Approaches to Food Digestion*, pp. 29–67 [Gouseti O, Bornhorst GM, Bakalis S *et al.*, editors]. Cham: Springer International Publishing.
- Marze S (2017) Bioavailability of nutrients and micronutrients: advances in modeling and *in vitro* approaches. *Annu Rev Food Sci Technol* **8**, 35–55.
- Verhoeckx K, Cotter P, López-Expósito I *et al.* (editors) (2015) *The Impact of Food Bioactives on Health*. Cham: Springer International Publishing.
- Minekus M, Alminger M, Alvito P *et al.* (2014) A standardised static *in vitro* digestion method suitable for food-an international consensus. *Food Funct* **5**, 1113–1124.
- Kong F & Singh RP (2008) A model stomach system to investigate disintegration kinetics of solid foods during gastric digestion. *J Food Sci* **73**, 202–210.
- Minekus M, Marteau P, Havenaar R *et al.* (1995) A multi-compartmental dynamic computer-controlled model simulating the stomach and small intestine. *ATLA Altern Lab Anim* **23**, 197–209.
- de Wiele T, den Abbeele P, Ossieur W *et al.* (2015) The simulator of the human intestinal microbial ecosystem (SHIME®). In *Impact Food Bioactive Health Vitro ex vivo Model.*, pp. 305–317 [Verhoeckx K, Cotter P, López-Expósito I *et al.*, editors]. Cham: Springer International Publishing.
- Guo Q, Ye A, Lad M *et al.* (2015) Disintegration kinetics of food gels during gastric digestion and its role on gastric emptying: an *in vitro* analysis. *Food Funct* **6**, 756–764.
- Deng R, Mars M, Van Der Sman RGM *et al.* (2020) The importance of swelling for *in vitro* gastric digestion of whey protein gels. *Food Chem* **330**, 127182.
- Singh J, Dartois A & Kaur L (2010) Starch digestibility in food matrix: a review. *Trends Food Sci Technol* **21**, 168–180.



28. Golding M & Wooster TJ (2010) The influence of emulsion structure and stability on lipid digestion. *Curr Opin Colloid Interface Sci* **15**, 90–101.
29. Fabek H, Messerschmidt S, Brulport V *et al.* (2014) The effect of *in vitro* digestive processes on the viscosity of dietary fibres and their influence on glucose diffusion. *Food Hydrocoll* **35**, 718–726.
30. Soukoulis C, Fisk ID, Gan HH *et al.* (2016) Intragastric structuring of anionic polysaccharide kappa-carrageenan filled gels under physiological *in vitro* digestion conditions. *J Food Eng* **191**, 105–114.
31. Lorieau L, Halabi A, Ligneul A *et al.* (2018) Impact of the dairy product structure and protein nature on the proteolysis and amino acid bioaccessibility during *in vitro* digestion. *Food Hydrocoll* **82**, 399–411.
32. Somaratne G, Reis MM, Ferrua MJ *et al.* (2019) Mapping the spatiotemporal distribution of acid and moisture in food structures during gastric juice diffusion using hyperspectral imaging. *J Agric Food Chem* **67**, 9399–9410.
33. Bordoni A, Picone G, Babini E *et al.* (2011) NMR Comparison of *in vitro* digestion of Parmigiano Reggiano cheese aged 15 and 30 months. *Magn Reson Chem* **49**, 61–70.
34. Zerbe O & Jurt S (2014) *Applied NMR Spectroscopy for Chemists and Life Scientists*, 2nd ed. Weinheim: Wiley-VCH Verlag GmbH.
35. Nieva-Echevarría B, Goicoechea E, Manzanos MJ *et al.* (2014) A method based on  $^1\text{H}$  NMR spectral data useful to evaluate the hydrolysis level in complex lipid mixtures. *Food Res Int* **66**, 379–387.
36. Nieva-Echevarría B, Goicoechea E & Guillén MD (2017) Effect of liquid smoking on lipid hydrolysis and oxidation reactions during *in vitro* gastrointestinal digestion of European sea bass. *Food Res Int* **97**, 51–61.
37. Nieva-Echevarría B, Goicoechea E, Manzanos MJ *et al.* (2016) A study by  $^1\text{H}$  NMR on the influence of some factors affecting lipid *in vitro* digestion. *Food Chem* **211**, 17–26.
38. Nieva-Echevarría B, Goicoechea E, Manzanos MJ *et al.* (2015) Usefulness of  $^1\text{H}$  NMR in assessing the extent of lipid digestion. *Food Chem* **179**, 182–190.
39. Sundekilde UK, Jarno L, Eggers N *et al.* (2018) Real-time monitoring of enzyme-assisted animal protein hydrolysis by NMR spectroscopy – an NMR reactomics concept. *LWT* **95**, 9–16.
40. Barclay T, Ginic-Markovic M, Johnston M *et al.* (2012) Analysis of the hydrolysis of inulin using real time H-1 NMR spectroscopy. *Carbohydr Res* **352**, 117–125.
41. Dona AC, Pages G, Gilbert RG *et al.* (2010) Digestion of starch: *in vivo* and *in vitro* kinetic models used to characterise oligosaccharide or glucose release. *Carbohydr Polym* **80**, 599–617.
42. Mariette F (2009) Investigations of food colloids by NMR and MRI. *Curr Opin Colloid Interface Sci* **14**, 203–211.
43. Kirtil E, Cikrikci S, McCarthy MJ *et al.* (2017) Recent advances in time domain NMR & MRI sensors and their food applications. *Curr Opin Food Sci* **17**, 9–15.
44. Ziegler GR, MacMillan B & Balcom BJ (2003) Moisture migration in starch molding operations as observed by magnetic resonance imaging. *Food Res Int* **36**, 331–340.
45. Ozel B, Uguz SS, Kilercioglu M *et al.* (2017) Effect of different polysaccharides on swelling of composite whey protein hydrogels: a low field (LF) NMR relaxometry study. *J Food Process Eng* **40**, 1–9.
46. Peters JPCM, Vergeldt FJ, Van As H *et al.* (2016) Time domain nuclear magnetic resonance as a method to determine and characterize the water-binding capacity of whey protein microparticles. *Food Hydrocoll* **54**, 170–178.
47. Gianferri R, D’Aiuto V, Curini R *et al.* (2007) Proton NMR transverse relaxation measurements to study water dynamic states and age-related changes in Mozzarella di Bufala Campana cheese. *Food Chem* **105**, 720–726.
48. Le Dean A, Mariette F & Marin M (2004)  $^1\text{H}$  Nuclear magnetic resonance relaxometry study of water state in milk protein mixtures. *J Agric Food Chem* **52**, 5449–5455.
49. Bordoni A, Laghi L, Babini E *et al.* (2014) The foodomics approach for the evaluation of protein bioaccessibility in processed meat upon *in vitro* digestion. *Electrophoresis* **35**, 1607–1614.
50. Marciani L, Manoj P, Hills BP *et al.* (1998) Echo-planar imaging relaxometry to measure the viscosity of a model meal. *J Magn Reson* **135**, 82–86.
51. Deng R, Janssen AEM, Vergeldt FJ *et al.* (2020) Exploring *in vitro* gastric digestion of whey protein by time-domain nuclear magnetic resonance and magnetic resonance imaging. *Food Hydrocoll* **99**, 105348.
52. Wang Y-XJ, Zhang Q, Li X *et al.* (2015) T1ρ magnetic resonance: basic physics principles and applications in knee and intervertebral disc imaging. *Quant Imaging Med Surg* **5**, 858–885.
53. Palmer AG (2014) Chemical exchange in biomacromolecules: past, present, and future. *J Magn Reson* **241**, 3–17.
54. Ishima R (2014) CPMG relaxation dispersion. *Methods Mol Biol* **1084**, 29–49.
55. Ishima R & Bagby S (2018) Protein dynamics revealed by CPMG dispersion. In *Modern Magnetic Resonance*, pp. 435–452 [G Webb, editor]. Cham: Springer International Publishing.
56. Hansen AL, Lundström P, Velyvis A *et al.* (2012) Quantifying millisecond exchange dynamics in proteins by CPMG relaxation dispersion NMR using side-chain  $^1\text{H}$  probes. *J Am Chem Soc* **134**, 3178–3189.
57. Duvvuri U, Reddy R, Patel SD *et al.* (1997) T1rho-relaxation in articular cartilage: effects of enzymatic degradation. *Magn Reson Med* **38**, 862–867.
58. Henkelman M, Stanisz GJ & Graham SJ (2001) Magnetization transfer in MRI: a review. *NMR Biomed* **14**, 57–64.
59. Ye A, Liu W, Cui J *et al.* (2019) Coagulation behaviour of milk under gastric digestion: effect of pasteurization and ultra-high temperature treatment. *Food Chem* **286**, 216–225.
60. Van Zijl PCM & Yadav NN (2011) Chemical exchange saturation transfer (CEST): what is in a name and what isn’t? *Magn Reson Med* **65**, 927–948.
61. Longo D, di Gregorio E, Abategiovanni R *et al.* (2014) Chemical exchange saturation transfer (CEST): an efficient tool for detecting molecular information on proteins’ behaviour. *Analyst* **139**, 2687–2690.
62. Pavuluri K & McMahon MT (2017) Ph imaging using chemical exchange saturation transfer (CEST) MRI. *Isr J Chem* **57**, 862–879.
63. Zhou J, Payen J-F, Wilson DA *et al.* (2003) Using the amide proton signals of intracellular proteins and peptides to detect pH effects in MRI. *Nat Med* **9**, 1085–1090.
64. Sun PZ, Wang E & Cheung JS (2012) Imaging acute ischemic tissue acidosis with pH-sensitive endogenous amide proton transfer (APT) MRI – correction of tissue relaxation and concomitant RF irradiation effects toward mapping quantitative cerebral tissue pH. *Neuroimage* **60**, 1–6.
65. Consolino L, Anemone A, Capozza M *et al.* (2020) Non-invasive investigation of tumor metabolism and acidosis by MRI-CEST imaging. *Front Oncol* **10**, 161.

66. Berger A (2002) Magnetic resonance imaging. *Br Med J* **324**, 35.
67. de Zwart IM & de Roos A (2010) MRI for the evaluation of gastric physiology. *Eur Radiol* **20**, 2609–2616.
68. Schmidt SJ, Sun X, Litchfield JB *et al.* (1996) Applications of magnetic resonance imaging in food science. *Crit Rev Food Sci Nutr* **36**, 357–385.
69. Schwizer W, Fraser R, Borovicka J *et al.* (1994) Measurement of gastric emptying and gastric motility by magnetic resonance imaging (MRI). *Dig Dis Sci* **39**, 101S–103S.
70. Marciani L, Gowland PA, Spiller RC *et al.* (2001) Effect of meal viscosity and nutrients on satiety, intragastric dilution, and emptying assessed by MRI. *Am J Physiol Gastrointest Liver Physiol* **280**, 2001/05/16, G1227–33.
71. Camps G, Mars M, de Graaf C *et al.* (2016) Empty calories and phantom fullness: a randomized trial studying the relative effects of energy density and viscosity on gastric emptying determined by MRI and satiety. *Am J Clin Nutr* **104**, 73–80.
72. Abell TL, Camilleri M, Donohoe K *et al.* (2008) Consensus recommendations for gastric emptying scintigraphy: a joint report of the American neurogastroenterology and motility society and the society of nuclear medicine. *J Nucl Med Technol* **36**, 44–54.
73. Camps G, Mars M, Witteman BJM *et al.* (2018) Indirect vs direct assessment of gastric emptying: a randomized crossover trial comparing C-isotope breath analysis and MRI. *Neurogastroenterol Motil.* **30**, e13317.
74. Fruehauf H, Menne D, Kwiatek MA *et al.* (2011) Inter-observer reproducibility and analysis of gastric volume measurements and gastric emptying assessed with magnetic resonance imaging. *Neurogastroenterol Motil* **23**, 2011/07/12, 854–861.
75. Collins PJ, Houghton LA, Read NW *et al.* (1991) Role of the proximal and distal stomach in mixed solid and liquid meal emptying. *Gut* **32**, 615–9.
76. Santangelo A, Peracchi M, Conte D *et al.* (1998) Physical state of meal affects gastric emptying, cholecystokinin release and satiety. *Br J Nutr* **80**, 521–527.
77. Marciani L, Hall N, Pritchard SE *et al.* (2012) Preventing gastric sieving by blending a solid/water meal enhances satiety in healthy humans. *J Nutr* **142**, 1253–1258.
78. Camps G, Mars M, de Graaf C *et al.* (2017) A tale of gastric layering and sieving: gastric emptying of a liquid meal with water blended in or consumed separately. *Physiol Behav* **176**, 26–30.
79. Murray K, Placidi E, Schuring EA *et al.* (2015) Aerated drinks increase gastric volume and reduce appetite as assessed by MRI: a randomized, balanced, crossover trial. *Am J Clin Nutr* **101**, 2015/02/04, 270–278.
80. Kwiatek MA, Menne D, Steingoetter A *et al.* (2009) Effect of meal volume and calorie load on postprandial gastric function and emptying: studies under physiological conditions by combined fiber-optic pressure measurement and MRI. *Am J Physiol Gastrointest Liver Physiol* **297**, G894–G901.
81. Mackie AR, Rafiee H, Malcolm P *et al.* (2013) Specific food structures suppress appetite through reduced gastric emptying rate. *Am J Physiol Gastrointest Liver Physiol* **304**, 2013/04/13, G1038–43.
82. Aguilera JM (2019) The food matrix: implications in processing, nutrition and health. *Crit Rev Food Sci Nutr* **59**, 3612–3629.
83. Alyami J, Whitehouse E, Yakubov GE *et al.* (2019) Glycaemic, gastrointestinal, hormonal and appetitive responses to pearl millet or oats porridge breakfasts: a randomised, crossover trial in healthy humans. *Br J Nutr* **122**, 2019/11/12, 1142–1154.
84. Kunz P, Feinle-Bisset C, Faas H *et al.* (2005) Effect of ingestion order of the fat component of a solid meal on intragastric fat distribution and gastric emptying assessed by MRI. *J Magn Reson Imaging* **21**, 383–390.
85. Liu D, Parker HL, Curcic J *et al.* (2016) The visualisation and quantification of human gastrointestinal fat distribution with MRI: a randomised study in healthy subjects. *Br J Nutr* **115**, 2016/01/20, 903–912.
86. Scheuble N, Schaffner J, Schumacher M *et al.* (2018) Tailoring emulsions for controlled lipid release: establishing *in vitro-in vivo* Correlation for digestion of lipids. *ACS Appl Mater Interfaces* **10**, 17571–17581.
87. Coletta M, Gates FK, Marciani L *et al.* (2016) Effect of bread gluten content on gastrointestinal function: a crossover MRI study on healthy humans. *Br J Nutr* **115**, 2015/11/03, 55–61.
88. van Aken GA, Bomhof E, Zoet FD *et al.* (2011) Differences in *in vitro* gastric behaviour between homogenized milk and emulsions stabilised by Tween 80, whey protein, or whey protein and caseinate. *Food Hydrocoll* **25**, 781–788.
89. Ye A, Cui J, Dalgleish D *et al.* (2016) Formation of a structured clot during the gastric digestion of milk: impact on the rate of protein hydrolysis. *Food Hydrocoll* **52**, 478–486.
90. Ye A, Cui J, Dalgleish D *et al.* (2017) Effect of homogenization and heat treatment on the behavior of protein and fat globules during gastric digestion of milk. *J Dairy Sci* **100**, 36–47.
91. Kong F & Singh RP (2008) Disintegration of solid foods in human stomach. *J Food Sci* **73**, R67–R80.
92. Boirie Y, Dangin M, Gachon P *et al.* (1997) Slow and fast dietary proteins differently modulate postprandial protein accretion. *Proc Natl Acad Sci* **94**, 14930–14935.
93. Mahé S, Roos N, Benamouzig R *et al.* (1996) Gastrojejunal kinetics and the digestion of [15N]beta-lactoglobulin and casein in humans: the influence of the nature and quantity of the protein. *Am J Clin Nutr* **63**, 546–552.
94. Mulet-Cabero A-I, Mackie AR, Wilde PJ *et al.* (2019) Structural mechanism and kinetics of *in vitro* gastric digestion are affected by process-induced changes in bovine milk. *Food Hydrocoll* **86**, 172–183.
95. Wang X, Ye A, Lin Q *et al.* (2018) Gastric digestion of milk protein ingredients: study using an *in vitro* dynamic model. *J Dairy Sci* **101**, 6842–6852.
96. Hodgkinson AJ, Wallace OAM, Boggs I *et al.* (2018) Gastric digestion of cow and goat milk: impact of infant and young child *in vitro* digestion conditions. *Food Chem* **245**, 275–281.
97. Fardet A, Dupont D, Rioux L-E *et al.* (2019) Influence of food structure on dairy protein, lipid and calcium bioavailability: a narrative review of evidence. *Crit Rev Food Sci Nutr* **59**, 1987–2010.
98. Mulet-Cabero A-I, Torcello-Gómez A, Saha S *et al.* (2020) Impact of caseins and whey proteins ratio and lipid content on *in vitro* digestion and *ex vivo* absorption. *Food Chem* **319**, 126514.