


Erratum: ESHRE guideline: ovarian stimulation for IVF/ICSI

The ESHRE Guideline Group on Ovarian Stimulation, Ernesto Bosch¹, Simone Broer², Georg Griesinger³, Michael Grynberg⁴, Peter Humaidan⁵, Estratios Kolibianakis⁶, Michal Kunicki⁷, Antonio La Marca⁸, George Lainas⁹, Nathalie Le Clef¹⁰, Nathalie Massin¹¹, Sebastiaan Mastenbroek¹², Nikolaos Polyzos¹³, Sesh Kamal Sunkara¹⁴, Tanya Timeva¹⁵, Mira Töyli¹⁶, Janos Urbancsek¹⁷, Nathalie Vermeulen¹⁰, and Frank Broekmans  ^{2*}

¹IVI-RMS Valencia, Valencia, Spain ²Department of Reproductive Medicine and Gynecology, University Medical Center Utrecht, The Netherlands ³Department of Gynecological Endocrinology and Reproductive Medicine, University Hospital Schleswig-Holstein, Lübeck, Germany ⁴Department of Reproductive Medicine & Fertility Preservation, Hopital Antoine Bécélère, Clamart, France ⁵The Fertility Clinic, Skive Regional Hospital, Faculty of Health, Aarhus University, Skive, Denmark ⁶Unit for Human Reproduction, 1st Dept of ObGyn, Medical School, Aristotle University, Thessaloniki, Greece ⁷INVICTA Fertility and Reproductive Centre, Department of Gynaecological Endocrinology, Medical University of Warsaw, Poland ⁸Department of Obstetrics and Gynaecology, University of Modena Reggio Emilia and Clinica Eugin, Modena, Italy ⁹Eugonia, Athens, Greece ¹⁰European Society of Human Reproduction and Embryology, Belgium ¹¹Department of Obstetrics, Gynaecology and Reproduction, University Paris-Est Créteil, Centre Hospitalier Intercommunal, Créteil, France ¹²University Medical Center Amsterdam, Amsterdam, The Netherlands ¹³Dexeus University Hospital, Barcelona, Spain ¹⁴Department of Women and Children's Health, King's College London, UK ¹⁵Hospital "Dr. Shterev", Sofia, Bulgaria ¹⁶Kanta-Häme Central Hospital, Hämeenlinna, Mehiläinen Clinics, Helsinki, Finland ¹⁷Department of Obstetrics and Gynaecology, Semmelweis University Faculty of Medicine, Budapest, Hungary

*Corresponding author: Frank Broekmans, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands. E-mail: f.broekmans@umcutrecht.nl; guidelines@eshre.eu  <https://orcid.org/0000-0003-1366-2727>

Human Reproduction Open, Volume 2020, Issue 2, 2020, hoaa009, <https://doi.org/10.1093/hropen/hoaa009>

Oxford University Press would like to apologise to the Authors and readers for the introduction of two errors into this paper during the Production of the final version.

In the version published online on 01 May 2020 the text recommended that progesterone administration for luteal support after IVF/ICSI is conditional when this should be a strong recommendation. The correct recommendation is:

Progesterone is recommended for luteal phase support after IVF/ICSI. (van der Linden et al., 2015)	Strong ⊕○○○
--	----------------

The following recommendation was adjusted, although the recommendation remains correct, the GRADE mark format was altered. The correct recommendation is:

In urgent (oncology) fertility preservation cycles, random-start OS is an important option. (Boots et al., 2016)	Conditional ⊕⊕○○
--	---------------------

Further an extra z was added to the following recommendation although the recommendation remains correct. The correct recommendation is:

The association of follicle size as a triggering criterion with outcome has not been sufficiently studied. Physicians may choose the follicle size upon which final oocyte maturation is triggered on a case to case basis. (Chen et al., 2014)	Conditional ⊕⊕○○
---	---------------------
