


# Genetic, biochemical, and clinical spectrum of patients with mitochondrial trifunctional protein deficiency identified after the introduction of newborn screening in the Netherlands

Marit Schwantje<sup>1,2</sup> | Sabine A. Fuchs<sup>1</sup> | Lonneke de Boer<sup>3</sup> | Annet M. Bosch<sup>4</sup> | Inge Cuppen<sup>5</sup> | Eugenie Dekkers<sup>6</sup> | Terry G. J. Derks<sup>7</sup> | Sacha Ferdinandusse<sup>2</sup> | Lodewijk Ijlst<sup>2</sup> | Riekelt H. Houtkooper<sup>2</sup> | Rose Maase<sup>6</sup> | W. Ludo van der Pol<sup>5</sup> | Maaïke C. de Vries<sup>3</sup> | Rendelien K. Verschoof-Puite<sup>8</sup> | Ronald J. A. Wanders<sup>2</sup> | Monique Williams<sup>9</sup> | Frits Wijburg<sup>4</sup> | Gepke Visser<sup>1,2</sup> 

<sup>1</sup>Department of Metabolic Diseases, Wilhelmina Children's Hospital, University Medical Center Utrecht, Utrecht, The Netherlands

<sup>2</sup>Laboratory Genetic Metabolic Diseases, and Metabolism Institute, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands

<sup>3</sup>Department of Metabolic Diseases, Amalia Children's Hospital, Radboud University Medical Centre, Nijmegen, The Netherlands

<sup>4</sup>Department of Metabolic Diseases, Emma Children's Hospital, and Metabolism Institute, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands

<sup>5</sup>Department of Neurology and Neurosurgery, Wilhelmina Children's Hospital, University Medical Center Utrecht, Utrecht, The Netherlands

<sup>6</sup>National Institute for Public Health and the Environment (RIVM) Reference Laboratory for Pre- and Neonatal Screening, Center for Health Protection (R.M.) and Center for Population Screening (E.D), Bilthoven, The Netherlands

<sup>7</sup>Department of Metabolic Diseases, Beatrix Children's Hospital, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

<sup>8</sup>Department for Vaccine Supply and Prevention Programs, National Institute for Public Health and the Environment, Bilthoven, The Netherlands

<sup>9</sup>Department of Pediatrics, Center for Lysosomal and Metabolic Diseases, Erasmus MC, University Medical Center Rotterdam, Rotterdam, Netherlands

## Correspondence

Gepke Visser, Department of Metabolic Diseases, Wilhelmina Children's Hospital, University Medical Center Utrecht, Lundlaan 6. 3584 EA, Utrecht, The Netherlands.

Email: [gvisser4@umcutrecht.nl](mailto:gvisser4@umcutrecht.nl)

## Funding information

This research was funded by Metakids and PNOzorg. The content of the article has not been influenced by the sponsor.

## Abstract

Long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency (LCHADD) is included in many newborn screening (NBS) programs. Acylcarnitine-based NBS for LCHADD not only identifies LCHADD, but also the other deficiencies of the mitochondrial trifunctional protein (MTP), a multi-enzyme complex involved in long-chain fatty acid  $\beta$ -oxidation. Besides LCHAD, MTP harbors two additional enzyme activities: long-chain enoyl-CoA hydratase (LCEH) and long-chain ketoacyl-CoA thiolase (LCKAT). Deficiency of one or more MTP

[Click here to access the podcast for this paper.](#)

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](#) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Journal of Inherited Metabolic Disease* published by John Wiley & Sons Ltd on behalf of SSIEM.

**Communicating Editor:** Manuel Schiff

activities causes generalized MTP deficiency (MTPD), LCHADD, LCEH deficiency (not yet reported), or LCKAT deficiency (LCKATD). To gain insight in the outcomes of MTP-deficient patients diagnosed after the introduction of NBS for LCHADD in the Netherlands, a retrospective evaluation of genetic, biochemical, and clinical characteristics of MTP-deficient patients, identified since 2007, was carried out. Thirteen patients were identified: seven with LCHADD, five with MTPD, and one with LCKATD. All LCHADD patients (one missed by NBS, clinical diagnosis) and one MTPD patient (clinical diagnosis) were alive. Four MTPD patients and one LCKATD patient developed cardiomyopathy and died within 1 month and 13 months of life, respectively. Surviving patients did not develop symptomatic hypoglycemia, but experienced reversible cardiomyopathy and rhabdomyolysis. Five LCHADD patients developed subclinical neuropathy and/or retinopathy. In conclusion, patient outcomes were highly variable, stressing the need for accurate classification of and discrimination between the MTP deficiencies to improve insight in the yield of NBS for LCHADD. NBS allowed the prevention of symptomatic hypoglycemia, but current treatment options failed to treat cardiomyopathy and prevent long-term complications. Moreover, milder patients, who might benefit from NBS, were missed due to normal acylcarnitine profiles.

#### KEYWORDS

LCHAD deficiency, LCKAT deficiency, long-chain fatty acid oxidation, mitochondrial trifunctional protein complex, MTP deficiency, newborn screening

#### Synopsis

Accurate classification of and discrimination between the different MTP deficiencies are required to improve insight in the true yield of NBS for LCHADD.

## 1 | INTRODUCTION

Mitochondrial trifunctional protein (MTP) is a multi-enzyme complex, which catalyzes the last three steps of mitochondrial long-chain fatty acid  $\beta$ -oxidation (lcFAO) following the first step catalyzed by very-long-chain acyl-CoA dehydrogenase.<sup>1,2</sup> The MTP complex harbors three enzyme activities: long-chain enoyl-CoA hydratase (LCEH), long-chain 3-hydroxyacyl-CoA dehydrogenase (LCHAD), and long-chain ketoacyl-CoA thiolase (LCKAT).<sup>1,2</sup> Deficient function of one or more of these enzyme activities leads to either generalized MTP deficiency (MTPD, OMIM #609015), isolated LCHAD deficiency (LCHADD, OMIM #609016), or isolated LCKAT deficiency (LCKATD, no OMIM entry).<sup>3,4</sup> Until now, isolated LCEH deficiency has not been reported.

In many countries, LCHADD is included in population newborn screening (NBS) programs. NBS for LCHADD is generally based on abnormal acylcarnitine profiles measured by tandem mass spectrometry (MS/MS).<sup>2</sup> In the Netherlands, positive NBS results are followed by genetic

and enzymatic analysis to confirm or rule out the diagnosis, as required to reduce the risk of misdiagnosis.<sup>5</sup> Acylcarnitine profiles, however, do not fully discriminate between LCHADD, MTPD, and LCKATD. Thus, screening for LCHADD results in screening for MTPD and LCKATD as well.

Structurally, the MTP complex is a hetero-octamer consisting of four  $\alpha$ - and four  $\beta$ -subunits. LCHAD and LCEH activities are catalyzed by the  $\alpha$ -subunit and both are encoded by the *HADHA* gene. LCKAT activity is catalyzed by the  $\beta$ -subunit, which is encoded by the *HADHB* gene.<sup>4,6</sup> Both *HADHA* and *HADHB* are located on chromosome 2p23. Variants in *HADHA* may cause a reduced LCHAD activity. The most common cause of LCHADD is the c.1528G > C (p.Glu510Gln) variant, which has been identified in at least one allele in most patients.<sup>7</sup> Homozygosity for this variant causes LCHADD, whereas heterozygosity for this variant in combination with another *HADHA* variant may result in either LCHADD or MTPD, depending on the effect of the second variant on the stability of the MTP

complex.<sup>7,8</sup> Similarly, some *HADHB* variants leave the MTP complex intact and only affect the enzymatic activity of the  $\beta$ -unit, resulting in LCKATD, whereas other variants affect the proper formation or the stability of the MTP complex, causing loss of all three enzyme activities resulting in MTPD. The fact that *HADHA* and *HADHB* variants can cause LCHADD and LCKATD, respectively, but also MTPD complicates the diagnostic process. Hence, both genetic diagnostics and analysis of the enzyme activities of the MTP complex are essential for accurate discrimination between these related but distinct diseases.<sup>9</sup>

Patients with MTPD and LCHADD display heterogeneous clinical phenotypes varying from early-onset life-threatening cardiomyopathy, hypoketotic hypoglycemia, and liver failure, to a late-onset form with myopathy, episodic rhabdomyolysis, peripheral neuropathy, and/or pigmentary retinopathy.<sup>3</sup> LCKATD has only been reported in three patients: two with cardiomyopathy and all with fatal outcomes within 2 months of life.<sup>10,11</sup>

In the Netherlands, LCHADD has been included in the NBS program since 2007. Although LCHADD is the intended target disorder of NBS, also, patients with MTPD and LCKATD can be identified by NBS. To increase insight in patient outcomes, we evaluated the genetic, biochemical, and clinical characteristics of patients with MTP deficiency identified since the introduction of LCHADD in the Dutch NBS program.

## 2 | METHODS

### 2.1 | Study design and setting

We conducted a retrospective evaluation of genetic, biochemical, and clinical characteristics of patients with MTP deficiency (comprising LCHADD, MTPD, and LCKATD), diagnosed since the introduction of LCHADD in the Dutch NBS program (2007) until May 2021. We included both patients identified by NBS and patients diagnosed after clinical presentation. The study was approved by the Medical Ethics Committee of the University Medical Center Utrecht (METC10-430/C; METC19-234/M). Informed consent was obtained from living patients.

### 2.2 | Patients and diagnosis

Patients were identified via the Dutch Diagnosis Registration Metabolic Diseases (DDRMD, [www.ddrmd.nl](http://www.ddrmd.nl)). In the Netherlands, NBS is performed within 72–168 h after birth by measuring hydroxy-C16-carnitine levels with MS/MS in dried blood spots (DBS) (initial cutoff point [January 2007–October 2010]:  $\geq 0.20$   $\mu\text{mol/L}$ , adjusted

cutoff point since October 2010:  $\geq 0.08$   $\mu\text{mol/L}$ ). NBS results were available within 10 days after birth. The diagnosis was confirmed with biochemical and genetic analysis. Patients were classified based on enzyme activities in LCHADD, MTPD, and LCKATD patients.

### 2.3 | Biochemical studies

Primary patient skin fibroblasts were cultured in Ham's F-10 medium supplemented with 10% fetal calf serum (Invitrogen, Carlsbad, CA), 25-mmol/L HEPES, 100-U/ml penicillin, 100- $\mu\text{g/ml}$  streptomycin, and 250- $\mu\text{g/ml}$  amphotericin in a humidified atmosphere of 5%  $\text{CO}_2$  at 37°C. To investigate thermosensitivity in patient #5, cells were cultured at 40°C for 2 weeks, followed by enzyme and long-chain fatty acid  $\beta$ -oxidation (lcFAO) flux analysis. lcFAO flux analysis was performed in fibroblasts, as described earlier by measuring the production of radiolabeled  $\text{H}_2\text{O}$  from [9,10- $^3\text{H(N)}$ ]-oleic acid.<sup>12–14</sup> The palmitate loading test was performed in fibroblasts by adding 120  $\mu\text{mol/L}$  [U-13C]palmitate and 0.4 mmol/L L-carnitine to the medium, essentially as described by Diekman et al.<sup>15</sup> After 96 h, acylcarnitines in the medium were measured by MS/MS.

LCHAD and LCKAT activities were measured in lymphocyte and/or fibroblast homogenates using 3-ketopalmitoyl-CoA as substrate. Since approximately 75% of 3-hydroxyacyl-CoA dehydrogenase activity as measured with 3-keto-palmitoyl-CoA is catalyzed by LCHAD and the remaining 25% by short-chain hydroxyacyl-CoA dehydrogenase (SCHAD), the 3-ketopalmitoyl-CoA dehydrogenase activity is measured after a pre-incubation (30 min) with and without N-ethylmaleimide (30 mmol/L), which inactivates LCHAD but not SCHAD. LCHAD incubations were performed in 0.2-mol/L potassium phosphate, 0.1-mol/L 2-(N-morpholino)ethanesulfonic acid (MES) pH 6.2, 1-g/L Triton X-100, and 0.4-mmol/L NADH with 0.066-mg/ml protein at 37°C. The LCKAT incubations were performed in a solution of 0.1-mol/L TRIS pH 8.6, 25-mmol/L  $\text{MgCl}_2$ , 0.2-mg BSA/ml, 3-mmol/L Coenzyme A, and 1-g/L Triton X-100 with 0.066-mg/ml protein at 37°C. Reactions were started with 0.18-mmol/L 3-keto-palmitoyl-CoA and stopped after 5 min for LCHAD and after 10 min for LCKAT with 0.33-mol/L HCl and cooling on ice. After 5 min, the samples were neutralized with 2-mol/L KOH/0.6 M MES, and 30% (v/v) acetonitrile was added. After a further 5 min on ice, the samples were centrifuged for 5 min at 20 000g at 4°C. The supernatant was subjected to ultra-high-performance liquid chromatography on a C18 column (Waters Acquity HSS C18 1.8  $\mu\text{m}$  2.1  $\times$  100 mm). Resolution of the different CoA-esters (substrate and products) was achieved by a linear gradient of

TABLE 1 Baseline and birth characteristics of all patients with MTP deficiency

Patient number	MTP deficiency	Sex	Current age	Birth weight (in gr)	Pregnancy duration (wk + days)	Apgar scores (1 min, 5 min)	Pregnancy complications	Consanguinity	Ethnicity	Phenotype at time of diagnosis		
										Diagnosed by NBS?	Signs and symptoms	Other diseases?
#1	MPTD	F	Deceased at day 3	1580	33 + 2	1, 7	AFLP	Unknown	Other African	Yes <sup>c</sup>	Prenatal cardiomyopathy, lactic acidosis (33 mmol/L), hypoglycemia ( <i>hosp</i> )	Trombopenia
#2		M	Deceased at day 5	2455	36 + 2	8, 10	HELLP	No	Other African	Yes <sup>c</sup>	Cardiomyopathy, lactic acidosis (10.3 mmol/L) ( <i>hosp</i> )	No
#3		M	Deceased at day 31	1275	30 + 0	7, 8	Pre-eclampsia, IUGR	Yes	Unknown	Yes	Cardiomyopathy, lactic acidosis (>10 mmol/L), hypoglycemia ( <i>hosp</i> )	IRDS
#4		F	Deceased at day 10	2110	35 + 1	4, 8	Pre-eclampsia	Yes	Northern African	Yes	Cardiomyopathy ( <i>hosp</i> )	NEC
#5		F	11.9y	3700	38	“good start”	Pre-eclampsia, maternal diabetes	Yes	Northern African	No	Fever-induced muscle weakness, exercise intolerance, leg pain	No
#6	LCHADD	M	10.4y	3240	38 + 0	>9, >9	No	No	Caucasian	Yes	Weight loss, jaundice, fever, desaturation ( <i>hosp</i> )	No
#7		M	12.5y	1130	31 + 2	7, 8	HELLP, IUGR	No	Caucasian	Yes	No	No
#8		M	13.3y	3030	41	5, 9	No	No	Caucasian	Missed by NBS	Gastroenteritis with increased CK (1796 U/L) and ALAT (470 U/L)	No
#9		M	7.2y	2560	36 + 5	2, 7	ICP, HELLP	No	Caucasian	Yes	Fetal distress during delivery, postpartum hypoglycemia, increased NT-proBNP (3100 pg/ml) ( <i>hosp</i> )	CP due to asphyxia
#10		F	5.9y	2990	39 + 0	9, 10	No	No	Caucasian	Prenatal <sup>d</sup>	n.a.	No
#11		F	10.3y	2970	38 + 6	“good start”	No	No	Caucasian	Yes	Weight loss ( <i>hosp</i> )	No
#12		F	3.9y	3120	37 + 6	“good start”	No	No	Caucasian	Yes	No	No

(Continues)

TABLE 1 (Continued)

Patient number	MTP deficiency	Sex	Current age	Birth weight (in gr)	Pregnancy duration (wk + days)	Apgar scores		Pregnancy complications	Consanguinity	Ethnicity	Diagnosed by NBS?	Phenotype at time of diagnosis	
						(1 min)	(5 min)					Signs and symptoms	Other diseases?
#13	LCKATD	F	Deceased at 13mo	2194	38 + 5	9, 10	No	No	Caucasian	Yes <sup>c</sup>	Hypoglycemia, rhabdomyolysis (CK: 10004 U/L), cardiomyopathy (NT-proBNP >35 000 pg/ml), lactic acidosis (8.3 mmol/L) ( <i>hosp</i> )	Epilepsy	

Note: Reference values: lactate <2.0 mmol/L, CK < 145–200 U/L, ALAT < 57 U/L, NT-proBNP < 125–320 pg/ml.

Abbreviations: AFLP, acute fatty liver of pregnancy; ALAT, alanine aminotransferase; CK, creatine kinase; CP, cerebral palsy; gr, grams; HELLP, hemolysis, elevated liver enzymes, and low platelets syndrome; Hosp, the patient was hospitalized preceding the abnormal NBS results; IUGR, intrauterine growth retardation; ICP, intrahepatic cholestasis of pregnancy; IRDS, infantile respiratory distress syndrome; min, minutes; NEC, necrotizing enterocolitis; NT-proBNP, N-terminal pro-brain natriuretic peptide; Y, years; wk, weeks.

<sup>a</sup>According to the Dutch Perinatal Dictionary and Data (PWD).

<sup>b</sup>An lcFAO disorder was already suspected based on clinical presentation, before NBS results became available.

<sup>c</sup>Prenatal diagnosis through family screening (sibling of patient #11).

acetonitrile (from 33% to 37% (v/v)) in 16.9 mmol/L sodium phosphate buffer (pH 6.9) at a flow rate of 0.5 ml/min under continuous monitoring at 260 nm.

## 2.4 | Genetic analysis

Genetic analysis was performed by Sanger sequencing of all exons and flanking intronic sequences of the *HADHA* and *HADHB* genes. Sequence data were compared to the reference sequences NM\_000182.4 (*HADHA*) and NM\_000183.2 (*HADHB*) with nucleotide numbering starting at the first adenine of the translation initiation codon ATG.

## 2.5 | Clinical characteristics

Historical data were collected from the treating metabolic centers and the Dutch Expertise Center for lcFAO disorders. At the expertise center, patients were seen by a pediatric multidisciplinary team comprising metabolic specialists, dietitians, neurologists, and cardiologists. Ophthalmologic investigations were performed in the treating metabolic center.

## 3 | RESULTS

From January 2007 to May 2, 2021, 463 575 children were born in the Netherlands.<sup>16</sup> NBS was performed in over 99% of the newborns, leading to the referral of 41 newborns (January 2007–October 2010): 3 newborns, October 2010–May 2021: 38 newborns). With 30 false positives, 11 of the referred newborns were diagnosed with MTP deficiency. Eight were hospitalized preceding the abnormal NBS results (Table 1). Two additional patients were diagnosed after the clinical presentation (patients #5 and #8). Of the 13 patients, five had MTPD, seven had LCHADD, and one had LCKATD. Hydroxy-C16-carnitine levels in NBS samples and blood plasma after referral or clinical presentation are shown in Table 2.

### 3.1 | Biochemical characteristics

The results of genetic and enzymatic analyses are summarized in Table 3. MTPD patients had both reduced LCHAD and LCKAT activities, LCHADD patients had only reduced LCHAD activity, and the LCKATD patient had only reduced LCKAT activity. Although LCKAT activity in fibroblasts was also slightly reduced in patients #10 and #12, LCKAT activity was normal in lymphocytes, resulting in the final diagnosis of LCHADD.



**TABLE 2** Acylcarnitine levels measured in (1) newborn screening bloodspots, (2) blood plasma directly after referral by NBS or during diagnostic trajectory (for patients #5 and #8), (3) the palmitate loading test performed in patient fibroblasts

Patient number	MTP deficiency	NBS results (in $\mu\text{mol/L}$ , in DBS)												
		C16-OH	C0 (ref)	C14:1 (ref)	C14-OH (ref)	C16-OH (ref)	C18-OH (ref)	C12 (0.0–0.8)	C14 (0.0–0.4)	C16 (0.0–4.3)	C16-OH (0–0.07)			
#1	MTPD	1.67 <sup>a</sup>	23.53 (22.3–54.8)	1.70 (0.02–0.18)	0.34 (0–0.04)	1.78 (0–0.02)	0.48 (0–0.4)	0.8	2.5	26.4	2.5	26.4	2.5	
#2		1.40	25.17 (22.3–54.8)	2.17 (0.02–0.18)	0.35 (0–0.04)	1.50 (0–0.02)	0.53 (0–0.4)	–	–	–	–	–	–	
#3		0.63	9.82 (22.35–54.8)	0.41 (0.02–0.18)	–	0.44 (0)	–	0.2	1.6	34.2	1.6	34.2	1.6	
#4		1.44	6.9 (22.35–54.8)	0.1 (0.02–0.18)	0.04 (0)	0.59 (0)	–	–	–	–	–	–	–	
#5		0.03	43 (22.3–54.8)	0.06 (0.02–0.18)	0.01 (0–0.04)	0.00 (0–0.02)	0.00 (0–0.02)	0.6	0.2	2.4	0.2	2.4	0.0	
#6	LCHADD	1.47	12.55 (22.3–54.8)	0.02 (0–0.04)	0.04 (0–0.04)	0.14 (0–0.02)	0.14 (0–0.04)	1.2	1.1	7.6	1.1	7.6	1.1	
#7		0.32	17.99 (20–55)	0.56 (0–0.17)	0.12 (0–0.04)	0.23 (0–0.02)	0.00 (0–0.05)	1.0	0.8	6.9	0.8	6.9	1.0	
#8		0.11	30.85 (22.30–54.8)	1.12 (0.02–0.18)	0.18 (0–0.04)	0.14 (0–0.02)	0.03 (0–0.04)	0.1	0.2	2.0	0.2	2.0	0.0	
#9		0.19	12 (5–35)	0.53 (0.01–0.34)	0.07 (0–0.01)	0.15 (0–0.01)	0.12 (0–0.01)	–	–	–	–	–	–	
#10		0.11	11.41 (20–55) <sup>b</sup>	0.10 (0–0.17) <sup>b</sup>	0.04 (0–0.04) <sup>b</sup>	0.15 (0–0.02) <sup>b</sup>	0.12 (0–0.05) <sup>b</sup>	1.5	1.4	13.7	1.4	13.7	1.8	
#11		2.19	30.36 (20–55)	0.92 (0–0.17)	0.25 (0–0.04)	0.62 (0–0.02)	0.00 (0–0.05)	–	–	–	–	–	–	
#12		1.02	26.0 (12–46)	2.37 (0–0.15)	0.52 (0–0.01)	1.00 (0–0.02)	–	1.3	1.1	9.9	1.1	9.9	1.0	
#13	LCKATD	4.49	75.62 (25–65)	0.32 (0–0.04)	0.42 (0–0.04)	2.81 (0–0.02)	0.77 (0–0.05)	0.2	0.5	14.0	0.2	14.0	0.9	

Note: Reference values and the number of decimals are shown as measured and reported by the performing metabolic laboratory.

Abbreviation: DBS, dried blood spots.

<sup>a</sup>NBS was performed too early (at day 1 of life).

<sup>b</sup>For patient #10, acylcarnitines were measured in umbilical cord blood, after prenatal diagnosis through family screening.

TABLE 3 Enzymatic and genetic characteristics of all patients with MTP deficiency

Patient number	MTP deficiency	Affected gene	Genetic variants		Allele 2	Enzyme activities in lymphocytes		Enzyme activities in skin fibroblasts		lcFAO-flux in skin fibroblasts
			Allele 1			LCHAD	LCKAT	LCHAD	LCKAT	
#1	MTPD	HADHB	c.181C > T (p.Arg61Cys)		c.254 + 5 G > A (suspected splicing defect)	20%	5%	20%	5%	-
#2		HADHB	c.18C > A (p.Tyr6X)		c.631-1 G > A (splicing defect)	29%	3%	-	-	-
#3		HADHB	c.354 + 5delG (skipping exon 6)		c.354 + 5delG (skipping exon 6)	-	-	14% <sup>a</sup>	18% <sup>a</sup>	-
#4		HADHB	c.209 + 1G > C (suspected splicing defect)		c.209 + 1G > C (suspected splicing defect)	50% <sup>a</sup>	8% <sup>a</sup>	-	-	-
#5		HADHB	c.397A > G (p.Thr133Ala)		c.397A > G (p.Thr133Ala)	-	-	40% At 40°C: 10%	33% At 40°C: 7%	99% At 40°C: 34%
#6	LCHADD	HADHA	c.1528G > C (p.Glu510Gln)		c.1528G > C (p.Glu510Gln)	23%	161%	10%	138%	24%
#7		HADHA	c.1528G > C (p.Glu510Gln)		c.1528G > C (p.Glu510Gln)	41%	89%	7%	103%	28%
#8		HADHA	c.1528G > C (p.Glu510Gln)		c.982G > A (p.Gly328Arg) + c.1072C > A (p.Gln358Lys)	26%	88%	10%	92%	83%
#9		HADHA	c.1528G > C (p.Glu510Gln)		c.1528G > C (p.Glu510Gln)	19%	133%	-	-	-
#10		HADHA	c.1528G > C (p.Glu510Gln)		c.2099delG (p.Gly700GlufsX30)	19%	82%	10%	43%	27%
#11		HADHA	c.1528G > C (p.Glu510Gln)		c.2099delG (p.Gly700GlufsX30)	25%	89%	-	-	-
#12		HADHA	c.1528G > C (p.Glu510Gln)		c.1432delG (p.Ala478Leufs*17)	41%	89%	14%	56%	22%
#13	LCKATD	HADHB	c.182G > A (p.Arg61His)		c.1289 T > C (p.Phe430Ser)	146%	9%	74%	4%	17%

Notes: LCHAD and LCKAT activities in the patient samples are expressed as % of the mean of the reference values. Shown long-chain fatty acid  $\beta$ -oxidation (lcFAO)-flux is the mean of two independent experiments and is expressed as % of the mean lcFAO-flux in two or three control cell lines measured in the same experiment.

<sup>a</sup>Measurements were performed with the formerly used spectrophotometric assay, as described by Wanders et al.<sup>9</sup>

Patient #5 had relatively high residual LCHAD and LCKAT activities in fibroblasts cultured at the standard temperature of 37°C. After culturing the fibroblasts at 40°C, LCHAD and LCKAT activities decreased three and five times compared to 37°C, respectively. Additionally, the lcFAO flux decreased from 99% of controls at 37°C to 34% at 40°C. These results show a thermosensitive MTP deficiency (Supplementary Material 1).

In the LCHADD patients, lcFAO flux was below 28% except for patient #8, who was missed by NBS and showed a lcFAO flux of 83%. Interestingly, the palmitate loading test was completely normal in patient #8, while all other LCHADD patients showed the characteristic accumulation of hydroxy-C16-carnitine (Table 2).

### 3.2 | Genetic characteristics

All MTPD patients were homozygous or compound heterozygous for *HADHB* variants (Table 3). All LCHADD patients carried *HADHA* variants, with at least one allele carrying the common c.1528G > C (p.Glu510Gln) variant. The LCKATD patient was compound heterozygous for the following *HADHB* variants: c.182G > A (p.Arg61His) and c.1289 T > C (p.Phe430Ser).

### 3.3 | Clinical characteristics

#### 3.3.1 | Generalized MTPD

Two male and three female patients had MTPD. In all patients, pregnancies were complicated with hemolysis, elevated liver enzymes and low platelets (HELLP) syndrome, acute fatty liver of pregnancy (AFLP), or pre-eclampsia (Table 1). Four MTPD patients died due to cardiomyopathy within the 1st month of life (median age of death: 8 days, range: 3–31 days); one MTPD patient is currently aged 11 years and experiences illness- and exercise-induced muscle symptoms.

The deceased patients were hospitalized before NBS could be performed or before the results became available. All had hypertrophic or dilated cardiomyopathy with a severely reduced left ventricle (LV) function, before the 8th day of life. Lactic acidosis was reported for three of them. At the time of death, MTP deficiency was suspected in all, and treatment by means of glucose infusion and inotropic and diuretic drugs had been initiated. In addition to multiple hypoglycemic events and cardiomyopathy, patient #3 had infantile respiratory distress syndrome without improvement despite treatment with surfactant and high-frequency ventilation. Patient #4 also developed necrotizing enterocolitis. Creatine kinase

(CK) levels were elevated up to 5876 U/L. Patients #3 and #4 were previously described by Diekman et al.<sup>17</sup>

MTPD patient #5 is now an 11-year-old girl. NBS showed normal levels of hydroxy-C16-carnitine. From the age of 3 years, she experienced fever-associated episodes of muscle weakness and exercise intolerance. Episodes lasted 2–4 weeks and began when body temperature normalized. Between episodes, she experienced exercise intolerance with a maximum walking time of 15 min and exercise-induced muscle weakness. On clinical examinations, muscle tendon reflexes were normal. Extensive biochemical, metabolic, and genetic diagnostics did not reveal abnormalities. Acylcarnitine profiles and CK levels, measured at seven different occasions, were always normal. At 10 years of age, whole-exome sequencing showed homozygosity for the following *HADHB* variant: c.397A > G (p.Thr133Ala). Subsequent enzyme analyses showed a thermosensitive MTP deficiency.

Cardiac screening with echocardiography showed normal cardiac function. Sensory nerve action potential (SNAP) amplitudes and conduction velocities (NCVs) were normal, but electromyography showed sporadic and small motor unit potentials.

#### *Development and growth*

Patient #5 had normal motor and cognitive development. She had a normal height corrected for target height but was overweight (Table 4).

#### *Dietary treatment*

Patients #1 to #4 received intravenous glucose infusion. Patient #3 was additionally treated with parenteral feeding, carnitine supplementation (100 mg/kg/day), long-chain triglyceride (LCT)-restriction, and medium-chain triglyceride (MCT)-enrichment. Since diagnosis, patient #5 was treated with but poorly complied with an LCT-restricted, MCT-supplemented, and carbohydrate-enriched diet during illness.

#### 3.3.2 | Isolated LCHADD

Four male and three female patients had LCHADD. All are alive with a current median age of 10 years (range: 3.9–13.3y) and a similar follow-up duration. In two patients, pregnancies were complicated with HELLP syndrome (Table 1). Clinical characteristics are summarized in Table 4.

Five LCHADD patients were identified by NBS. Patient #10, the younger sibling of patient #11, was diagnosed prenatally. Despite the initiation of dietary measures immediately after birth, NBS was still positive (Table 2). Patient #8 was missed by NBS (false-negative NBS results) and presented with increased CK and ALAT



TABLE 4 Clinical characteristics of all surviving patients

Cardiac	Muscular	Hepatic	Neuropathic	Ocular	Growth	Diet
<b>Abnormalities on echocardiography</b>						
	<b>Muscle pain and/or rhabdomyolysis</b>			<b>Pigmentary ophthalmoscopy</b>		
<b>Abnormalities on ECG (age detected)</b>	<b>Illness-induced (max CK)</b>	<b>Hypoglycemia, lowest known glucose level (age)</b>	<b>Abnormal reflexes (age first detected)</b>	<b>Peripheral neuropathy on NCS (if yes: age detected, if no: age last NCS)</b>	<b>Height corrected for TH (SD)</b>	<b>Total fat, MCT, and LCT (in energy%)</b>
<b>Abnormalities detected</b>	<b>Reversibility</b>	<b>Symptoms</b>				
#1 n.r.	Hypertrophic CMP, lowest LV FS 2.7% (prenatal)	Cardiac failure, death	Cardiac failure, death	Cardiac failure, death	n.a.	n.a.
#2 n.r.	Dilated CMP, lowest LV FS 9% (day 4)	Cardiac failure, death	Cardiac failure, death	Cardiac failure, death	n.a.	n.a.
#3 n.r.	Hypertrophic, dilated CMP, low cardiac output (day 6)	Cardiac and respiratory failure, death (day 6)	Cardiac and respiratory failure, death (day 6)	Cardiac and respiratory failure, death (day 6)	n.a.	n.a.
#4 n.r.	Dilated CMP, low cardiac output (1 wk)	Cardiac failure, death	Cardiac failure, death	Cardiac failure, death	n.a.	n.a.
#5 No	No	No	No	No	-0.06	+3.5
#6 No	Mild LV hypertrophy, normal LV function (1wk)	Yes, after start MCT-diet	Yes, muscle weakness (CK measured, 9y)	Yes (7y)	+1.76	-0.04
#7 No	Dilated CMP, lowest LVFS: 16% (4mo)	Yes, with spironolactone, furosemide, captopril	2 episodes (CK not measured, 11y)	No (11y)	-0.32	+1.0
#8 No	No	1 episode before diagnosis (1796 U/L)	1 episode with myoglobinuria (42 900 U/L, 8y)	Yes (7y)	+1.19	Total: 38 LCT: 39 MCT: 61
#9 No	No	1 episode (63 238 U/L)	2 episodes (CK not measured, 11y)	No (6y)	+0.38	Total: 35 LCT: 23 MCT: 77
#10 Aspecific abnormal repolarization (4y)	No	4 episodes (76 937 U/L)	No	No (5y)	-0.82	Total: 18 LCT: 45 MCT: 55
#11 No	Hypertrophic CMP, lowest LVFS: 20% (5mo)	Feeding problems	3 episodes* (6038 U/L, 8y)	Yes (5y)	-2.39	Total: 22 LCT: 40 MCT: 60

TABLE 4 (Continued)

Patient	Cardiac		Muscular		Hepatic		Neuropathic		Ocular		Growth		Diet	
	Abnormalities on ECG (age detected)	Abnormalities on echocardiography	Abnormalities on ECG (age detected)	Symptoms	Reversibility	Illness-induced (max CK)	Exercise-induced (max CK, age 1st episode)	Hypoglycemia, lowest known glucose level (age)	Abnormal reflexes (age first detected)	Peripheral neuropathy on NCS (if yes: age detected, if no: age last NCS)	Pigmentary retinopathy on ophthalmoscopy (if yes: age detected, if no: age last ophthalmoscopy)	Height corrected for TH (SD)	Weight for height (SD) energy%	Total fat, MCT, and LCT (in energy%)
#12	Aspecific abnormal repolarization (2y)	No	No	-	-	No	No	No	No	Yes (2y) <sup>d</sup>	-2.38	-2.55	-1.08	Total: 34 LCT: 100 MCT: 0
#13	n.r.	CMP, lowest LV FS 14% (day 4)	Cardiac failure, death	5 episodes (33 715 U/L)	n.a.	Yes, levels unknown (postpartum)	No	No	Not performed	Not performed	+0.9	n.r.	+0.1	Not calculated

Notes: The most recent growth and dietary characteristics of the expertise center for IcfAO disorders are shown. For patient #5, who did not visit the expertise center, the most recent characteristics of the treating metabolic center are shown.

Abbreviations: CK, creatine kinase (reference value: <145–200); CMP, compound muscle action potential; CMP, cardiomyopathy; ECG, electrocardiogram; energy%, energy percentage of total calorie intake; LCT, long-chain triglycerides; LV, left ventricle; LVFS, left ventricle fractional shortening; MCT, medium-chain triglycerides; mo, months; n.r., not reported; n.a., not applicable; NCS, nerve conduction studies; SD, standard deviation; SNAP, sensory nerve action potential; wk, week; Y, years.

<sup>a</sup>Patient #11 already experienced exercise-induced muscle pain from infancy but increased CK levels were not measured until the age of 8 years.

<sup>b</sup>Glucose concentrations were undetectable after birth (“low”).

<sup>c</sup>Electromyography was also performed. Results were nonspecific, showing sporadic and small motor unit potential, not specific for a myopathy.

<sup>d</sup>NCS showed decreased CMAP in the n. peroneus and decreased sensory nerve conduction velocity in the n. suralis. These abnormalities were most likely caused by a low temperature of the lower extremities.

during a bout of gastroenteritis at the age of 9 months. Mild muscular hypotonia and microcephaly ( $-2.7$  SD) were noticed. An abnormal acylcarnitine profile suggestive of LCHADD led to confirmatory enzymatic and genetic testing. The identification of this patient resulted in a lowering of the NBS cutoff point for hydroxy-C16-carnitine in DBS from  $\geq 0.20$   $\mu\text{mol/L}$  to  $\geq 0.08$   $\mu\text{mol/L}$ . A retrospective study was conducted on all newborns born in the period January 1st, 2007 til October 1st, 2010 ( $n = 500.000$ ). In the 23 DBS of newborns with hydroxy-C16-carnitine  $> 0.08$   $\mu\text{mol/L}$  and  $< 0.22$   $\mu\text{mol/L}$ , homozygosity or heterozygosity for the common c.1528G > C (p.Glu510Gln) variant was not found, suggesting that no cases were missed in this period.

Three patients had mild reversible abnormalities on echocardiography. Patient #6 only needed dietary measures for reversibility, whereas patients #11 and #7 were also temporarily treated with diuretics and ACE inhibitors. At the time of diagnosis, patients #11 and #7 had feeding problems. Patient #7 had asymptomatic hypoglycemia, low carnitine plasma concentration (free carnitine: 6.2  $\mu\text{mol/L}$ , ref: 20–55), and a slightly delayed motor development, normalizing after treatment initiation.

In patient #9, glucose concentrations were undetectable after birth (“low”). On the 1st day of life, the total glucose intake could not be decreased below 9.6 mg/kg/min. Besides patient #7, no other patient experienced a hypoglycemic event after diagnosis. ALAT was normal in all, except during rhabdomyolysis.

Illness-induced rhabdomyolysis was reported in three patients (patients #9, #10, and #11). Two patients experienced one (patient #6) or two (patient #7) episodes of severe muscle pain preceded by physical activity from the age of 8 and 11 years, respectively. Patient #11 reported muscle pain after minimal exercise from the age of 4 years onward and experienced several episodes of muscle pain after physical activity with increased CK levels from 8 years onward.

On neurologic examination, muscle strength was normal in all patients. Four showed decreased or absent tendon reflexes from the age of 8 to 11 years onward (Table 4). Clinical sensory examination was normal in most patients, with the exception of patient #8, who had abnormal vibration sense at the age of 12 years. Only patient #8 showed abnormal nerve conduction studies (NCS). NCS revealed decreased SNAP amplitudes in the n. suralis and decreased compound muscle action potential amplitudes in the n. peroneus at the age of 9 years, which remained stable during follow-up.

Pigmentary retinopathy without visual impairment or night blindness was diagnosed by ophthalmoscopy in three patients at the age of 2, 5, and 7 years in patients #12, #11, and #6, respectively.

### Development and growth

Most patients had a normal height and weight (Table 4). All had normal early cognitive development and all patients older than 4 years attended a regular primary school. Patient #7 was slightly delayed in motor development at 4 months, which recovered after treatment of his cardiomyopathy. Patient #9 had cerebral palsy due to birth asphyxia and showed delayed motor development, improving with physiotherapy. Patient #12 was on a bottom scooter and started walking at 22 months.

### Dietary treatment

After diagnosis, all patients were advised a maximal fasting time according to age. All but patient #7 used an LCT-restricted and MCT-enriched diet from diagnosis, in two (patients #11 and #12) combined with breastfeeding. Patient #7 started an LCT-restricted and MCT-enriched diet at 4 months of age and used carnitine supplementation temporarily. For patient #12, the diet was stopped gradually upon initiation of solid foods but restarted at the age of 2.5 years. All patients on the dietary treatment used supplementation of Omega-3 and Omega-6 fatty acids. Four patients (patients #6, #7, #10, and #11) needed a percutaneous endoscopic gastrostomy because of feeding problems (behavioral problems and frequent vomiting).

### 3.3.3 | Isolated LCKATD

The LCKATD patient was born after an uncomplicated pregnancy and delivery. Because of low birth weight (<2.3 SD), she was admitted and treated for hypoglycemia. She gradually deteriorated and developed cardiorespiratory insufficiency with lactic acidosis. On day 4, echocardiography showed reduced LV function (LVFS: 15%) with initially normal cardiac dimensions, later biventricular hypertrophy. She developed rhabdomyolysis. On treatment with intravenous glucose, LCT restriction, MCT- and ketone supplementation, inotropic drugs, diuretics, and ACE-inhibitors, she improved and cardiac function recovered.

After discharge at the age of 1 month, she had a relatively stable period. At 4 months, she developed epilepsy, which was treated with levetiracetam and later zonisamide. From 9 months onward, she experienced several metabolic exacerbations with deterioration of cardiac function and increased CK levels. During the fourth exacerbation, cardiac function did not recover despite intravenous glucose and cardiac medication. She died at the age of 13 months.

### Development and growth

She had a motor developmental delay. At the age of 12 months, her length was +0.9 SD, and her weight for height: +0.1 SD.

### Dietary treatment

She was treated with an LCT-restricted and MCT- and ketone-supplemented diet through tube feeding. Ketones were supplemented as sodium-D,L-3-hydroxybutyrate (300 mg/kg/day increasing to 500 mg/kg/day). Extensive details of this patient have been described in a separate report (under submission).

## 4 | DISCUSSION

Since the introduction of acylcarnitine-based NBS for LCHADD in the Netherlands, seven LCHADD patients (resulting birth prevalence: approximately 1 in 350 000) were diagnosed, of whom one was missed by NBS. Thereafter, the cutoff point for referral was adjusted. Since acylcarnitine profiles in LCHADD, LCKATD, and MTPD are similar, not only patients with LCHADD, but also four with MTPD and one with LCKATD (birth prevalence: 1 in 2 500 000) were diagnosed by NBS. During the study period, an additional patient with MTPD (birth prevalence: 1 in 500 000) was diagnosed based on clinical symptoms. All LCHADD patients and this latter patient are still alive. The four patients with MTPD referred by NBS died within their 1st month of life and the LCKATD patient at the age of 13 months, all because of cardiac failure.

None of the LCHADD patients developed symptomatic hypoglycemia or neurological symptoms, whereas, in an international pre-NBS cohort study, hypoglycemia was reported as one of the presenting symptoms in 78% of the LCHADD patients. Moreover, a quarter of the pre-NBS patients had psychomotor retardation before or at the time of diagnosis.<sup>18</sup> These differences point to a significant beneficial effect of NBS. However, LCHADD patients detected by NBS still developed cardiomyopathy (three out of six, excluding patient #8, who was missed by NBS) and episodes of rhabdomyolysis (five out of six), despite treatment with a maximum fasting time according to age and an LCT-restricted and MCT-enriched diet since diagnosis by NBS for most.

Furthermore, peripheral neuropathy and pigmentary retinopathy were detected preclinically in four of the LCHADD patients detected by NBS during routine evaluation. The course of these complications is often progressive and longer follow-up is needed to investigate the clinical consequences in this population. It has been

reported that the long-term complications are less severe in LCHADD patients under optimal and/or earlier initiated dietary therapy.<sup>19,20</sup> However, the early development of subclinical long-term complications in our patients and patients previously reported in the literature show that NBS diagnosis and early initiation of current treatment strategies cannot fully prevent these complications.

During 14 years of NBS for LCHADD in the Netherlands, 30 referred newborns had false-positive NBS results. The percentage of false positives was calculated to be approximately 0.0003% before adjustment of the cutoff point for NBS and 0.002% after adjustment. The increase of false-positive NBS results was a logical consequence of lowering the cutoff point of NBS for LCHADD. Although the amount of false positives was comparable to numbers previously reported by Sander et al. (0.001%),<sup>10</sup> comparison to other NBS programs is difficult because of variability between used screening markers, cutoff points, and classification of MTP deficiencies. One LCHADD patient had false-negative NBS results (14%). Other LCHADD patients missed by NBS have been reported in literature,<sup>21</sup> but the true amount of false negatives worldwide is thus far unclear.<sup>22</sup>

The LCHADD patient missed by NBS and detected by increased CK and ALAT during a bout of gastroenteritis at the age of 9 months had no metabolic derangements during the 12 years of follow-up but developed asymptomatic peripheral neuropathy. Grünert et al.<sup>23</sup> described a similar presentation in four patients with MTP deficiency due to biallelic *HADHA* variants (enzyme activities not reported) who were also missed by NBS. These four patients showed isolated peripheral neuropathy as the presenting symptom, and in three of them, this was the only symptom. Since MTP-related peripheral neuropathy is hypothesized to be caused by the toxicity of long-chain 3-hydroxyacyl fatty acid metabolites, it is surprising that this was the presenting symptom in patients without a characteristic NBS acylcarnitine profile. Although plasma acylcarnitines may not fully represent the accumulation of acylcarnitines in nerves,<sup>24</sup> this may suggest a different pathophysiological mechanism causing neuropathy in MTP deficiencies.

The target disorder for NBS in the Netherlands is LCHADD and therefore, the MTPD patient clinically diagnosed at the age of 9 years can officially not be classified as a false-negative NBS result. The diagnosis was established by whole-exome sequencing in combination with enzyme diagnostics because of unexplained muscle symptoms. Interestingly, her acylcarnitine profile was repeatedly normal. Normal acylcarnitine profiles under healthy conditions, and in one patient even during metabolic decompensation, have been described in the literature for other MTPD patients with a (neuro)myopathic

phenotype.<sup>25,26</sup> Dietary treatment combined with avoidance of provoking factors proved beneficial for most of them. Consequently, our patient might have had a more favorable outcome with less metabolic decompensations and a shorter diagnostic trajectory had she been identified by NBS.

MTPD patients identified by NBS died early in life despite early diagnosis and treatment initiation. Although MTPD is known for its high mortality rates,<sup>27</sup> the broad clinical spectrum also comprises milder phenotypes with peripheral neuropathy and/or adult-onset myopathy.<sup>28,29</sup> Knowledge regarding the precise distribution of mild and severe phenotypes among the whole MTPD population is insufficient, but an overrepresentation of the severe phenotype in our cohort is possible. This might be the result of the genotypic spectrum with (expected) frameshift variants and deletions, which are associated with severe disease phenotypes.<sup>28</sup> However, given the fact that the deceased MTPD patients were already admitted before NBS results became available, it is also plausible that this severe MTPD phenotype is underreported in other patient cohorts.

It seems unlikely that an earlier diagnosis either by prenatal diagnosis or cord blood analysis would have prevented the fatal outcome of the severe MTPD patients. This implies that the currently available treatment options are insufficient and improved treatment strategies are needed. Recent studies suggest a beneficial effect of treatment with beta-hydroxybutyrate and triheptanoin in patients with (long-chain) fatty acid oxidation disorders.<sup>30–33</sup> The reported LCKATD patient, who was treated with sodium-D,L-3-hydroxybutyrate, had a longer survival time than LCKATD patients previously reported in the literature and potentially benefited from this treatment. However, both treatment strategies require tolerance for enteral feeding, which is a sometimes unavoidable obstacle in the treatment of critically ill patients. Intravenous beta-hydroxybutyrate has been investigated in adults with cardiac disease and might be a promising future treatment option.<sup>34</sup>

NBS might be more beneficial for patients with a milder, myopathic phenotype of MTPD, but for these patients, acylcarnitine-based NBS might be inadequate due to their often normal acylcarnitine profiles. An alternative approach is NBS by next-generation sequencing (NGS). NGS has already identified patients with homozygous or compound heterozygous variants in *HADHA* and *HADHB* within cohorts of patients with myopathic symptoms or peripheral neuropathy, not identified by acylcarnitine profiling.<sup>35</sup> To further classify the patients and prevent the identification of variants without functional consequences, genetic-based NBS should always be combined with enzyme analyses.



Prognostication after referral by NBS by acylcarnitine profiling for a possible MTP deficiency is difficult, because of the broad phenotypic spectrum. Spiekerkoetter et al.<sup>28</sup> reported a genotype–phenotype correlation for MTPD, where *HADHB* missense variants were expected to present with milder phenotypes than premature termination or frameshift variants. Although functional studies were also expected to allow prognostication, a clear correlation between MTP enzyme activities and clinical phenotype has not been established.<sup>28,36</sup> Despite the detection of a correlation between the results of metabolic flux studies in cultured skin fibroblasts and phenotypic severity,<sup>28,36</sup> the current knowledge on a relation between enzymatic or genotypic characteristics and clinical outcomes is still too limited for accurate prognostication. Even LCHADD patients with identical genotypes (homozygosity for the common c.1528G > C [p.Glu510Gln] variant) show heterogeneous clinical phenotypes, suggesting the influence of environmental and other genetic factors.<sup>18</sup>

## 5 | CONCLUSION

We report the outcome of 13 patients diagnosed since the introduction of LCHADD in the Dutch NBS program. Since NBS for LCHADD by acylcarnitine profiling identifies all three MTP deficiencies, the identified patients comprised seven LCHADD, five MTPD, and one LCKATD. Their clinical presentation and outcomes were highly variable. Additionally, milder patients with normal acylcarnitine profiles, for whom dietary treatment might be most beneficial, were missed. This could be solved with NBS by NGS, followed by enzyme analysis.

The most apparent benefit of NBS for LCHADD was the prevention of symptomatic hypoglycemia. However, the currently available treatment options could not prevent all symptoms and long-term complications. To enhance the benefits of NBS, more effective treatment strategies are needed. Moreover, accurate classification of and discrimination between the different MTP deficiencies utilizing both enzymatic and genetic analysis is required to improve insight in the yield of NBS, prognosis prediction, and patient outcomes.

## CONFLICT OF INTEREST

Marit Schwantje, Sabine Fuchs, Lonneke de Boer, Annet Bosch, Inge Cuppen, Eugenie Dekkers, Terry Derks, Sacha Ferdinandusse, Lodewijk IJlst, Riekelt Houtkooper, Rose Maase, Ludo van der Pol, Maaïke de Vries, Rendelien Verschoof-Puite, Ronald Wanders, Monique Williams, Frits Wijburg, and Gepke Visser declare to have no potential conflicts of interests. None of the authors have accepted reimbursements, fees, funds, or salaries

from an organization that may in any way gain or lose financially from the results reported in this manuscript. None of the authors have any competing interests regarding relevant financial activities outside the submitted work, intellectual property, or any other relationships.

## AUTHOR CONTRIBUTIONS

M.S. and G.V. were involved in acquisition of data, interpretation of data and drafting of the manuscript. All authors were involved in the interpretation of data and reviewing and editing the manuscript. L.B., A.M.B, T.G.J. D, M.C.V., M.W. G.V., S.A.F., I.C., W.L.P. and M.S. were involved in the long-term care of patients. S.F. and L.I. were involved in the biochemical analysis for all patients. E.D., R.M. and R.K.V.P. are involved in the implementation in and evaluation of the Dutch NBS program. M.S. and G.V. take responsibility for the collection of data, the interpretation and publication. All authors have given final approval of the version to be published.

## ETHICS APPROVAL

The study has been approved by the Medical Ethics Committee of the University Medical Center Utrecht (METC10-430/C; METC19-234/M).

## PATIENT CONSENT STATEMENT

All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2000. All living patients/patients' parents gave their written consent to participate in this study.

## DATA AVAILABILITY STATEMENT

Additional clinical data, not described in this article, will be shared upon (reasonable) request to the corresponding author.

## ORCID

Gepke Visser  <https://orcid.org/0000-0002-7618-6767>

## REFERENCES

1. Uchida Y, Izai K, Orii T, Hashimoto T. Novel fatty acid beta-oxidation enzymes in rat liver mitochondria. II. Purification and properties of enoyl-coenzyme a (CoA) hydratase/3-hydroxyacyl-CoA dehydrogenase/3-ketoacyl-CoA thiolase trifunctional protein. *J Biol Chem*. 1992;267(2):1034-1041. [published Online First: January 15, 1992].
2. Carpenter K, Pollitt RJ, Middleton B. Human liver long-chain 3-hydroxyacyl-coenzyme a dehydrogenase is a multifunctional membrane-bound beta-oxidation enzyme of mitochondria. *Biochem Biophys Res Commun*. 1992;183(2):443-448. doi:10.1016/0006-291x(92)90501-b [published Online First: 1992/03/16].



3. Knottnerus SJG, Bleeker JC, Wust RCI, et al. Disorders of mitochondrial long-chain fatty acid oxidation and the carnitine shuttle. *Rev Endocr Metab Disord.* 2018;19(1):93-106. doi:10.1007/s11154-018-9448-1 [published Online First: 2018/06/22].
4. Bennett MJ, Rinaldo P, Strauss AW. Inborn errors of mitochondrial fatty acid oxidation. *Crit Rev Clin Lab Sci.* 2000;37(1):1-44. doi:10.1080/10408360091174169 [published Online First: 2000/03/29].
5. Lotz-Havla AS, Roschinger W, Schiergens K, et al. Fatal pitfalls in newborn screening for mitochondrial trifunctional protein (MTP)/long-chain 3-Hydroxyacyl-CoA dehydrogenase (LCHAD) deficiency. *Orphanet J Rare Dis.* 2018;13(1):122. doi:10.1186/s13023-018-0875-6 [published Online First: 2018/07/22].
6. Kamijo T, Aoyama T, Miyazaki J, Hashimoto T. Molecular cloning of the cDNAs for the subunits of rat mitochondrial fatty acid beta-oxidation multienzyme complex. Structural and functional relationships to other mitochondrial and peroxisomal beta-oxidation enzymes. *J Biol Chem.* 1993;268(35):26452-26460. [published Online First: December 15, 1993].
7. IJlst L, Ruiten JP, Hoovers JM, Jakobs ME, Wanders RJA. Common missense mutation G1528C in long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency. Characterization and expression of the mutant protein, mutation analysis on genomic DNA and chromosomal localization of the mitochondrial trifunctional protein alpha subunit gene. *J Clin Invest.* 1996;98(4):1028-1033. doi:10.1172/JCI118863 [published Online First: August 15, 1996].
8. Boutron A, Acquaviva C, Vianey-Saban C, et al. Comprehensive cDNA study and quantitative analysis of mutant HADHA and HADHB transcripts in a French cohort of 52 patients with mitochondrial trifunctional protein deficiency. *Mol Genet Metab.* 2011;103(4):341-348. doi:10.1016/j.ymgme.2011.04.006 [published Online First: 2011/05/10].
9. Wanders RJA, Ruiten JP, IJlst L, Waterham HR, Houten SM. The enzymology of mitochondrial fatty acid beta-oxidation and its application to follow-up analysis of positive neonatal screening results. *J Inherit Metab Dis.* 2010;33(5):479-494. doi:10.1007/s10545-010-9104-8 [published Online First: May 22, 2010].
10. Sander J, Sander S, Steuerwald U, et al. Neonatal screening for defects of the mitochondrial trifunctional protein. *Mol Genet Metab.* 2005;85(2):108-114. doi:10.1016/j.ymgme.2005.02.002 [published Online First: 2005/05/18].
11. Das AM, Illsinger S, Lucke T, et al. Isolated mitochondrial long-chain ketoacyl-CoA thiolase deficiency resulting from mutations in the HADHB gene. *Clin Chem.* 2006;52(3):530-534. doi:10.1373/clinchem.2005.062000 [published Online First: 2006/01/21].
12. Manning NJ, Olpin SE, Pollitt RJ, Webley J. A comparison of [9,10-3H]palmitic and [9,10-3H]myristic acids for the detection of defects of fatty acid oxidation in intact cultured fibroblasts. *J Inherit Metab Dis.* 1990;13(1):58-68. doi:10.1007/BF01799333 [published Online First: 1990/01/01].
13. Olpin SE, Manning NJ, Pollitt RJ, Bonham JR, Downing M, Clark S. The use of [9,10-3H]myristate, [9,10-3H]palmitate and [9,10-3H]oleate for the detection and diagnosis of medium and long-chain fatty acid oxidation disorders in intact cultured fibroblasts. *Adv Exp Med Biol.* 1999;466:321-325. doi:10.1007/0-306-46818-2\_37 [published Online First: 2000/03/10].
14. Olpin SE, Manning NJ, Pollitt RJ, Clarke S. Improved detection of long-chain fatty acid oxidation defects in intact cells using [9,10-3H]oleic acid. *J Inherit Metab Dis.* 1997;20(3):415-419. doi:10.1023/a:1005358802096 [published Online First: 1997/07/01].
15. Diekman EF, Ferdinandusse S, van der Pol L, et al. Fatty acid oxidation flux predicts the clinical severity of VLCAD deficiency. *Genet Med.* 2015;17(12):989-994. doi:10.1038/gim.2015.22 [published Online First: 2015/04/04].
16. CBS. Population development; month and year. Centraal Bureau voor de Statistiek - Dutch Institute for Statistics; 2021. <https://opendata.cbs.nl/statline/#/CBS/nl/dataset/83474NED/table?ts=16330130245752021>
17. Diekman EF, Boelen CC, Prinsen BH, et al. Necrotizing enterocolitis and respiratory distress syndrome as first clinical presentation of mitochondrial trifunctional protein deficiency. *JIMD Rep.* 2013;7:1-6. doi:10.1007/8904\_2012\_128 [published Online First: 2013/02/23].
18. den Boer ME, Wanders RJA, Morris AA, et al. Long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency: clinical presentation and follow-up of 50 patients. *Pediatrics.* 2002;109(1):99-104. doi:10.1542/peds.109.1.99 [published Online First: 2002/01/05].
19. Immonen T, Ahola E, Toppila J, Lapatto R, Tyni T, Lauronen L. Peripheral neuropathy in patients with long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency - a follow-up EMG study of 12 patients. *Eur J Paediatr Neurol.* 2016;20(1):38-44. doi:10.1016/j.ejpn.2015.10.009 [published Online First: 2015/12/15].
20. Gillingham MB, Connor WE, Matern D, et al. Optimal dietary therapy of long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency. *Mol Genet Metab.* 2003;79(2):114-123. doi:10.1016/s1096-7192(03)00073-8 [published Online First: 2003/06/18].
21. Karall D, Brunner-Krainz M, Kogelnig K, et al. Clinical outcome, biochemical and therapeutic follow-up in 14 Austrian patients with long-chain 3-Hydroxy acyl CoA dehydrogenase deficiency (LCHADD). *Orphanet J Rare Dis.* 2015;10:21. doi:10.1186/s13023-015-0236-7 [published Online First: 2015/04/19].
22. Stinton C, Fraser H, Geppert J, et al. Newborn screening for long-chain 3-Hydroxyacyl-CoA dehydrogenase and mitochondrial trifunctional protein deficiencies using Acylcarnitines measurement in dried blood spots-a systematic review of test accuracy. *Front Pediatr.* 2021;9:606194. doi:10.3389/fped.2021.606194 [published Online First: 2021/04/06].
23. Grunert SC, Eckenweiler M, Haas D, et al. The spectrum of peripheral neuropathy in disorders of the mitochondrial trifunctional protein. *J Inherit Metab Dis.* 2021;44(4):893-902. doi:10.1002/jimd.12372 [published Online First: 2021/02/28].
24. Schooneman MG, Achterkamp N, Argmann CA, Soeters MR, Houten SM. Plasma acylcarnitines inadequately reflect tissue acylcarnitine metabolism. *Biochim Biophys Acta.* 2014;1841(7):987-994. doi:10.1016/j.bbali.2014.04.001 [published Online First: 2014/04/22].
25. Yagi M, Lee T, Awano H, et al. A patient with mitochondrial trifunctional protein deficiency due to the mutations in the HADHB gene showed recurrent myalgia since early childhood and was diagnosed in adolescence. *Mol Genet Metab.* 2011;104(4):556-559. doi:10.1016/j.ymgme.2011.09.025 [published Online First: 2011/10/18].

26. Spiekeroetter U, Bennett MJ, Ben-Zeev B, Strauss AW, Tein I. Peripheral neuropathy, episodic myoglobinuria, and respiratory failure in deficiency of the mitochondrial trifunctional protein. *Muscle Nerve*. 2004;29(1):66-72. doi:[10.1002/mus.10500](https://doi.org/10.1002/mus.10500) [published Online First: 2003/12/25].
27. den Boer ME, Dionisi-Vici C, Chakrapani A, van Thuijl AOJ, Wanders RJA, Wijburg FA. Mitochondrial trifunctional protein deficiency: a severe fatty acid oxidation disorder with cardiac and neurologic involvement. *J Pediatr*. 2003;142(6):684-689. doi:[10.1067/mpd.2003.231](https://doi.org/10.1067/mpd.2003.231) [published Online First: 2003/07/03].
28. Spiekeroetter U, Sun B, Khuchua Z, Bennett MJ, Strauss AW. Molecular and phenotypic heterogeneity in mitochondrial trifunctional protein deficiency due to beta-subunit mutations. *Hum Mutat*. 2003;21(6):598-607. doi:[10.1002/humu.10211](https://doi.org/10.1002/humu.10211) [published Online First: 2003/05/20].
29. Miyajima H, Orii KE, Shindo Y, et al. Mitochondrial trifunctional protein deficiency associated with recurrent myoglobinuria in adolescence. *Neurology*. 1997;49(3):833-837. doi:[10.1212/wnl.49.3.833](https://doi.org/10.1212/wnl.49.3.833) [published Online First: 1997/09/26].
30. Van Hove JL, Grunewald S, Jaeken J, et al. D,L-3-hydroxybutyrate treatment of multiple acyl-CoA dehydrogenase deficiency (MADD). *Lancet*. 2003;361(9367):1433-1435. doi:[10.1016/S0140-6736\(03\)13105-4](https://doi.org/10.1016/S0140-6736(03)13105-4) [published Online First: 2003/05/03].
31. Norris MK, Scott AI, Sullivan S, et al. Tutorial: triheptanoin and nutrition management for treatment of long-chain fatty acid oxidation disorders. *J Parenter Enteral Nutr*. 2021;45(2):230-238. doi:[10.1002/jpen.2034](https://doi.org/10.1002/jpen.2034) [published Online First: 2020/10/22].
32. Zoggeler T, Stock K, Jorg-Streller M, et al. Long-term experience with triheptanoin in 12 Austrian patients with long-chain fatty acid oxidation disorders. *Orphanet J Rare Dis*. 2021;16(1):28. doi:[10.1186/s13023-020-01635-x](https://doi.org/10.1186/s13023-020-01635-x) [published Online First: 2021/01/16].
33. Vockley J, Burton B, Berry G, et al. Effects of triheptanoin (UX007) in patients with long-chain fatty acid oxidation disorders: results from an open-label, long-term extension study. *J Inherit Metab Dis*. 2021;44(1):253-263. doi:[10.1002/jimd.12313](https://doi.org/10.1002/jimd.12313) [published Online First: 2020/09/05].
34. White H, Heffernan AJ, Worrall S, Grunfeld A, Thomas M. A systematic review of intravenous beta-Hydroxybutyrate use in humans—a promising future therapy? *Front Med (Lausanne)*. 2021;8:740374. doi:[10.3389/fmed.2021.740374](https://doi.org/10.3389/fmed.2021.740374) [published Online First: 2021/10/09].
35. Diebold I, Schon U, Horvath R, et al. HADHA and HADHB gene associated phenotypes—identification of rare variants in a patient cohort by next generation sequencing. *Mol Cell Probes*. 2019;44:14-20. doi:[10.1016/j.mcp.2019.01.003](https://doi.org/10.1016/j.mcp.2019.01.003) [published Online First: 2019/01/27].
36. Olpin SE, Clark S, Andresen BS, et al. Biochemical, clinical and molecular findings in LCHAD and general mitochondrial trifunctional protein deficiency. *J Inherit Metab Dis*. 2005;28(4):533-544. doi:[10.1007/s10545-005-0533-8](https://doi.org/10.1007/s10545-005-0533-8) [published Online First: 2005/05/20].

## SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** Schwantje M, Fuchs SA, de Boer L, et al. Genetic, biochemical, and clinical spectrum of patients with mitochondrial trifunctional protein deficiency identified after the introduction of newborn screening in the Netherlands. *J Inherit Metab Dis*. 2022;45(4):804-818. doi:[10.1002/jimd.12502](https://doi.org/10.1002/jimd.12502)