

most people; its use to determine what lung function an individual should expect should also raise alarm. Using specific genes that may explain lung function would be useful and doesn't have the same problem, though the information may still be abused. However, where specific genes are known there is no longer any need to know the "% African" inheritance or any other information on ethnicity. This is the logic that supports the screening of neonates for sickle cell genotypes without regard to reported ethnicity. Unfortunately, there has still been relatively little success in identifying the genes that are believed from family studies to explain around 50% of the variation in lung function. Until this situation improves, genes are of limited use in interpreting lung function. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Peter Burney, M.A., M.D.
Imperial College
London, United Kingdom

Corresponding author (e-mail: p.burney@imperial.ac.uk).

References

1. Witonsky J, Elhawary JR, Eng C, Rodríguez-Santana JR, Borrell LN, Burchard EG. Genetic ancestry to improve precision of race/ethnicity-based lung function equations in children. *Am J Respir Crit Care Med* 2022;205:725–727.
2. Hughes M, Hertel BR. The significance of color remains: a study of life chances, mate selection, and ethnic consciousness among Black Americans. *Soc Forces* 1990;68:1105–1120.
3. Hill ME. Color differences in the socioeconomic status of African American Men: results of a longitudinal study. *Soc Forces* 2000;78:1437–1460.

Copyright © 2022 by the American Thoracic Society



Finite Element Modeling of Pulmonary Mechanics in Severe Acute Respiratory Distress Syndrome: Explaining the Inclination Angle?

To the Editor:

With great interest, we read the research letter by Marazzo and colleagues (1) in which the effect of changing trunk inclination on respiratory function in patients with coronavirus disease (COVID-19)-associated acute respiratory distress syndrome is described. In our clinical practice, we repeatedly observed such rapid improvement of V_T s upon change of trunk position from

■ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0. For commercial usage and reprints, please e-mail Diane Gern (dgern@thoracic.org).

Originally Published in Press as DOI: 10.1164/rccm.202205-0931LE on June 2, 2022

semirecumbent to supine flat. The rapidity of altered respiratory mechanics does not, in our view, favor a major role for alveolar recruitment but rather a varying distention of different lung areas compatible with a patchy pathomorphological manifestation of COVID-19. This clinical observation led us to hypothesize that different configurations of, for example, fibrotic and emphysematous pulmonary parenchyma differentially influence lung aeration depending on the inclination of the individual patient's trunk.

Therefore, we set out to model diseased COVID-19 lungs using a finite element method-based mathematical pulmonary model using Ansys (2020 R2 ANSYS, Inc.), allowing simulation of distinct geometrical configurations of heterogeneously distributed parenchymal pathomorphology and its related local tissue mechanics (Figure 1). The model consists of five rows and three columns of rectangular blocks, representing a single lung, surrounded by a rigid supporting structure representing the thoracic cage. Each of the 15 blocks is composed of homogeneous material with mechanical properties described as elastic moduli, density, and yield strength as derived from the pathoanatomical literature and related to the modeled pathomorphological state of a specific anatomical lung area: fibrotic, emphysematous, or edematous (2, 3). The diaphragm at the caudal lung border is modeled as a plane with an imposed dynamic force simulating the intraabdominal pressure. The intraabdominal pressure depends on the degree of bed inclination: 8.4 mm Hg for 0°, 9.5 mm Hg for 15°, and 11 mm Hg for 30° of inclination (4). The gravitational force is represented by the x - and y -components, adapted accordingly for 15° and 30°. To test a representative set of clinically relevant pathomorphological manifestations as pulmonary COVID-19, more than 30 configurations of mechanical properties of the 15 blocks were tested, starting from homogeneous edema with an increasing number of blocks with fibrosis and emphysema in different and opposite positions (caudal versus cranial and dorsal versus ventral).

The output was defined as aeration of the lung, summed over the 15 individual modeled blocks. After simulating all different configurations in three bed inclination angles, it was particularly the modeled phenotype of an edematous lung configured with both cranial dorsal emphysema (≥ 3 blocks) and caudal dorsal fibrosis (≥ 3 blocks) that appeared to predict an increase in aeration when changing the bed inclination both from 0° to 15° and from 15° to 30° (Figure 1).

Although this straightforward computational finite element method approach is merely hypothesis generating, it is tempting to speculate whether combining these insights with pulmonary imaging techniques (e.g., computational tomography or electrical impedance tomography) would allow us to further validate our findings on a patient-specific basis. Ultimately, such modeling approach of pulmonary pathomorphological heterogeneity may advance our understanding of more patient-specific mechanical ventilation at an individualized degree of bed inclination in severe acute respiratory distress syndrome. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.




Figure 1. Example configuration of the finite element method lung model (upper part) and representation of the resulting lung aeration in 0°, 15°, and 30° bed inclination (lower part).

Eline Oppersma, Ph.D.*
 Stijn Geraats, B.Sc.
 Nicolien H. Heikens, B.Sc.
 Arjen Welleweerd, B.Sc.
 Chiel A. Woerts, B.Sc.
 Libera Fresiello, Ph.D.
 University of Twente
 Enschede, the Netherlands

Christiaan L. Meuwese, M.D., Ph.D.
 Erasmus Medical Center
 Rotterdam, the Netherlands

Dirk W. Donker, M.D., Ph.D.
 University of Twente
 Enschede, the Netherlands
 and

University Medical Center Utrecht
 Utrecht, the Netherlands

ORCID IDs: 0000-0002-0150-306X (E.O.); 0000-0003-0972-8340 (L.F.);
 0000-0002-1901-1653 (C.L.M.); 0000-0001-6496-2768 (D.W.D.).

*Corresponding author (e-mail: e.mos-oppersma@utwente.nl).

4. Samimian S, Ashrafi S, Khaleghdoost Mohammadi T, Yeganeh MR, Ashraf A, Hakimi H, *et al.* The correlation between head of bed angle and intra-abdominal pressure of intubated patients; a pre-post clinical trial. *Arch Acad Emerg Med* 2021;9:e23.

Copyright © 2022 by the American Thoracic Society



Reply to Oppersma *et al.*



From the Authors:

We thank Dr. Oppersma and colleagues for their interest in our study, in which we described the effects of changes in trunk inclination on respiratory mechanics in mechanically ventilated patients with COVID-associated acute respiratory distress syndrome (ARDS) (1). Reducing trunk inclination from semi-recumbent (40° head-of-bed elevation) to supine-flat (0° head-of-bed elevation) increased markedly (and reversibly) the compliance of the respiratory system, both due to an increase in chest wall and lung compliance. We did not measure changes in lung aeration (end-expiratory lung volume [EELV]); however, we think that the most relevant underlying mechanisms is the decrease in EELV in supine-flat position, favoring a reduction of

© This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0. For commercial usage and reprints, please e-mail Diane Gern (dgern@thoracic.org).

Originally Published in Press as DOI: 10.1164/rccm.202205-0987LE on June 2, 2022

References

- Marrazzo F, Spina S, Forlini C, Guarnieri M, Giudici R, Bassi G, *et al.* Effects of trunk inclination on respiratory mechanics in patients with COVID-19-associated acute respiratory distress syndrome: let's always report the angle! [letter] *Am J Respir Crit Care Med* 2022; 205:582–584.
- de Ryk J, Thiesse J, Namati E, McLennan G. Stress distribution in a three dimensional, geometric alveolar sac under normal and emphysematous conditions. *Int J Chron Obstruct Pulmon Dis* 2007;2:81–91.
- Lee HJ, Lee HJ, Lee JS, Kang YN, Koo JC. A built-up-type deformable phantom for target motion control to mimic human lung respiration. *Rev Sci Instrum* 2020;91:054106.