



## Sleep assessment in critically ill adults: A systematic review and meta-analysis

Ellaha Kakar, MD <sup>a,b,\*</sup>, Matthijs Priester, BSc <sup>c,1</sup>, Pascale Wessels, BSc <sup>d</sup>, Arjen J.C. Slooter, MD, PhD <sup>e,f</sup>, M. Louter, MD, PhD <sup>g</sup>, M. van der Jagt, MD, PhD <sup>b</sup>

<sup>a</sup> Department of Surgery, Erasmus MC, University Medical Center Rotterdam, the Netherlands

<sup>b</sup> Department of Intensive Care Adults, Erasmus MC, University Medical Center Rotterdam, the Netherlands

<sup>c</sup> Erasmus University Rotterdam, Rotterdam, the Netherlands

<sup>d</sup> Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

<sup>e</sup> Department of Intensive Care Medicine, UMC Utrecht Brain Center, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands

<sup>f</sup> Department of Neurology, UZ Brussel and Vrije Universiteit Brussel, Brussels, Belgium

<sup>g</sup> Department of Neurology, Erasmus MC, University Medical Center Rotterdam, the Netherlands

### ARTICLE INFO

#### Keywords:

Sleep  
Intensive care unit  
Critical care

### ABSTRACT

**Purpose:** To systematically review sleep evaluation, characterize sleep disruption, and explore effects of sleep disruption on outcomes in adult ICU patients.

**Materials and methods:** We systematically searched databases from May 1969 to June 2021 (PROSPERO protocol number: CRD42020175581). Prospective and retrospective studies were included studying sleep in critically ill adults, excluding patients with sleep or psychiatric disorders. Meta-regression methods were applied when feasible.

**Results:** 132 studies (8797 patients) were included. Fifteen sleep assessment methods were identified, with only two validated. Patients had significant sleep disruption, with low sleep time, and low proportion of restorative rapid eye movement (REM). Sedation was associated with higher sleep efficiency and sleep time. Surgical versus medical patients had lower sleep quality. Patients on ventilation had a higher amount of light sleep. Meta-regression only suggested an association between total sleep time and occurrence of delirium ( $p < 0.001$ , 15 studies, 519 patients). Scarce data precluded further analyses. Sleep characterized with polysomnography (PSG) correlated well with actigraphy and Richards Campbell Sleep Questionnaire (RCSQ).

**Conclusions:** Sleep in critically ill patients is severely disturbed, and actigraphy and RCSQ seem reliable alternatives to PSG. Future studies should evaluate impact of sleep disruption on outcomes.

© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

### 1. Introduction

Patients admitted to the intensive care unit (ICU) are known to be at risk for disrupted sleep. Important causes for sleep disruption are environmental factors, such as noise and light; physiologic factors, such as pain, immobility, discomfort and coughing; care-related factors, such as nursing care and medication, psychologic factors, and most importantly the severity of illness [1–8]. Disturbed sleep in critically ill patients has been associated with cognitive impairment, delirium, prolonged duration of mechanical ventilation, altered immune function, and

long-term psychological comorbidities, which may subsequently contribute to increased morbidity and mortality [1–8].

Polysomnography (PSG) is considered the reference standard to assess sleep quality and provides data on sleep architecture [9–11]. However, its use is logistically challenging and costly, especially in the ICU setting [1,7,12,13]. Therefore, many other sleep assessment tools, both objective (e.g. biosignals such as movements) and subjective (e.g. sleep questionnaires), are applied in the ICU setting.

Although previous studies have summarized sleep assessment methods [14] and risk factors for disrupted sleep [15], to our knowledge, no meta-analysis has been published on the characterization of sleep disruption, on sleep disruption in relation to mechanical ventilation and sedation or the impact of sleep disruption on relevant outcomes in critically ill patients. Therefore, we aimed to summarize current sleep assessment methods as a basis to characterize sleep disruption in ICU patients as compared to normal sleep, to assess the impact of

\* Corresponding author at: Doctor Molewaterplein 40, 3015 GD Rotterdam Room, NA-2123, the Netherlands.

E-mail address: [e.kakar@erasmusmc.nl](mailto:e.kakar@erasmusmc.nl) (E. Kakar).

<sup>1</sup> Shared first authorship.

additional ICU factors (e.g. mechanical ventilation and sedation) on sleep, and to study the associations between sleep characteristics and relevant health outcomes.

## 2. Materials and methods

### 2.1. Protocol and registration

This systematic review follows the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) [16] guidelines (see supplemental material S1) and was registered with PROSPERO database (<https://www.crd.york.ac.uk/PROSPERO>) as record number CRD42020175581.

### 2.2. Literature search strategy

A systematic literature search was performed with assistance of a biomedical information specialist, searching articles between May 14th, 1969 and June 24th, 2021. Main keywords included 'assessment tool', 'sleep', and 'intensive care unit'. The Embase, Medline Ovid, Cochrane Central, Web of Science and Google Scholar databases were searched. The pediatric and neonatal population was excluded from the search. The full search strategy is described in supplemental material S2.

### 2.3. Study screening and selection

Title and abstract screening were performed independently by three reviewers (EK/MP/PW) using the methods described by Bramer et al. [17] Prospective and retrospective studies were included if they studied adult patients ( $\geq 18$  years) admitted to the ICU in whom sleep quality and/or quantity was studied, using objective or subjective sleep assessment tools. These sleep assessment tools included PSG as an objective tool or patient questionnaires as subjective tools, evaluating sleep by healthcare professionals or investigators. Other inclusion criteria were a sample size of at least ten patients, full text availability and manuscript written in the English language. Two reviewers (MP/PW) assessed full texts independently. Studies were found eligible for full-text analysis if 'intensive care unit' or a closely related term such as 'coronary care unit' or 'acute care unit' were among the keywords of the paper, when focus on intensive care setting was not immediately clear from title/abstract screening. After full text screening, studies not performed in the intensive care were excluded. In randomized controlled trials that investigated the effect of an intervention on sleep, only the non-intervention control groups were included for this review, since the intervention may affect sleep outcomes. If randomization was performed in a crossover design, the outcomes of the 'before' study treatment were extracted. All disagreements between reviewers were resolved by discussion.

### 2.4. Main outcomes

This review first provides a summary of sleep assessment tools that are used in the critically ill. Furthermore, we assessed sleep characteristics based on these sleep assessment tools and compared them with sleep of healthy subjects. Further, we aimed to study the association of several patient or ICU management related factors on sleep characteristics (these subgroup analyses pertained to mechanical ventilation [MV], sedation, sedation level assessed with a validated scale [Richmond agitation sedation scale (RASS), Ramsay score], reason for ICU admission [e.g. medical versus surgical], severity of disease [acute physiology and chronic health evaluation: APACHE, Simplified Acute Physiology Score: SAPS]). Finally, we assessed associations between sleep characteristics and relevant health outcomes for ICU patients (delirium, length of ICU stay, length of mechanical ventilation, complications during ICU stay, and mortality). Whether tools were validated or not in the ICU

population was based on the Clinical Practice Guidelines for the Prevention and Management of Pain, Agitation/Sedation, Delirium, Immobility, and Sleep Disruption in Adult Patients in the ICU (PADIS) [1], the American Academy of Sleep Medicine [22], and a more recent study on actigraphy [154].

### 2.5. Data extraction and quality assessment


Two reviewers independently (MP/PW) extracted study data. The following study and patient characteristics of the included studies were extracted; author name, year of publication, type of study (e.g. retrospective, prospective, cohort, randomized controlled trial), sample size, reason for ICU admission (medical, surgical, or trauma), age, sex, sedation during ICU stay (yes/no), MV during ICU stay (yes/no), RASS, and APACHE scores. For the sleep assessment tools, we extracted type of sleep assessment tool (e.g. PSG, questionnaire, actigraphy, or other), the variables assessed in these tool, and whether the tool was validated in the ICU population. Furthermore, a senior author (MJ) checked at least 10% of the articles finally included by the junior authors to confirm and establish full consensus on the selection process. Studies were appraised, by two reviewers (MP/EK) during the data extraction phase, based on the method of Boulos et al. [18], in which appraisal was based on whether patients with sleep or psychiatric disorders were explicitly excluded.

### 2.6. Statistical analysis

Descriptive statistics were expressed in numbers (N) and percentages (%). Pooled means of the sleep variables were separately calculated for each assessment tool. Data were pooled, if  $> 3$  studies were available, using the "Metamean" function of R version 4.0.3 and the random effects model. Studies were included in the meta-analysis if mean values and standard deviations (SD) of the main outcomes were reported. If only medians were reported, these were used to calculate an approximation of the mean based on the methods of Hozo et al. [19] Approximations of the SD and interquartile ranges (IQR) were calculated as previously described [19-21]. Additionally, a variable named "slow wave sleep (SWS) merged" was created, in which data of N3 and N4 sleep stages (corresponding to deep sleep) were merged [22]. Heterogeneity was assessed using the  $I^2$  statistics. Comparison between the nocturnal PSG sleep variables assessed in this review and healthy volunteers from the study of Boulos et al. [18] was calculated using the two-sample  $t$ -test. Nocturnal pooled means and SD's were used for this review and compared with those in healthy volunteers reported by Boulos et al. [18]. Subgroup analysis of the pooled means of the sleep variables was performed, stratified by; MV, sedation, RASS, APACHE score, and reason for ICU admission (surgical, medical, trauma, and mixed [including surgical, medical, and/or trauma patients]) to explore associations of these variables with the sleep characteristics. Results of significant subgroup analysis were reported using the  $Q$ -test for heterogeneity ( $Q$ ). Meta-regression (linear for continuous normally distributed variables and logistic for binary variables) was performed to assess the associations between sleep characteristics and health outcomes, when data were available for the sleep variable and health outcome in  $\geq 10$  studies [20]. Statistical significance was inferred at  $p$ -value  $< 0.05$ .

## 3. Results

The initial search yielded 3421 articles, and after removing duplicates, 2151 were screened on title and abstract. After title and abstract screening, 280 articles were included for full-text assessment. Finally, 132 studies (8797 patients) were included (Fig. 1) [12,23-153].



**Fig. 1.** Flow diagram of article selection. N = number of studies, ICU = Intensive Care Unit.

### 3.1. Sleep assessment methods

Results on sleep assessment method and their characteristics are elaborated in detail and tabulated in supplemental material S3-S13. In brief, sixty-four studies used objective measurements (measuring bio-signals), 62 studies used previously validated subjective methods, and 32 studies used non-validated subjective methods.

### 3.2. Appraisal of studies

Appraisal of the studies can be found in supplemental material S14. Fifty-one studies (38.1%) excluded patients based on sleep disorders and 47 studies (35.1%) excluded patients based on psychiatric disorders. Eighteen of the 48 (37.5%) PSG studies excluded patients based on sleep disorders and 11/48 (22.9%) excluded patients based on psychiatric disorders. For the RCSQ these numbers were respectively 26/49 (53.1%) and 21/49 (42.9%).

### 3.3. Quantitative analysis

#### 3.3.1. Sleep characteristics

Data on sleep characterization could be pooled for PSG, RCSQ, and actigraphy, since sufficient data for analysis were available. These tools have been previously validated for sleep quality assessment [22,157].

**Table 1** Comparison of nocturnal PSG data between ICU population and healthy adults.

PSG variables	ICU			Healthy adults			P-value
	Mean	SD	N	Mean	SD	N	
TST, minutes	6.2	3.1	971	6.6	0.6	4038	<0.001
SE, %	52.2	18.2	554	85.7	6.8	4217	<0.001
WASO, minutes	122.9	58.8	155	48.2	21.7	2757	<0.001
N1, %	25.0	14.3	656	7.9	3.2	2940	<0.001
N2, %	49.7	12.7	775	51.4	6.2	2940	<0.001
SWS, %	13.0	10.0	893	20.4	7.3	2995	<0.001
REM, %	5.6	4.1	637	19.0	2.9	3012	<0.001
SOL, minutes	26.3	21.1	279	15.4	7.1	3828	<0.001
REML, %	134.9	15.9	44	97.4	16.9	2859	<0.001
AI, number per hour	18.0	9.5	317	12.6	3.9	2847	<0.001
Actigraphy Variables	ICU			Healthy adults			P-value
	Mean	SD	N	Mean	SD	N	
TST, minutes	8.1	6.3	491	7.2	0.8	21	0.006
SE, %	81.1	3.4	91	97.1	2.2	21	<0.001
WASO, minutes	71.8	17.3	100	12.6	9.6	21	<0.001

TST = Total Sleep Time, SE = Sleep Efficiency, WASO = Wake After Sleep Onset, N1 = light sleep stage, N2 = light sleep stage, REM = Rapid-Eye Movement, REML = REM Latency, SOL = Sleep Onset Latency, AI = Arousal Index, SD = standard deviation, N = number of patients.

We compared our pooled data to normal data of the concerned sleep assessment tool in healthy volunteers as the reference (quantitatively for actigraphy (Table 1) and qualitatively for RCSQ (S18A-E)). Supplemental material S15 provides an overview and specification of the items assessed with these sleep assessment methods and supplemental material S16A-C show the results of the pooled analyses of sleep characteristics. Comparison of PSG data of healthy volunteers [18] versus ICU-patients is shown in Table 1. Critically ill patients had a significantly lower sleep quality. Compared to normal actigraphy values [156], we also found that ICU patients had a lower total sleep time, sleep efficiency, and consequently a higher wake after sleep onset (Table 1). The RCSQ consists of six separate variables, as described in supplemental material S15, which can be converted to one score. A score between 0 and 25 is considered as very poor sleep, 26–50 as poor sleep, 51–75 as good sleep, and 76–100 as very good sleep [158]. A score of <50 on the RCSQ was found for the total RCSQ score, sleep depth, awakening, and quality.

### 3.3.2. Subgroup analysis

Subgroup analysis of sleep characteristics was performed for critically ill patients with or without MV, stratified by sedation administered or not, and ICU admission category. In the supplemental material, we only show statistically significant associations between sleep characteristics and subgroups. When analyzing PSG data, sedation was associated with a longer total sleep time (supplemental material S17A,  $p < 0.001$ ) and a higher sleep efficiency (supplemental material S17B,  $p = 0.01$ ) compared to patients without sedation. Patients with MV had a significantly lower amount of the sleep stage N1 (light sleep) (supplemental material S17C,  $p < 0.001$ ), a higher amount of sleep stage N2 (light sleep, supplemental material S17D,  $p = 0.005$ ) and a lower arousal index (S17E,  $p < 0.001$ ). Surgical patients had a significant lower sleep efficiency (supplemental material S17F,  $p = 0.004$ ), a higher sleep stage N1 (supplemental material S17G,  $p < 0.001$ ), and a lower wake after sleep onset (supplemental material S17H,  $p = 0.03$ ) compared with medical patients. Subgroup analysis for the RCSQ resulted in sedated patients scoring significantly higher on all RCSQ variables, except for the total score of the RCSQ (average all the variables) and noise, and thus having a more favorable sleep characterization (supplemental material S18A-E,  $p$ -values ranged from 0.01 to <0.001). No significant results were found in the subgroup analysis of actigraphy.

### 3.3.3. Associations of sleep variables with health outcomes

Sufficient data was available for PSG variables to perform only two meta-regression analyses, which showed significant associations between the total sleep time and occurrence of delirium (%) and between slow wave sleep (deep sleep) and duration of MV (days). Results of the meta-regression are depicted in Fig. 2. This shows that one hour increase in the total sleep time may be associated with 5.8% increase in risk of delirium ( $p < 0.001$ , 15 studies, 519 patients) and for each percent increase of slow wave sleep (deep sleep) there is an association with 0.58 days increase of MV duration ( $p = 0.01$ , 18 studies, 384 patients).

## 4. Discussion


In this review, we have provided an overview of sleep assessment methods studied in critically ill patients, including objective and subjective tools. Furthermore, based on qualitative analysis of PSG, RCSQ, and actigraphy data, we quantified in more detail the clear presence of significant sleep disruption, referring to significant differences in sleep duration and architecture, in critically ill patients as compared with healthy subjects, which has not been done in previous work. Also, we found that sedation, mechanical ventilation, and ICU admission type (surgical, medical or trauma) had significant effect on sleep characteristics as assessed with PSG, RCSQ, and actigraphy. Finally, increase in the total sleep time might be associated with increase in occurrence of

delirium, but scarce data precluded further analyses on relevant sleep-outcome associations and further research is warranted on these preliminary associations, given that these are likely influenced by other factors as well, and therefore unlikely to be uni-directional, or uni-factorial (e.g. associations between delirium and sleep are likely to be affected and confounded by sedation, disease severity, pain and its management, among others).

PSG is considered the reference standard in assessment of sleep quality, but unfortunately it is costly and labour intensive [1,7,9-13]. Recently, actigraphy has been found to be a reliable alternative to the PSG in the ICU population [154]. Based on the PSG and actigraphy analyses in this study, it seems that admission to the ICU causes difficulty in staying asleep and when asleep a reduction of deeper sleep stages occurs, which leads to loss of the restorative and resting function of sleep [6,66,159]. Actigraphy may be a more feasible tool to measure sleep quality at the ICU than PSG, although we found similar detection of sleep disruption based on sleep efficiency and awake after sleep onset-time, but a higher total sleep time as measured with actigraphy versus PSG (Table 1 and S16A-C). Regarding the RCSQ, we found that characteristics indicating sleep disruption seemed to correlate well with PSG, based on the results (S18A-E) of the RCSQ, which show the same disturbances in the equivalent sleep variables of PSG. RCSQ was previously validated against PSG [154,157,160]. Since the patient-nurse interrater reliability for the RCSQ is moderate, the RCSQ assessment is more reliable when assessed by the patient themselves [161].

Compared to PSG data of healthy volunteers [18], ICU patients' sleep is of lower quality. In the PADIS guidelines (2018) it is stated that total sleep time and sleep efficiency are normal in critically ill patients and that the percentage of light sleep is higher than deep sleep [1]. This review confirms the statement regarding the total sleep time and higher proportion of light sleep stages, but shows that sleep efficiency is reduced during critically illness. Interpretation of the total sleep time should be done with caution, since it constitutes an incomplete parameter to characterize the sleep quality and was also slightly different in critically ill versus healthy subjects hampering clinical relevance of this difference (although statistically significant). Also, in the studies that reported a TST the total recording time ranges from 7 to 24 h, which impacts on the total sleeping time. This study shows that ICU patients spend three times more time in the N1 sleep stage (light sleep) than healthy people, the same amount of time in the N2 sleep stage. Also, ICU patients spend 1.6 times and 3.4 times less time in the deeper sleep stage and REM, which are important for memory respectively emotion regulation, and may be relevant for recovery from illness [162,163]. Recent reviews have studied sleep assessment tools and risk factors for sleep disruption, but did not focus on sleep characterization in detail nor attempted meta-regression [14,15]. Therefore, the findings of these previous studies are complementary to this review. It should be noted that data from patients in the included studies who received an intervention of some kind which may influence the sleep quality were excluded from the analysis, however this concerned only a small minority of the total patients included for analysis.

Within the PSG studies a significant higher total sleep time (14.4 h. vs. 5.3 h.) and sleep efficiency (88.7% vs. 46.4%) in sedated patients compared to non-sedated ICU patients was found. This finding is interesting since sedatives have been considered as disrupting, rather than improving sleep quality [1]. Nevertheless, the clinical importance of these findings is unclear, since total sleep time and sleep efficiency are regarded as less important variables when it comes to the restorative function of sleep. Based on our results, it can be concluded that independent of sedative administration, all ICU patients have an impaired nocturnal sleep. Based on the RCSQ in this review, in which all items of the RCSQ were scored as "good" to "very good" in the sedated group compared to as "poor" in the non-sedated group, it seems that non-sedated patients experience more sleep



MV = Mechanical Ventilation, SWS = Slow Wave Sleep, TST = Total Sleep Time

Fig. 2. Bubble plot of the meta-regression.

MV = Mechanical Ventilation, SWS = Slow Wave Sleep, TST = Total Sleep Time.

disruption than sedated patients. In this study it also appears that mechanically ventilated patients seem to spend less time in lighter sleep stage (N1) compared to non-ventilated patients (22.2% vs. 51.2% (normal = 7.9% (SD = 3.2))) and have less arousals during sleep based on PSG data (13.5 vs. 50.0). The level of sedation may importantly influence these associations, but it was not possible to analyze this due to limited granularity of the pertinent data.

Regarding the differences in sleep characteristics found between ICU admission types, it seems that surgery also negatively influences sleep efficiency and thus the ability to reach deeper sleep stages, which is also concluded by a large multicentre study [164], assessing sleep in hospitalized patients. This difference may be caused by the level of pain in the surgical population, since the main reported reasons for sleep problems were noise and pain in that study [164]. Otherwise severity of illness or other variables may be confounding factors.

## 5. Strengths and limitations

To the best of our knowledge, this is the first meta-analysis that assessed and quantified sleep, using multiple sleep assessment methods, in the critically ill population in detail. However, several limitations should be discussed. First, there is a large heterogeneity in study populations, e.g. differences in diagnosis, medication usage, and ventilation protocols. Second, the sleep disruption-associated health outcomes we aimed to investigate (delirium, ICU LOS, length of MV, complications, and mortality) were scarcely reported. In addition, for MV we did not specify what type of ventilation mode was applied, but this has been investigated recently by Honarmand et al. [15] Third, pooled data for most of the questionnaires evaluating sleep quality could not be assessed. Fourth, although based on a large previous study [18], the quality assessment in our study may have been suboptimal and

therefore we want to emphasize the results should be interpreted with caution. Lastly, mere statistical associations cannot be distinguished from causal relations based on our analyses. The associations in our study should therefore be considered hypothesis generating rather than facts and therefore could guide future studies. For example, better sleep quality seems of high importance in general, but possibly the definition of 'better' sleep quality is different for the ICU population vs. healthy subjects.

## 6. Future studies

Although we provided an overview of sleep disruption in the critically ill and did so in more detail than was done in the PADIS guidelines and summarized sleep assessment methods and their possible yield in clinical practice, further study is needed to determine how assessment of sleep could be used to improve clinical management and outcomes. This review confirms that RCSQ and actigraphy may be reliable, valid, patient-friendly, and feasible alternatives to PSG for assessing sleep quality in the critically ill. It is known that the RCSQ assessment is most reliable when assessed by patients themselves [13]. Still, more studies are needed to define good sleep quality in the ICU population. Our analysis suggests that longer TST, while portending longer and possibly better quality of sleep (indicated by more SWS) in ICU patients, associates with poor outcomes. Although counterintuitive at first sight, this association may indeed signal that wakefulness in critically ill patients, rather than prolonged and even good quality sleep, is preferably maximized.

An important point should be noted regarding the use of PSG to characterize sleep disruption: PSG assesses important parts of the sleep architecture which currently can only be studied by the PSG. Therefore, although other measurement tools (mainly actigraphy and RCSQ) may provide promising results and may have practical advantages, PSG should not be discarded for further studies on sleep quality at the ICU. Therefore, other measurement techniques may provide promising results, but still validation of these new techniques using PSG are needed. A recent review regarding this subject also concludes the above ("ICU patient have atypical brain waves"), indicating that PSG evaluation to assess abnormality of sleep quality may be hampered) [14]. Compared with this previous review, our review characterized and quantified in more detail the clear presence of significant sleep disruption in critically ill patients compared with healthy subjects. Also, in this study associations between sleep and health outcomes were analyzed which has not been done in previous literature using quantified sleep data.

## 7. Conclusions

This review is the first meta-analysis to provide a detailed overview of sleep characteristics in ICU patients. Sleep in the critically ill is severely disturbed and can be characterized as short, light, and fragmented. Many methods in assessing sleep exist, with only a minority validated or immediately suitable for the ICU setting. Actigraphy and RCSQ seem to be reliable tools in assessing sleep in the critically ill as alternatives to PSG. However their usability and applicability seem to be more dependent on the type of management of the patient (e.g. sedation level) than PSG and therefore comparability is suboptimal. Furthermore, sedation, ventilation, and reason for admission alter sleep characteristics. Future studies investigating consequences of disrupted sleep for clinical outcomes are warranted.

## Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Author contributions

Conceptualization; MJ, EK Data curation; EK, MP, PW, Formal analysis; EK, MP, PW, Investigation; MJ, ML, AS, EK, MP, PW, Methodology; MJ, ML, AS, Supervision; MJ, ML, AS, Roles/Writing - original draft; EK, MP, Writing - review & editing; MJ, ML, AS, EK, MP, PW.

## Declaration of Competing Interest

None.

## Acknowledgements

The authors would like to thank W. Bramer, biomedical information specialist at the Medical Library of the Erasmus University Medical Center, for developing and updating the search strategies for this review. Author contribution: Each author certifies that he has made a direct and substantial contribution to the work reported in the manuscript. All authors approved the final version of the manuscript.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jcrr.2022.154102>.

## References

- [1] Devlin JW, Skrobik Y, Gelinas C, Needham DM, Slooter AJC, Pandharipande PP, et al. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adult patients in the ICU. *Crit Care Med*. 2018;46(9):e825–73.
- [2] Durmer JS, Dinges DF. Neurocognitive consequences of sleep deprivation. *Semin Neurol*. 2005;25(1):117–29.
- [3] Friese RS. Sleep and recovery from critical illness and injury: a review of theory, current practice, and future directions. *Crit Care Med*. 2008;36(3):697–705.
- [4] Opp MR. Sleep and psychoneuroimmunology. *Immunol Allergy Clin North Am*. 2009;29(2):295–307.
- [5] Orvelius L, Nordlund A, Nordlund P, Edell-Gustafsson U, Sjöberg F. Prevalence of sleep disturbances and long-term reduced health-related quality of life after critical care: a prospective multicenter cohort study. *Crit Care*. 2008;12(4):R97.
- [6] Pisani MA, Friese RS, Gehlbach BK, Schwab RJ, Weinhouse GL, Jones SF. Sleep in the intensive care unit. *Am J Respir Crit Care Med*. 2015;191(7):731–8.
- [7] Pulak LM, Jensen L. Sleep in the intensive care unit: a review. *J Intensive Care Med*. 2016;31(1):14–23.
- [8] Sanders RD, Maze M. Contribution of sedative-hypnotic agents to delirium via modulation of the sleep pathway. *Can J Anaesth*. 2011;58(2):149–56.
- [9] Bourne RS, Minelli C, Mills GH, Kandler R. Clinical review: sleep measurement in critical care patients: research and clinical implications. *Crit Care*. 2007;11(4):226.
- [10] Gong X, Huang L, Liu X, Li C, Mao X, Liu W, et al. Correlation analysis between polysomnography diagnostic indices and heart rate variability parameters among patients with obstructive sleep apnea hypopnea syndrome. *PLoS One*. 2016;11(6):e0156628.
- [11] Xie J, Yu W, Wan Z, Han F, Wang Q, Chen R. Correlation analysis between obstructive sleep apnea syndrome (OSAS) and heart rate variability. *Iran J Public Health*. 2017;46(11):1502–11.
- [12] Beecroft JM, Ward M, Younes M, Crombach S, Smith O, Hanly PJ. Sleep monitoring in the intensive care unit: comparison of nurse assessment, actigraphy and polysomnography. *Intensive Care Med*. 2008;34(11):2076–83.
- [13] Kamdar BB, Shah PA, King LM, Kho ME, Zhou X, Colantuoni E, et al. Patient-nurse interrater reliability and agreement of the Richards-Campbell sleep questionnaire. *Am J Crit Care*. 2012;21(4):261–9.
- [14] Richards KC, Wang YY, Jun J, Ye L. A systematic review of sleep measurement in critically ill patients. *Front Neurol*. 2020;11:542529.
- [15] Honarmand K, Rafay H, Le J, Mohan S, Rochweg B, Devlin JW, et al. A systematic review of risk factors for sleep disruption in critically ill adults. *Crit Care Med*. 2020;48(7):1066–74.
- [16] Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol*. 2009;62(10):1006–12.
- [17] Bramer WM, Milic J, Mast F. Reviewing retrieved references for inclusion in systematic reviews using EndNote. *J Med Libr Assoc*. 2017;105(1):84–7.
- [18] Boulos MI, Jairam T, Kendzerska T, Im J, Mekhael A, Murray BJ. Normal polysomnography parameters in healthy adults: a systematic review and meta-analysis. *Lancet Respir Med*. 2019;7(6):533–43.
- [19] Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol*. 2005;5:13.

- [20] Higgins JP, Altman DG, Gøtzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928.
- [21] Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol*. 2014;14:135.
- [22] Iber C, Ancoli-Israel S, Chesson AL, Quan S. The AASM manual for the scoring of sleep and associated events: Rules, terminology and technical specifications. Westchester, IL: American Academy of Sleep Medicine; 2007.
- [23] Aitken LM, Elliott R, Mitchell M, Davis C, Macfarlane B, Ullman A, et al. Sleep assessment by patients and nurses in the intensive care: an exploratory descriptive study. *Aust Crit Care*. 2017;30(2):59–66.
- [24] Alexopoulou C, Kondili E, Diamantaki E, Psarologakis C, Kokkini S, Bolaki M, et al. Effects of dexmedetomidine on sleep quality in critically ill patients: a pilot study. *Anesthesiology*. 2014;141:801–7.
- [25] Alexopoulou C, Kondili E, Plataki M, Georgopoulos D. Patient-ventilator synchrony and sleep quality with proportional assist and pressure support ventilation. *Intensive Care Med*. 2013;39(6):1040–7.
- [26] Alexopoulou C, Kondili E, Vakouti E, Klimathianaki M, Priniyanakis G, Georgopoulos D. Sleep during proportional-assist ventilation with load-adjustable gain factors in critically ill patients. *Intensive Care Med*. 2007;33(7):1139–47.
- [27] Alsulami G, Rice AM, Kidd L. Prospective repeated assessment of self-reported sleep quality and sleep disruptive factors in the intensive care unit: acceptability of daily assessment of sleep quality. *BMJ Open*. 2019;9(6).
- [28] Andréjak C, Monconduit J, Rose D, Toublanc B, Mayeux I, Rodenstein D, et al. Does using pressure-controlled ventilation to rest respiratory muscles improve sleep in ICU patients? *Respir Med*. 2013;107(4):534–41.
- [29] Arık E, Dolgun H, Hanalioglu S, Sahin OS, Ucar F, Yazicioglu D, et al. Prospective randomized study on the effects of improved sleep quality after craniotomy on melatonin concentrations and inflammatory response in neurosurgical intensive-care patients. *World Neurosurg*. 2020;140:253–9.
- [30] Bani Younis M, Hayajneh F, Alshraideh JA. Effect of noise and light levels on sleep of intensive care unit patients. *Nurs Crit Care*. 2019;26:73–8.
- [31] Bani Younis MK, Hayajneh FA, Alduraidei H. Effectiveness of using eye mask and earplugs on sleep length and quality among intensive care patients: a quasi-experimental study. *Int J Nurs Pract*. 2019;25(3):e12740.
- [32] Baumstarck K, Boucekine M, Estagnasie P, Geantot MA, Berric A, Simon G, et al. Assessment of patients' self-perceived intensive care unit discomforts: validation of the 18-item version of the IPREA. *Health Qual Life Outcomes*. 2019;17(1).
- [33] Bihari S, Doug McEvoy R, Matheson E, Kim S, Woodman RJ, Bersten AD. Factors affecting sleep quality of patients in intensive care unit. *J Clin Sleep Med*. 2012;8(3):301–7.
- [34] Boesen HC, Andersen JH, Bendtsen AO, Jennum PJ. Sleep and delirium in unsedated patients in the intensive care unit. *Acta Anaesthesiol Scand*. 2016;60(1):59–68.
- [35] Bosma K, Ferreyra G, Ambrogio C, Pasero D, Mirabella L, Braghioroli A, et al. Patient-ventilator interaction and sleep in mechanically ventilated patients: pressure support versus proportional assist ventilation. *Crit Care Med*. 2007;35(4):1048–54.
- [36] Bourne RS, Mills GH, Minelli C. Melatonin therapy to improve nocturnal sleep in critically ill patients: encouraging results from a small randomised controlled trial. *Crit Care*. 2008;12(2).
- [37] Boyko Y, Jennum P, Oerding H, Lauridsen JT, Nikolic M, Toft P. Sleep in critically ill, mechanically ventilated patients with severe sepsis or COPD. *Acta Anaesthesiol Scand*. 2018;62(8):1120–6.
- [38] Boyko Y, Toft P, Ørding H, Lauridsen JT, Nikolic M, Jennum P. Atypical sleep in critically ill patients on mechanical ventilation is associated with increased mortality. *Sleep Breath*. 2019;23(1):379–88.
- [39] Broughton R, Baron R. Sleep patterns in the intensive care unit and on the ward after acute myocardial infarction. *Electroencephalogr Clin Neurophysiol*. 1978;45(3):348–60.
- [40] Cabello B, Thille AW, Drouot X, Galia F, Mancebo J, D'Ortho MP, et al. Sleep quality in mechanically ventilated patients: comparison of three ventilatory modes. *Crit Care Med*. 2008;36(6):1749–55.
- [41] Campo FR, Drouot X, Thille AW, Galia F, Cabello B, D'Ortho MP, et al. Poor sleep quality is associated with late noninvasive ventilation failure in patients with acute hypercapnic respiratory failure. *Crit Care Med*. 2010;38(2):477–85.
- [42] Carrera-Hernandez L, Aizpitarte-Pejenaute E, Zugazagoitia-Ciarrusta N, Goni-Viguria R. Patients' perceptions of sleep in a critical care unit. *Enferm Intensiva*. 2018;29(2):53–63.
- [43] Casida JM, Davis JE, Zaleski A, Yang JJ. Night-time care routine interaction and sleep disruption in adult cardiac surgery. *J Clin Nurs*. 2018;27(7–8):e1377–84.
- [44] Chen JH, Chao YH, Lu SF, Shiang TF, Chao YF. The effectiveness of valerian acupuncture on the sleep of ICU patients: a randomized clinical trial. *Int J Nurs Stud*. 2012;49(8):913–20.
- [45] Chen LX, Ji DH, Zhang F, Li JH, Cui L, Bai CJ, et al. Richards-Campbell sleep questionnaire: psychometric properties of Chinese critically ill patients. *Nurs Crit Care*. 2019;24(6):362–8.
- [46] Cho MY, Min ES, Hur MH, Lee MS. Effects of aromatherapy on the anxiety, vital signs, and sleep quality of percutaneous coronary intervention patients in intensive care units. *Evid Based Complement Alternat Med*. 2013;2013.
- [47] Cochen V, Arnulf I, Demeret S, Neulat ML, Gourlet V, Drouot X, et al. Vivid dreams, hallucinations, psychosis and REM sleep in Guillain-Barré syndrome. *Brain*. 2005;128(11):2535–45.
- [48] Cooper AB, Thornley KS, Young GB, Slutsky AS, Stewart TE, Hanly PJ. Sleep in critically ill patients requiring mechanical ventilation. *Chest*. 2000;117(3):809–18.
- [49] Corbett SM, Rebeck JA, Greene CM, Callas PW, Neale BW, Healey MA, et al. Dexmedetomidine does not improve patient satisfaction when compared with propofol during mechanical ventilation. *Crit Care Med*. 2005;33(5):940–5.
- [50] Córdoba-Izquierdo A, Drouot X, Thille AW, Galia F, Roche-Campo F, Schortgen F, et al. Sleep in hypercapnic critical care patients under noninvasive ventilation: conventional versus dedicated ventilators. *Crit Care Med*. 2013;41(1):60–8.
- [51] Czempik PF, Jarosinska A, Machlowska K, Pluta M. Impact of light intensity on sleep of patients in the intensive care unit: a prospective observational study. *Indian J Crit Care Med*. 2020;24(1):33–7.
- [52] Darbyshire JL, Borthwick M, Edmonds P, Vollam S, Hinton L, Young JD. Measuring sleep in the intensive care unit: electroencephalogram, actigraphy, or questionnaire? *J Intensive Care Soc*. 2018;21(1):22–7.
- [53] Delisle S, Ouellet P, Bellemare P, Tetrault JP, Arsenault P. Sleep quality in mechanically ventilated patients: comparison between NAVA and PSV modes. *Ann Intensive Care*. 2011;1(1):42.
- [54] Demoule A, Carreira S, Lavault S, Pallanca O, Morawiec E, Mayaux J, et al. Impact of earplugs and eye mask on sleep in critically ill patients: a prospective randomized study. *Crit Care*. 2017;21(1).
- [55] Dennis CM, Lee R, Woodard EK, Szalaj JJ, Walker CA. Benefits of quiet time for neuro-intensive care patients. *J Neurosci Nurs*. 2010;42(4):217–24.
- [56] Díaz-Alonso J, Smith-Plaza AM, Suárez-Mier B, Lana A. Impact of a nurse intervention to improve sleep quality in intensive care units: results from a randomized controlled trial. *Dimens Crit Care Nurs*. 2018;37(6):310–7.
- [57] Diby M, Romand JA, Frick S, Heidegger CP, Walder B. Reducing pain in patients undergoing cardiac surgery after implementation of a quality improvement postoperative pain treatment program. *J Crit Care*. 2008;23(3):359–71.
- [58] Dres M, Younes M, Rittayamai N, Kendzerska T, Telias I, Grieco DL, et al. Sleep and pathological wakefulness at the time of liberation from mechanical ventilation (SLEWE) a prospective multicenter physiological study. *Am J Respir Crit Care Med*. 2019;199(9):1106–15.
- [59] Duclos C, Dumont M, Paquet J, Blais H, Van der Maren S, Menon DK, et al. Sleep-wake disturbances in hospitalized patients with traumatic brain injury: association with brain trauma but not with an abnormal melatonin circadian rhythm. *Sleep*. 2019;43(1):zsz191.
- [60] Elbaz M, Léger D, Sauvet F, Champigneulle B, Rio S, Strauss M, et al. Sound level intensity severely disrupts sleep in ventilated ICU patients throughout a 24-h period: a preliminary 24-h study of sleep stages and associated sound levels. *Ann Intensive Care*. 2017;7(1).
- [61] Elliott R, McKinley S, Cistulli P, Fien M. Characterisation of sleep in intensive care using 24-hour polysomnography: an observational study. *Crit Care*. 2013;17(2).
- [62] Engelmann C, Wallenborn J, Olthoff D, Kaisers UX, Ruffert H. Propofol versus flunitrazepam for inducing and maintaining sleep in postoperative ICU patients. *Indian J Crit Care Med*. 2014;18(4):212–9.
- [63] Engwall M, Fridh I, Johansson L, Bergbom I, Lindahl B. Lighting, sleep and circadian rhythm: an intervention study in the intensive care unit. *Intensive Crit Care Nurs*. 2015;31(6):325–35.
- [64] Ersoy EO, Ocal S, Kara A, Ardic S, Topeli A. Sleep in mechanically ventilated patients in the intensive care unit. *J Turk Sleep Med Turk Uykü Tbb Dergisi*. 2016;3(1):10–3.
- [65] Faraklas I, Holt B, Tran S, Lin H, Saffle J, Cochran A. Impact of a nursing-driven sleep hygiene protocol on sleep quality. *J Burn Care Res*. 2013;34(2):249–54.
- [66] Freedman NS, Gazendam J, Levan L, Pack AI, Schwab RJ. Abnormal sleep/wake cycles and the effect of environmental noise on sleep disruption in the intensive care unit. *Am J Respir Crit Care Med*. 2001;163(2):451–7.
- [67] Friese RS, Diaz-Arastia R, McBride D, Frankel H, Gentilello LM. Quantity and quality of sleep in the surgical intensive care unit: are our patients sleeping? *J Trauma*. 2007;63(6):1210–4.
- [68] Frisk U, Nordström G. Patients' sleep in an intensive care unit - Patients' and nurses' perception. *Intensive Crit Care Nurs*. 2003;19(6):342–9.
- [69] Hamze FL, de Souza CC, Chianca TC. The influence of care interventions on the continuity of sleep of intensive care unit patients. *Rev Lat Am Enfermagem*. 2015;23(5):789–96.
- [70] Hansen IP, Langhorn L, Dreyer P. Effects of music during daytime rest in the intensive care unit. *Nurs Crit Care*. 2018;23(4):207–13.
- [71] Hardin KA, Seyal M, Stewart T, Bonekat HW. Sleep in critically ill chemically paralyzed patients requiring mechanical ventilation. *Chest*. 2006;129(6):1468–77.
- [72] Hsu WC, Guo SE, Chang CH. Back massage intervention for improving health and sleep quality among intensive care unit patients. *Nurs Crit Care*. 2019;24(5):313–9.
- [73] Hu RF, Jiang XY, Hegadoren KM, Zhang YH. Effects of earplugs and eye masks combined with relaxing music on sleep, melatonin and cortisol levels in ICU patients: a randomized controlled trial. *Crit Care*. 2015;19(1).
- [74] Huttmann SE, Wilms K, Hamm C, Magnet FS, Windisch W, Storre JH. Assessment of sleep in patients receiving invasive mechanical ventilation in a specialized weaning unit. *Lung*. 2017;195(3):361–9.
- [75] Ibrahim MG, Bellomo R, Hart GK, Norman TR, Goldsmith D, Bates S, et al. A double-blind placebo-controlled randomised pilot study of nocturnal melatonin in tracheostomised patients. *Crit Care Resusc*. 2006;8(3):187–91.
- [76] Jean R, Shah P, Yudelevich E, Genese F, Gershner K, Leventowski D, et al. Effects of deep sedation on sleep in critically ill medical patients on mechanical ventilation. *J Sleep Res*. 2019;29(3):e12894.
- [77] Jones C, Dawson D. Eye masks and earplugs improve patient's perception of sleep. *Nurs Crit Care*. 2012;17(5):247–54.
- [78] Kalfon P, Mimos O, Auquier P, Loundou A, Gauzit R, Lepape A, et al. Development and validation of a questionnaire for quantitative assessment of perceived discomforts in critically ill patients. *Intensive Care Med*. 2010;36(10):1751–8.

- [79] Kamdar BB, Combs MP, Colantuoni E, King LM, Niessen T, Neufeld KJ, et al. The association of sleep quality, delirium, and sedation status with daily participation in physical therapy in the ICU. *Crit Care*. 2016;20(1).
- [80] Kamdar BB, Kadden DJ, Vangala S, Elashoff DA, Ong MK, Martin JL, et al. Feasibility of continuous Actigraphy in patients in a medical intensive care unit. *Am J Crit Care*. 2017;26(4):329–35.
- [81] Karadag E, Samancioglu S, Ozden D, Bakir E. Effects of aromatherapy on sleep quality and anxiety of patients. *Nurs Crit Care*. 2017;22(2):105–12.
- [82] Karaman Özlü Z, Özer N. The effect of enhancing environmental factors on the quality of patients' sleep in a cardiac surgical intensive care unit. *Biol Rhythm Res*. 2017;48(1):85–98.
- [83] Knauert MP, Yaggi HK, Redeker NS, Murphy TE, Araujo KL, Pisani MA. Feasibility study of unattended polysomnography in medical intensive care unit patients. *Heart Lung J Acute Crit Care*. 2014;43(5):445–52.
- [84] Kondili E, Alexopoulou C, Xirouchaki N, Georgopoulos D. Effects of propofol on sleep quality in mechanically ventilated critically ill patients: a physiological study. *Intensive Care Med*. 2012;1:7.
- [85] Kroll RR, McKenzie ED, Boyd JG, Sheth P, Howes D, Wood M, et al. Use of wearable devices for post-discharge monitoring of ICU patients: a feasibility study. *J Intensive Care*. 2017;5(1).
- [86] Krotsetis S, Richards KC, Behncke A, Köpke S. The reliability of the German version of the Richards Campbell sleep questionnaire. *Nurs Crit Care*. 2017;22(4):247–52.
- [87] Lewandowska K, Mędrzycka-Dąbrowska W, Kwiecień-Jaguś K, Czyż-Szympenbejl K. Factors determining sleep in patients hospitalized in ICUs in a hospital in northern Poland. *Sleep Biol Rhythms*. 2019;17(2):243–50.
- [88] Li SY, Wang TJ, Vivienne Wu SF, Liang SY, Tung HH. Efficacy of controlling nighttime noise and activities to improve patients' sleep quality in a surgical intensive care unit. *J Clin Nurs*. 2011;20(3–4):396–407.
- [89] Litton E, Elliott R, Thompson K, Watts N, Seppelt I, Webb SAR. Using clinically accessible tools to measure sound levels and sleep disruption in the ICU: a prospective multicenter observational study. *Crit Care Med*. 2017;45(6):966–71.
- [90] Longley L, Simons T, Glanzer L, Du C, Trinks H, Letzkus L, et al. Evaluating sleep in a surgical trauma burn intensive care unit: an elusive dilemma. *Dimens Crit Care Nurs*. 2018;37(2):97–101.
- [91] Louis M, Treger K, Ashby T, Smotherman C, Gautum S, Seeram V, et al. Patient-related factors may influence nursing perception of sleep in the intensive care unit. *PLoS One*. 2020;15(1).
- [92] Louzon P, Andrews J, Torres X, Pyles E, Ali M, Du Y, et al. Characterisation of ICU sleep by a commercially available activity tracker and its agreement with patient-perceived sleep quality. *BMJ Open Respir Res*. 2020;7(1).
- [93] McKinley S, Fien M, Elliott R, Elliott D. Sleep and psychological health during early recovery from critical illness: an observational study. *J Psychosom Res*. 2013;75(6):539–45.
- [94] Meneer A, Elliott R, L MA, Lal S, McKinley S. Repeated sleep-quality assessment and use of sleep-promoting interventions in ICU. *Nurs Crit Care*. 2017;22(6):348–54.
- [95] Mistraletti G, Umbrello M, Sabbatini G, Miori S, Taverna M, Cerri B, et al. Melatonin reduces the need for sedation in ICU patients: a randomized controlled trial. *Minerva Anestesiol*. 2015;81(12):1298–310.
- [96] Murata H, Oono Y, Sanui M, Saito K, Yamaguchi Y, Takinami M, et al. The Japanese version of the Richards-Campbell sleep questionnaire: reliability and validity assessment. *Nurs Open*. 2019;6(3):808–14.
- [97] Nagatomo K, Masuyama T, Iizuka Y, Makino J, Shiotsuka J, Sanui M. Validity of an under-mattress sensor for objective sleep measurement in critically ill patients: a prospective observational study. *J Intensive Care*. 2020;8(1).
- [98] Naik RD, Gupta K, Soneja M, Elavarasi A, Sreenivas V, Sinha S. Sleep quality and quantity in intensive care unit patients: a cross-sectional study. *Indian J Crit Care Med*. 2018;22(6):408–14.
- [99] Navarro-García MÁ, de Carlos Alegre V, Martínez-Oroz A, Irigoyen-Aristorena MI, Elizondo-Sotro A, Indurain-Fernández S, et al. Calidad del sueño en pacientes sometidos a cirugía cardíaca durante el postoperatorio en cuidados intensivos. *Enferm Intensiva*. 2017;28(3):114–24.
- [100] Nelson JE, Meier DE, Oei EJ, Nierman DM, Senzel RS, Manfredi PL, et al. Self-reported symptom experience of critically ill cancer patients receiving intensive care. *Crit Care Med*. 2001;29(2):277–82.
- [101] Nicolás A, Aizpitarte E, Iruarrizaga A. Perception of night-time sleep by surgical patients in an intensive care unit. *Nursing in critical care*. 2008;13(1):25–33.
- [102] Novaes MA, Aronovich A, Ferraz MB, Knobel E. Stressors in ICU: patients' evaluation. *Intensive Care Med*. 1997;23(12):1282–5.
- [103] Oto J, Yamamoto K, Koike S, Imanaka H, Nishimura M. Effect of daily sedative interruption on sleep stages of mechanically ventilated patients receiving midazolam by infusion. *Anaesth Intensive Care*. 2011;39(3):392–400.
- [104] Oto J, Yamamoto K, Koike S, Onodera M, Imanaka H, Nishimura M. Sleep quality of mechanically ventilated patients sedated with dexmedetomidine. *Intensive Care Med*. 2012;38(12):1982–9.
- [105] Papatheanassoglou EDE, Hadjilalasi M, Miltiadous P, Lambrinou E, Papastavrou E, Paikousis L, et al. Effects of an integrative nursing intervention on pain in critically ill patients: a pilot clinical trial. *Am J Crit Care*. 2018;27(3):172–85.
- [106] Parthasarathy S, Tobin MJ. Effect of ventilator mode on sleep quality in critically ill patients. *Am J Respir Crit Care Med*. 2002;166(11):1423–9.
- [107] Richardson A, Allsop M, Coghil E, Turnock C. Earplugs and eye masks: do they improve critical care patients' sleep? *Nurs Crit Care*. 2007;12(6):278–86.
- [108] Richardson S. Effects of relaxation and imagery on the sleep of critically ill adults. *Dimens Crit Care Nurs*. 2003;22(4):182–90.
- [109] Ritmala-Castren M, Virtanen I, Leivo S, Kaukonen KM, Leino-Kilpi H. Sleep and nursing care activities in an intensive care unit. *Nurs Health Sci*. 2015;17(3):354–61.
- [110] Roche-Campo F, Thille AW, Drouot X, Galia F, Margarit L, Córdoba-Izquierdo A, et al. Comparison of sleep quality with mechanical versus spontaneous ventilation during weaning of critically ill tracheostomized patients. *Crit Care Med*. 2013;41(7):1637–44.
- [111] Romagnoli S, Villa G, Fontanarosa L, Tofani L, Pinelli F, De Gaudio AR, et al. Sleep duration and architecture in non-intubated intensive care unit patients: an observational study. *Sleep Med*. 2020;70:79–87.
- [112] Rompaey BV, Elseviers MM. The effect of earplugs during the night on the onset of delirium and sleep perception: a randomized controlled trial in intensive care patients. [ccforum.biomedcentral.com](https://www.ccforum.biomedcentral.com/); 2012.
- [113] Shilo L, Dagan Y, Smorjick Y, Weinberg U, Dolev S, Komptel B, et al. Patients in the intensive care unit suffer from severe lack of sleep associated with loss of normal melatonin secretion pattern. *Am J Med Sci*. 1999;317(5):278–81.
- [114] Simons KS, Verweij E, Lemmens PMC, Jelfs S, Park M, Spronk PE, et al. Noise in the intensive care unit and its influence on sleep quality: a multicenter observational study in Dutch intensive care units. *Crit Care*. 2018;22(1).
- [115] Stewart JA, Green C, Stewart J, Tiruvoipati R. Factors influencing quality of sleep among non-mechanically ventilated patients in the intensive care unit. *Aust Crit Care*. 2017;30(2):85–90.
- [116] Su CP, Lai HL, Chang ET, Yiin LM. ... randomized controlled trial of the effects of listening to non-commercial music on quality of nocturnal sleep and relaxation indices in patients in medical intensive care. *J Adv*. 2013;69(6):1377–89.
- [117] Thille AW, Reynaud F, Marie D, Barrau S, Rousseau L, Rault C, et al. Impact of sleep alterations on weaning duration in mechanically ventilated patients: a prospective study. *Eur Respir J*. 2018;51(4).
- [118] Toublanc B, Rose D, Gléran JC, Francois G, Mayeux I, Rodenstein D, et al. Assist-control ventilation vs. low levels of pressure support ventilation on sleep quality in intubated ICU patients. *Intensive Care Med*. 2007;33(7):1148–54.
- [119] Uğraş GA, Babayigit S, Tosun K, Aksog Y, Turan Y. The effect of nocturnal patient care interventions on patient sleep and satisfaction with nursing care in neurosurgery intensive care unit. *J Neurosci Nurs*. 2015;47(2):104–12.
- [120] Uğraş GA, Öztekin SD. Patient perception of environmental and nursing factors contributing to sleep disturbances in a neurosurgical intensive care unit. *Tohoku J Exp Med*. 2007;212(3):299–308.
- [121] Vacas S, McInrue E, Gropper MA, Maze M, Zak R, Lim E, et al. The feasibility and utility of continuous sleep monitoring in critically ill patients using a portable electroencephalography monitor. *Anesth Analg*. 2016;123(1):206–12.
- [122] van de Pol I, van Itersson M, Maaskant J. Effect of nocturnal sound reduction on the incidence of delirium in intensive care unit patients: an interrupted time series analysis. *Intensive Crit Care Nurs*. 2017;41:18–25.
- [123] Wallace CJ. The effect of earplugs on polysomnographic sleep and description of sleep-disturbing factors in critically ill subjects. The University of Utah; 1998.
- [124] Watson PL, Pandharipande P, Gehlbach BK, Thompson JL, Shintani AK, Dittus BS, et al. Atypical sleep in ventilated patients: empirical electroencephalography findings and the path toward revised ICU sleep scoring criteria. *Crit Care Med*. 2013;41(8):1958–67.
- [125] Wu XH, Cui F, Zhang C, Meng ZT, Wang DX, Ma J, et al. Low-dose dexmedetomidine improves sleep quality pattern in elderly patients after noncardiac surgery in the intensive care unit: a pilot randomized controlled trial. *Anesthesiology*. 2016;125(5):979–91.
- [126] Yazdankar AR, Zareie A, Hasanpour M, Kashefi P. The effect of earplugs and eye mask on patients' perceived sleep quality in intensive care unit. *Iran J Nurs Midwifery Res*. 2014;19(6):673–8.
- [127] Zhang L, Sha YS, Kong QQ, Woo JAL, Miller AR, Li HW, et al. Factors that affect sleep quality: perceptions made by patients in the intensive care unit after thoracic surgery. *Support Care Cancer*. 2013;21(8):2091–6.
- [128] Bellapart J, Appadurai V, Lassig-Smith M, Stuart J, Zappala C, Boots R. Effect of exogenous melatonin Administration in Critically ill Patients on delirium and sleep: a randomized controlled trial. *Crit Care Res Pract*. 2020;2020:3951828.
- [129] Sun T, Sun Y, Huang X, Liu J, Yang J, Zhang K, et al. Sleep and circadian rhythm disturbances in intensive care unit (ICU)-acquired delirium: a case-control study. *J Int Med Res*. 2021;49(3).
- [130] Chen Q, Peng Y, Lin Y, Li S, Huang X, Chen LW. Atypical sleep and postoperative delirium in the cardiothoracic surgical intensive care unit: a pilot prospective study. *Nat Sci Sleep*. 2020;12:1137–44.
- [131] Thille AW, Barrau S, Beuvon C, Marie D, Reynaud F, Bardin J, et al. Role of sleep on respiratory failure after extubation in the ICU. *Ann Intensive Care*. 2021;11(1).
- [132] Aydın Sayılan A, Kulakaç N, Sayılan S. The effects of noise levels on pain, anxiety, and sleep in patients. *Nurs Crit Care*. 2021;26(2):79–85.
- [133] Andrews JL, Louzon PR, Torres X, Pyles E, Ali MH, Du Y, et al. Impact of a pharmacist-led intensive care unit sleep improvement protocol on sleep duration and quality. *Ann Pharmacother*. 2021;55(7):863–9.
- [134] Biazim SK, Souza DA, Junior HC. The Richards-Campbell sleep questionnaire and sleep in the intensive care unit questionnaire: translation to Portuguese and cross-cultural adaptation for ... *SciELO. Brasil*. 2020;46(4):e20180237.
- [135] Gandolfi JV, Bernardo APAD, Chanes DAV, Martin DF, Joles VB, Amendola CP, et al. The effects of melatonin supplementation on sleep quality and assessment of the serum melatonin in ICU patients: a randomized controlled trial. *Crit Care Med*. 2020;48(12):E1286–93.
- [136] Kim J, Choi D, Yeo MS, Yoo GE, Kim SJ, Na S. Effects of patient-directed interactive music therapy on sleep quality in postoperative elderly patients: a randomized-controlled trial. *Nat Sci Sleep*. 2020;12:791–800.
- [137] Kim JK, Park JH, Cho J, Lee SM, Lee J. Reliability of the Korean version of the Richards-Campbell sleep questionnaire. *Acute Crit Care*. 2020;35(3):164–8.
- [138] Koçak AT, Arslan S. The effect of using eye masks and earplugs on intensive care patients quality of sleep and vital signs. *J Neurosci Nurs*. 2021;53(1):29–33.



- [139] Lochová H, Axmann K, Žiaková K, Šerková D. Effect of a multicomponent sleep protocol on sleep quality in conscious patients in the intensive care unit. *J Turk Sleep Med*. 2020;7(3):140–8.
- [140] Miranda-Ackerman RC, Lira-Trujillo M, Gollaz-Cervantez AC, Cortés-Flores AO, Zuloaga-Fernández Del Valle CJ, García-González LA, et al. Associations between stressors and difficulty sleeping in critically ill patients admitted to the intensive care unit: a cohort study. *BMC Health Serv Res*. 2020;20(1):631.
- [141] Mori C, Boss K, Indermuhle P, Stahl E, Chiu SH, Shanks L. Is it noise? Factors linked with sleep interruption in hospitalized patients. *Clin Nurse Spec*. 2021;35(4):199–207.
- [142] Obanor OO, McBroom MM, Elia JM, Ahmed F, Sasaki JD, Murphy KM, et al. The impact of earplugs and eye masks on sleep quality in surgical ICU patients at risk for frequent awakenings. *Crit Care Med*. 2021;49(9):822–32.
- [143] Ritmala-Castren M, Axelin A, Richards KC, Mitchell ML, Vahlberg T, Leino-Kilpi H. Investigating the construct and concurrent validity of the Richards-Campbell sleep questionnaire with intensive care unit patients and home sleepers. *Aust Crit Care*. 2021;35(2):130–5.
- [144] Tonna JE, Dalton A, Presson AP, Zhang C, Colantuoni E, Lander K, et al. The effect of a quality improvement intervention on sleep and delirium in critically ill patients in a surgical intensive care unit. *Chest*. 2021;160:899–908.
- [145] Arita H, Takeno Y, Fujimoto E. Sleep disturbance in elderly patients after cardiac surgery during their stay in intensive care unit and surgical ward: pdfs.semanticscholar.org; 2014..
- [146] Munro CL, Liang Z, Elías MN, Ji M, Chen X, Calero K. Sleep and activity patterns are altered during early critical illness in mechanically ventilated adults. *Dimens Crit Care Nurs*. 2021;40(1):29–35.
- [147] Pedrão RAA, Riella RJ, Richards K, Valderramas SR. Viability and validity of the bispectral index to measure sleep in patients in the intensive care unit. *Rev Bras Ter Intensiva*. 2020;32(4):535–41.
- [148] Chaudhary A, Kumari V, Neetu N. Sleep promotion among critically ill patients: earplugs/eye mask versus ocean sound-a randomized controlled trial study. *Crit Care Res Pract*. 2020;2020:8898172.
- [149] Bode L, Fuchs S, Gehrke S, Ernst J, Schubert M, Nuñez DG, et al. Assessing delirium with nursing care instruments: evaluation of the cognitive and associated domains. *Palliat Support Care*. 2021;19(1):11–6.
- [150] Alinia-najjar R, Bagheri-Nesami M, Shorofi SA, Mousavinasab SN, Saatchi K. The effect of foot reflexology massage on burn-specific pain anxiety and sleep quality and quantity of patients hospitalized in the burn intensive care unit (ICU). *Burns*. 2020;46(8):1942–51.
- [151] Zhang Y, Li X, Chen S. Bilateral transversus thoracis muscle plane block provides effective analgesia and enhances recovery after open cardiac surgery. *J Card Surg*. 2021;36:2818–23.
- [152] Zhang Y, Wu Y, Xu D, Xiao P, Xie B, Huang H, et al. Very-short-term sleep deprivation slows early recovery of lymphocytes in septic patients. *Front Med (Lausanne)*. 2021;8:656615.
- [153] Lee YJ, Lee J, Lee SM, Cho J. Postextubation respiratory events in patients admitted to the intensive care unit: a prospective pilot study using overnight respiratory polygraphy. *Acute Crit Care*. 2020;35(4):271–8.
- [154] Delaney LJ, Litton E, Melehan KL, Huang HC, Lopez V, Van Haren F. The feasibility and reliability of actigraphy to monitor sleep in intensive care patients: an observational study. *Crit Care*. 2021;25(1):42.
- [156] Snyder-Halpern R, Verran JA. Instrumentation to describe subjective sleep characteristics in healthy subjects. *Res Nurs Health*. 1987;10(3):155–63.
- [157] Richards KC, O'Sullivan PS, Phillips RL. Measurement of sleep in critically ill patients. *J Nurs Meas*. 2000;8(2):131–44.
- [158] Richards K. Techniques for measurement of sleep in critical care. *Focus Crit Care*. 1987;14(4):34–40.
- [159] Boyko Y, Jennum P, Toft P. Sleep quality and circadian rhythm disruption in the intensive care unit: a review. *Nat Sci Sleep*. 2017;9:277–84.
- [160] Schwab KE, Ronish B, Needham DM, To AQ, Martin JL, Kamdar BB. Actigraphy to evaluate sleep in the intensive care unit. A systematic review. *Ann Am Thorac Soc*. 2018;15(9):1075–82.
- [161] Kamdar BB, Shah PA, King LM, Kho ME, Zhou X, Colantuoni E, et al. Patient-nurse interrater reliability and agreement of the Richards-Campbell sleep questionnaire. *Am J Crit Care*. 2012;21(4):261–9.
- [162] Siegel JM. Principles and practice of sleep medicine. Elsevier; 2017; 78–95.
- [163] Singer HS. Movement disorders in childhood. Elsevier; 2010; 205–18.
- [164] Wesselius HM, van den Ende ES, Alsmas J, Ter Maaten JC, Schuit SCE, Stassen PM, et al. Quality and quantity of sleep and factors associated with sleep disturbance in hospitalized patients. *JAMA Intern Med*. 2018;178(9):1201–8.