

Endoscopic embolization of a refractory bronchobiliary fistula by endoscopic retrograde cholangiography using coils and Histoacryl



Bronchobiliary fistulae are often related to hepatic tumors and also occur after ablative therapy [1,2]. Endoscopic sphincterotomy or stenting is considered first-line treatment. In case of refractory fistulas, sealing with glue or coils has been suggested, however data are scarce [3,4].

We present a 53-year-old woman who underwent locoregional treatments of colorectal liver metastasis, including segment 2/3 and wedge resections, radiofrequency ablation (segment 8), and radio-embolization. Plastic stents (after sphincterotomy) were previously inserted to treat sclerosis of both hepatic ducts related to intra-arterial chemotherapy. She was referred for endoscopic management of a refractory bronchobiliary fistula, resulting in biliptysis.

Endoscopic retrograde cholangiography (ERC) confirmed leakage from a peripheral bile duct in segment 8 towards the bronchial tree (► **Video 1**, ► **Fig. 1**). A 4-Fr tapered Glo-tip catheter (Cook Medical, Bloomington, Indiana, USA) was advanced just underneath the fistula. Two coils (Tornado 0.035 inch, 4–3 mm, and MRey 0.035 inch, 5 mm; Cook Medical) were deployed at the fistula by pushing them through and out the catheter with a standard 0.035-inch guidewire. Two plastic stents were re-inserted over the persistent hepatic duct strictures.

After initial resolution, biliptysis recurred after two weeks and a second ERC was performed showing persistent leakage alongside the coils. A cannula (Tandem XL; Boston Scientific, Marlborough, Massachusetts, USA) was advanced and was flushed with a 5% glucose solution. N-butyl-2-cyanoacrylate (Histoacryl; B. Braun, Melsungen, Germany) was diluted with Lipiodol (Guerbet GmbH, Sulzbach, Germany) (0.5 ml: 0.5 ml). The glue mixture was injected slowly by pushing it with a 5% glucose solution. The catheter was pulled back under fluoroscopic guidance during injection. An elongated glue



► **Video 1** Sealing of a bronchobiliary fistula by endoscopic retrograde cholangiography using coils and Histoacryl.



► **Fig. 1** Cholangiography demonstrating bronchobiliary fistula.



► **Fig. 2** Presence of two coils (blue arrow) and glue cast (red arrow) at the fistula and in the small peripheral bile duct.

cast was seen after injection of 0.3 cc around the coils and in the end of the peripheral bile duct. Two plastic stents were repositioned (► **Fig. 2**). No complications occurred and the patient remained symptom-free during 2 months' follow-up.

Endoscopy_UCTN_Code_TTT_1AR_2AG

Competing interests

The authors declare that they have no conflict of interest.

The authors

Paul Didden¹, Rutger C. G. Bruijnen², Evert-Jan P. A. Vonken², Frank P. Vleggaar¹

- 1 University Medical Center Utrecht, Department of Gastroenterology and Hepatology, Utrecht, Netherlands
- 2 University Medical Center Utrecht, Department of Radiology, Utrecht, Netherlands

Corresponding author

Paul Didden, MD

University Medical Center Utrecht,
Department of Gastroenterology and
Hepatology, Heidelberglaan 100, 3584 CX
Utrecht, Netherlands
pdidden@umcutrecht.nl

References

- [1] Liao GQ, Wang H, Zhu GY et al. Management of acquired bronchobiliary fistula: A systematic literature review of 68 cases published in 30 years. *World J Gastroenterol* 2011; 33: 3842–3849
- [2] Huang ZM, Zuo MX, Gu YK et al. Broncho-biliary fistula after ablation of hepatocellular carcinoma adjacent to the diaphragm: Case report and literature review. *Thorac Cancer* 2020; 5: 1233–1238
- [3] Suzuki E, Fujita Y, Matsushashi N. Case of bile duct coil embolization for broncho-biliary fistula after radiofrequency ablation (with video). *Dig Endosc* 2021; 4: e79–e80
- [4] Seewald S, Groth S, Sriram PV et al. Endoscopic treatment of biliary leakage with n-butyl-2 cyanoacrylate. *Gastrointest Endosc* 2002; 6: 916–919

Bibliography

Endoscopy 2023; 55: E268–E269
DOI 10.1055/a-1974-9202
ISSN 0013-726X
published online 25.11.2022
© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>