

Patients With and Without COVID-19 in the Intensive Care Unit: Physical Status Outcome Comparisons 3 Months After Discharge

Bastiaan Cijis, PT, MSc^{1,*}, Karin Valkenet, PT, PhD¹, Germijn Heijnen¹,
J.M. Anne Visser-Meily, MD¹, Marike van der Schaaf, PT, PhD^{2,3,4}

¹Department of Rehabilitation, Physical Therapy Science & Sports, University Medical Center Utrecht, Utrecht University, UMC Utrecht Brain Center, Utrecht, the Netherlands

²Department of Rehabilitation Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam, the Netherlands

³Amsterdam Movement Sciences, Ageing and Vitality, Amsterdam, the Netherlands

⁴Centre of Expertise Urban Vitality, Faculty of Health, Amsterdam University of Applied Sciences, Amsterdam, the Netherlands

*Address all correspondence to Mr Cijis at: bas.cijis@hu.nl

Abstract

Objective. Many patients with coronavirus disease 2019 (COVID-19) infections were admitted to an intensive care unit (ICU). Physical impairments are common after ICU stays and are associated with clinical and patient characteristics. To date, it is unknown if physical functioning and health status are comparable between patients in the ICU with COVID-19 and patients in the ICU without COVID-19 3 months after ICU discharge. The primary objective of this study was to compare handgrip strength, physical functioning, and health status between patients in the ICU with COVID-19 and patients in the ICU without COVID-19 3 months after ICU discharge. The second objective was to identify factors associated with physical functioning and health status in patients in the ICU with COVID-19.

Methods. In this observational, retrospective chart review study, handgrip strength (handheld dynamometer), physical functioning (Patient-Reported Outcomes Measurement Information System Physical Function), and health status (EuroQol 5 Dimension 5 Level) were compared between patients in the ICU with COVID-19 and patients in the ICU without COVID-19 using linear regression. Multilinear regression analyses were used to investigate whether age, sex, body mass index, comorbidities in medical history (Charlson Comorbidity Index), and pre-morbid function illness (Identification of Seniors At Risk-Hospitalized Patients) were associated with these parameters in patients in the ICU with COVID-19.

Results. In total, 183 patients ($N = 92$ with COVID-19) were included. No significant between-group differences were found in handgrip strength, physical functioning, and health status 3 months after ICU discharge. The multilinear regression analyses showed a significant association between sex and physical functioning in the COVID-19 group, with better physical functioning in men compared with women.

Conclusion. Current findings suggest that handgrip strength, physical functioning, and health status are comparable for patients who were in the ICU with COVID-19 and patients who were in the ICU without COVID-19 3 months after ICU discharge.

Impact. Aftercare in primary or secondary care in the physical domain of postintensive care syndrome after ICU discharge in patients with COVID-19 and in patients without COVID-19 who had an ICU length of stay >48 hours is recommended.

Lay Summary. Patients who were in the ICU with and without COVID-19 had a lower physical status and health status than healthy people, thus requiring personalized physical rehabilitation. Outpatient aftercare is recommended for patients with an ICU length of stay >48 hours, and functional assessment is recommended 3 months after hospital discharge.

Keywords: COVID-19, Health Status, Intensive Care Unit, Physical Functioning, Physical Therapy, Rehabilitation

Received: May 12, 2022. **Revised:** December 16, 2022. **Accepted:** February 15, 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of the American Physical Therapy Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Introduction

In December 2019, a new coronavirus, called coronavirus disease 2019 (COVID-19), was identified in Wuhan, China. COVID-19 causes severe respiratory illness and is associated with intensive care unit (ICU) admission and high mortality.¹ The Netherlands experienced a multiple wave pattern in reported cases of COVID-19.² A systematic review including studies in multiple countries showed that 26% of patients hospitalized with a COVID-19 infection were admitted to the ICU.³ Due to novelty of the new COVID-19 infection, little is known about the functional status of these patients several months after ICU discharge. On other timepoints, it is known that patients with COVID-19 who had a pneumosepsis perform worse on physical assessments on ICU discharge compared with patients without COVID-19 who had a pneumosepsis.⁴ Furthermore, most patients with COVID-19 in the ICU have lower physical functioning than the general population 6 months after ICU discharge.⁵ Moreover, at 6 months after ICU discharge, COVID-19-related ICU stays are not associated with a lower health-related quality of life compared with non-COVID-19-related ICU stays.⁶

Approximately half of the patients without COVID-19 in the ICU develop the post-intensive care syndrome (PICS). PICS includes new or worsening physical, cognitive, and/or mental health impairments arising after critical illness, persisting beyond ICU care.^{7,8} Known possible predictive variables for poor physical recovery after ICU stay are older age, female sex (reported as male/female), a high body mass index (BMI), presence of comorbidities, a low self-reported pre-morbid function, and a higher severity of illness.^{9,10} Rehabilitation seems to be a cornerstone in the management of PICS during all phases of recovery, starting at the ICU admission,¹¹ continuing on the general hospital ward, and after hospital discharge.¹² However, a Cochrane review, focusing on rehabilitation after hospital discharge, did not show an effect on physical functioning due to low study quality.¹³

Patient and clinical characteristics associated with physical impairments in patients without COVID-19 in the ICU might be different in patients with COVID-19 in the ICU, as the group of patients without COVID-19 in the ICU show a large heterogeneity with respect to diagnosis, age, comorbidities, and ICU treatment when compared with patients with a COVID-19 infection in the ICU.^{14–16}

Currently, it remains unknown whether outcomes of physical status are comparable between patients with a COVID-19 infection and patients without COVID-19 several months after ICU discharge. Therefore, the first aim of this study is to compare physical status outcomes (handgrip strength, physical functioning, and health status) between these 2 groups 3 months after ICU discharge. The second aim is to investigate which demographic and patient characteristics are associated with the physical status outcomes in patients who had COVID-19 in the ICU. Gaining a better understanding about physical status might yield possibilities for physical therapy.

Methods

This observational, retrospective chart review used usual care data from the outpatient ICU aftercare clinic of the University Medical Center Utrecht. In outpatient ICU aftercare 3 months after ICU discharge, a physician assistant and a rehabilitation specialist assess physical, mental, and cognitive functioning

and participation goals and, when necessary, provide advice for improving recovery. In this cross-sectional study, participants were included if they were 18 years old or older at ICU admission and visited the outpatient ICU aftercare of University Medical Center Utrecht between January 2020 and June 2021. Exclusion criteria were age \geq 80 years, ICU length of stay $<$ 48 hours, neurological disease as ICU admission diagnosis, or ICU discharge with home-based mechanical ventilation. The data were obtained by the research group from the electronic health record programs HiX (HiX Digital Health Services, Amsterdam, the Netherlands) and Metavision (Prophesee Metavision, Paris, France), pseudonymized, and entered into an electronic data collection application (CASTOR EDC, New York, NY, USA).

Measurements

During the COVID-19 crisis, The Netherlands Society of Rehabilitation Medicine has recommended a set of measurement instruments to use in outpatient ICU aftercare.¹⁷ In this study, data from measurements and questionnaires related to handgrip strength, physical functioning, and health status were used. Handgrip strength was measured with a JAMAR handheld dynamometer.¹⁸ Handheld dynamometry has good validity in patients in the ICU and good interrater reliability for measuring handgrip strength.¹⁹ The raw outcome values of the JAMAR handheld dynamometer were converted to a percentage of normative references by sex, age, and body height.²⁰ This percentage was used in the analyses of this study. Physical functioning was measured with the Dutch Short-Form Patient-Reported Outcomes Measurement Information System Physical Function (PROMIS-PF), a questionnaire measuring patient-reported physical functioning on a 5-point scale.²¹ The total scores on the PROMIS-PF were transformed into normalized *t* values and used in the analyses of this study.^{21,22} The validity and reliability of the PROMIS-PF are unknown for patients in the ICU; however, the PROMIS-PF is known to be sensitive to change in intervention studies on patients with chronic heart failure and cancer.²³ Furthermore, the PROMIS-PF is sensitive to change among different clinical samples.²³ Health status was measured with EuroQol 5 Dimensions 5 Levels (EQ-5D-5L), a questionnaire assessing mobility, self-care, daily activities, pain, and anxiety/depression.²⁴ Outcomes were transformed into an EQ-5D-5L index value and used in the analyses of this study. The EQ-5D-5L index is a valid extension of the EQ-5D-3L with improved discriminatory power and known-groups validity^{25,26} and has shown to be a valid and responsive measure of health status in patients with COPD and asthma, patients after stroke, and patients with chronic hepatic diseases.^{26–29} All outcome variables are continuous data.

For the secondary study aim, the following potential determinants for postdischarge physical status of the COVID-19 ICU group were selected based on earlier studies: age^{9,10,30} (continuous), sex^{10,30} (reported as male/female), BMI,^{10,31} comorbidities in medical history,^{9,10,30} and pre-morbid function^{9,10} at ICU admission. BMI was calculated according to the continuous variable Quetelet Index.³² The presence of comorbidities in medical history was measured by the continuous Charlson Comorbidity Index (CCI).³³ The CCI quantifies an individual's burden of disease and corresponding 1-year mortality risk. Pre-morbid function was measured by the ordinal variable Identification of Seniors At Risk-Hospitalized Patients (ISAR-HP).³⁴ The ISAR-HP is a screening instrument

to predict 90-day functional decline in older patients who had an acute care admission to the department of internal medicine.³⁴

Finally, patient characteristics and clinical data associated with the ICU stay were recorded to describe and compare the group of patients in the ICU without COVID-19 and the group of patients in the ICU with COVID-19. The severity of illness was scored with the Acute Physiology And Chronic Health Evaluation (APACHE) IV score.¹⁰ The APACHE IV score is a method for predicting hospital mortality among adults who are critically ill. Furthermore, the duration of mechanical ventilation,^{9,35} ICU length of stay, and hospital length of stay³⁵ were recorded in days, and organ dysfunction was measured with the Sequential Organ Failure Assessment score.^{36,37} Furthermore, the presence of sepsis was indicated by the highest C-reactive protein level (CRP). A CRP level > 50 ng/ml indicates sepsis.^{9,38} Finally, the presence of hyperglycemia was recorded, indicated by the blood glucose level during ICU stay. A blood glucose level > 150 mg/dl maintained for >5 days was defined as hyperglycemia.³⁹

Results from the JAMAR handheld dynamometer measurements, the PROMIS-PF, the EQ-5D-5L, the CCI, the CRP level, and the glucose level were extracted manually from the electronic health records of participants and entered into an electronic data capturing tool. Other study parameters were automatically extracted and imported into the electronic data capturing tool. To prevent bias due to input errors, a second researcher randomly checked 5% of the entered data. Pseudonymized data from the electronic data collection application were exported into SPSS version 27 (SPSS Inc, Chicago, IL, USA) for analyses.⁴⁰

Ethical Approval

Ethical approval for the usage of daily care data was obtained from the Medical Ethical Research Committee of University Medical Center Utrecht. Data from patients who signed an objection letter for the use of their health care data in research were excluded. The study complies with the Declaration of Helsinki.

Statistical Analyses

Descriptive statistics were performed to summarize demographic and clinical characteristics. These characteristics were compared between the non-COVID-19 ICU group and the COVID-19 ICU group by using univariate linear regression analyses.

To conduct the main analyses with complete data, multiple imputation was performed for missing data. For missing data qualified as missing completely at random and missing at random, 5 imputations were obtained. This gives an efficiency of 99% compared with using an infinite number of imputations.⁴¹ Consistent with the guidelines for variable selection for multiple imputation, patient characteristics were included in the multiple imputation procedure.⁴² Imputed variables and variables that were used as a predictor for imputation are described in the [Supplementary Appendix](#).

Primarily, handgrip strength, physical functioning, and health status of patients in the ICU without COVID-19 and patients in the ICU with COVID-19 were compared—as well as demographic and clinical characteristics—using univariate linear regression analyses. Subsequently, these comparisons were repeated while correcting for time between ICU discharge and visitation of outpatient ICU aftercare

using multivariate linear regression analyses. Furthermore, the mean handgrip strength, physical functioning, and health status of the non-COVID-19 ICU group and the COVID-19 ICU group were compared with reference values of healthy persons.^{20,21,43}

Secondarily, to investigate which factors were associated with the dependent variables of handgrip strength, physical functioning, and health status in patients in the ICU with COVID-19, multilinear regression analyses were used, with age, sex, BMI, number of comorbidities in medical history, and premorbid function illness as independent variables.^{9,27–29} Multicollinearity was checked for independent variables.

Furthermore, 2 post hoc analyses were performed. First, the multilinear regression analysis was repeated and supplemented with the variables that showed to be significantly different between the 2 groups in [Table 1](#) as independent variables, to correct for differences at baseline. Second, the Netherlands experienced a multiple wave pattern in reported cases of COVID-19. During our inclusion period, the first 3 COVID-19 waves occurred. As the medical treatment of patients with COVID-19 in the ICU evolved, and as the proportion of hospitalized patients requiring ICU treatment and mechanical ventilation dropped, subgroup analyses per wave of reported cases were performed. Therefore, the COVID-19 ICU group was divided into 3 groups, depending on the time of ICU admission, namely, in the first wave (March 2020–June 2020), second wave (July 2020–January 2021), and third wave (since February 2021).² The total group of patients in the ICU without COVID-19 and each subgroup of patients in the ICU with COVID-19 were compared on handgrip strength, physical functioning, and health status using univariate linear regression analyses.

All analyses were conducted with SPSS 27 (IBM, Armonk, NY, USA). Statistical assumptions were examined prior to the analyses. Statistical results at level $P \leq .05$ were considered to be significant.

Sample Size Calculation

To determine the minimal sample size for the primary analyses, the effect size was set at 0.5, indicating a medium effect size. This effect size was chosen because there are no sufficient data available to calculate an effect size.⁴⁴ Based on a 2-tailed outcome, an α error probability of 0.05 and a power of 0.80 with an equal allocation ratio were set leading to a minimal sample size of 128 (64 per group). For the secondary analyses, the rule of thumb of a maximum of 1 independent variable per 10 events was followed.⁴⁵

Results

We studied 183 patients (57 women) with a mean (SD) age of 57.6 (13.3) years. In total, 168 patients had a face-to-face consult at the outpatient ICU aftercare and, in the non-COVID-19 ICU group, 15 patients had an assessment conducted by phone due to the COVID-19 crisis. The COVID-19 ICU group consisted of 92 patients (27 women) with a mean (SD) age of 60.8 (10.5) years, and the non-COVID-19 ICU group consisted of 91 patients (30 women) with a mean (SD) age of 54.2 (15.1) years ($P = <.001$). The time between ICU discharge and visitation at the outpatient ICU aftercare appointment differed between the COVID-19 ICU group (mean [SD] = 84.6 [30.1] days) and the non-COVID-

Table 1. Sample Characteristics (N = 183)^a

Characteristic	Total Group (N = 183)		COVID-19 ICU Group (n = 92)		Non-COVID-19 ICU Group (n = 91)		P ^b
	No. (%)	Mean (SD)	No. (%)	Mean (SD)	No. (%)	Mean (SD)	
Sex, women	57 (31.1)		27 (29.3)		30 (33.0)		.59
Age	182	57.6 (13.3)	92	60.8 (10.5)	90	54.2 (15.1)	<.001
Body mass index	182	27.3 (5.6)	91	28.0 (5.9)	91	26.5 (5.3)	.08
Embolism			32 (34.8)				
Diagnosis type					91		
Cardiac					11 (12.1)		
Pulmonary					11 (12.1)		
Neurologic					8 (8.8)		
Surgical					52 (57.1)		
Orthopedic					1 (1.1)		
Internal					6 (6.6)		
Gastroenterologic					2 (2.2)		
Oncologic					0 (0.0)		
Surgical type					52		
Cardiac					33 (63.5)		
Pulmonary					5 (9.6)		
Neurologic					2 (3.8)		
Orthopedic					3 (5.8)		
Internal					2 (3.8)		
Gastroenterologic					1 (1.9)		
Oncologic					6 (11.5)		
Physical consult at ICU aftercare	168 (91.8)		91 (98.9)		77 (84.6)		<.001
Time between ICU discharge and ICU aftercare, d	175	118.1 (70.4)	87	84.6 (30.1)	88	151.1 (82.4)	<.001
Premorbid functioning ^c	113	0.6 (0.1)	55	0.5 (0.9)	58	0.7 (1.1)	.54
Comorbidities ^d	181	2.6 (2.3)	91	2.6 (2.1)	90	2.6 (2.5)	.99
Hospital stay, d	154	32.5 (21.8)	74	30.3 (15.9)	80	34.5 (26.0)	.36
ICU stay, d	174	15.4 (13.9)	87	18.7 (13.1)	87	12.1 (14.0)	<.001
Severity of illness ^e	166	66.9 (22.3)	86	66.3 (19.5)	80	67.6 (24.9)	.72
Readmission to ICU	23 (12.6)		6 (6.5)		17 (18.7)		.02
Mechanical ventilation	167 (91.3)		83 (90.2)		84 (92.3)		.60
Reintubation	21 (11.5)		5 (5.4)		16 (17.6)		.01
Mechanical ventilation, d	157	11.8 (11.7)	77	15.8 (11.3)	80	8.0 (10.8)	<.001
ECLS/ECMO	15 (8.2)		2 (2.2)		13 (14.3)		.003
ECLS/ECMO, d	15	10.1 (8.6)	2	13.0 (9.8)	13	9.7 (8.7)	.62
CRP level ^f	158	223.7 (115.1)	72	235.5 (121.3)	86	213.9 (109.4)	.33
Blood glucose level ^f	160	12.5 (3.8)	72	12.5 (4.0)	88	12.6 (3.7)	.99
Multiorgan failure ^g	177	9.4 (2.8)	91	9.2 (2.8)	86	9.7 (2.9)	.24
Rehabilitation	181		91		90		.07
None	25 (13.8)		8 (8.8)		17 (18.9)		
Primary care	61 (33.7)		35 (38.5)		26 (28.9)		
Secondary care	44 (24.3)		26 (28.6)		18 (20.0)		
Tertiary care	51 (28.1)		22 (24.2)		29 (32.2)		
Discharge destination	181		91		90		.03
Home	103 (56.9)		43 (47.3)		60 (66.7)		
Care hotel	2 (1.1)		2 (2.2)		0 (0.0)		
Secondary care	40 (22.1)		26 (28.6)		14 (15.6)		
Tertiary care	36 (19.9)		20 (22.0)		16 (17.8)		

^aCRP = C-reactive protein; ECLS = extracorporeal life support; ECMO = extracorporeal membrane oxygenation; ICU = intensive care unit. ^bValues in bold type were significant at $P < .05$. ^cDetermined with the Identification of Seniors at Risk–Hospitalized Patients instrument. ^dDetermined with the Charlson Comorbidity Index. ^eDetermined with the Acute Physiology and Chronic Health Evaluation IV. ^fHighest value during ICU stay. ^gHighest Sequential Organ Failure Assessment Score during ICU stay.

ICU group (mean [SD] = 151.1 [82.4] days) ($P = < .001$). Furthermore, patients in the COVID-19 ICU group were mechanically ventilated longer (15.8 days vs 8.0 days; $P = < .001$), and the use of extracorporeal life support/extracorporeal membrane oxygenation (ECLS/ECMO) was necessary less often (2.2% vs 14.3%; $P = < .003$). Patients in the non-COVID-19 ICU group were more often able to go home upon discharge (66.7%) than those in the COVID-19 ICU group (47.3%). In the COVID-19 group, 91.3% received some

form of rehabilitation (primary care, 38.5%; secondary care, 28.6%; tertiary care, 24.2%), whereas in the non-COVID-19 group, 81.1% received some form of rehabilitation (primary care, 28.9%; secondary care, 20.0%; tertiary care, 32.2%). Demographic details and patient characteristics are provided in Table 1.

There were missing data on primary parameters for hand-grip strength (37.7%), physical functioning (26.5%), and health status (32.2%) and on secondary parameters for age

Table 2. Comparison of Handgrip Strength, Physical Functioning, and Health Status Between the COVID-19 ICU Group and the Non-COVID-19 ICU Group ($N = 183$)^a

Parameter	COVID-19 ICU Group		Non-COVID 19 ICU Group		Unadjusted ^b			Adjusted ^c		
	No.	Mean (SEM)	No.	Mean (SEM)	B	95% CI	P^d	B	95% CI	P^d
Handgrip strength ^e	90	74.2 (2.93)	89	84.2 (4.44)	-10.03	-22.41 to 2.35	.10	-5.4	-17.41 to 6.54	.36
Physical functioning ^f	91	39.0 (0.99)	90	39.8 (0.99)	-0.82	-3.59 to 1.95	.56	-1.87	-5.07 to 1.34	.25
Health status ^g	91	0.7 (0.03)	90	0.7 (0.03)	-0.00	-0.09 to 0.11	.85	0.00	0.00 to 0.00	.85

^aICU = intensive care unit; SEM = standard error of the mean. ^bUnivariate linear regression. ^cMultivariate linear regression adjusted for time between ICU discharge and ICU aftercare. ^dSignificance was set at $P < .05$. ^eDetermined with a JAMAR handheld dynamometer and reported as percentages of reference values. ^fReported as a Patient-Reported Outcomes Measurement Information System physical function t value. ^gReported as a EuroQol 5 Dimensions 5 Levels index value.

(0.5%), BMI (0.5%), CCI (1%), and ISAR-HP (38.2%). All missing data were considered to be “missing completely at random,” with exception of the 14 patients who had an ICU aftercare appointment by phone; those missing data were qualified as “missing at random.” There were no missing values described as “missing not at random,” and therefore all missing data were eligible for multiple imputation.

Univariate linear regression analyses showed no significant differences in handgrip strength ($P = .104$), physical functioning ($P = .562$), and health status ($P = .848$) between the 2 groups. These findings remained unchanged after adjustment for the time between ICU discharge and the visitation at the ICU aftercare appointment (Tab. 2). The handgrip strength of the patients in the COVID-19 ICU group (74.2% of normative values) and non-COVID-19 ICU group (84.2% of normative values) was lower than healthy people.²⁰ The physical functioning of both groups was 1 SD below average,²¹ and the health status of both groups was lower than in healthy people (Fig. 1).⁴³

The multivariate linear regression analyses showed a significant ($P = .008$) association between sex and physical functioning in the COVID-19 ICU group, where men showed a higher level of physical functioning than women showed. Furthermore, a significant association ($P = .03$) between the number of comorbidities and physical functioning in the COVID-19 ICU group was shown. No other significant associations were found (all P values $> .05$) (Tab. 3).

The post hoc multivariate linear regression analyses in the COVID-19 ICU group showed significant associations between handgrip strength and ICU length of stay ($P = .035$), physical functioning and comorbidities ($P = .024$), and ICU length of stay ($P = .014$), after correction in the secondary analyses for duration of mechanical ventilation and use of ECLS/ECMO. The significant association between physical functioning and sex remained ($P = .003$), where men had a higher physical functioning score than women did. No other significant associations were found (Tab. 4).

The post hoc univariate linear regression analyses showed patients in the COVID-19 ICU group of the first wave ($n = 43$) having a significant lower score on handgrip strength ($P = .024$) compared with patients in the non-COVID-19 ICU group ($n = 89$). These differences were not found in the second wave ($n = 39$) or the third wave ($n = 8$). Furthermore, no differences were found in physical functioning and health status between patients in the COVID-19 ICU group of the first wave, the second wave, and the third wave and patients in the non-COVID-19 ICU group (Tab. 5).

Discussion

In this cross-sectional study, the first aim was to compare handgrip strength, physical functioning, and health status between the COVID-19 ICU group and the non-COVID-19 ICU group about 3 months after ICU discharge. We found no significant differences between the groups. The secondary aim was to investigate which factors were associated with the physical status outcomes in the patients in the COVID-19 ICU group. The secondary analyses showed that the number of comorbidities and sex was associated with physical functioning, where men had better physical functioning than women.

Based on the differences in patient and clinical characteristics, as well as the course during ICU stay, between the patients in the COVID-19 ICU group and the patients in the non-COVID-19 group, we expected a different physical recovery 3 months after ICU discharge. There are several possible explanations why similar handgrip strength, physical functioning, and health status were found in the COVID-19 ICU group and the non-COVID-19 ICU group. First, the demographic and clinical characteristics of both groups showed fewer differences than expected.¹⁴⁻¹⁶ Differences were confirmed only for age, duration of mechanical ventilation, and ICU length of stay. Although our post hoc analyses showed that a longer ICU length of stay was associated with a lower physical functioning after discharge in the COVID-19 ICU group, no significant difference in physical functioning was found between non-COVID-19 ICU group and the COVID-19 ICU group 3 months after ICU discharge. Perhaps these variables have less influence on physical functioning 3 months after ICU discharge than expected. Second, about 3 months after ICU discharge, the handgrip strength, physical functioning, and health status of patients in the COVID-19 ICU group and the non-COVID-19 ICU group were similar. In the non-COVID-19 ICU patient population, it is known that muscle strength and physical functioning increase mainly in the first 3 months and flatten between 3 and 6 months.^{46,47} It is possible that differences existed in the first 3 months because of an older age,^{9,10} a longer ICU length of stay,¹⁰ and a longer duration of mechanical ventilation,⁴⁸ but these possible differences were no longer present at the time of measurement. Furthermore, our data showed that physical functioning and health status of patients in the COVID-19 ICU group were lower than in healthy people approximately 3 months after ICU discharge. An earlier study showed that the physical functioning of patients in the ICU with COVID-19 remains lower compared with healthy people 6 months after ICU discharge.⁵ This may suggest that, in contrast to




Figure. Comparisons of handgrip strength, physical functioning, and health status of COVID-19 ICU, regular ICU patients and healthy people. * Steiber N. Strong or weak handgrip? Normative reference values for the German population across the life course stratified by sex, age, and body height. *PLoS One*. 2016;11.²⁰ ** Terwee CB, Roorda LD, De Vet HCW, et al. Dutch-Flemish translation of 17 item banks from the Patient-Reported Outcomes Measurement Information System (PROMIS). *Qual Life Res*. 2014;23:1733–1741.²¹ *** Hinz A, Kohlmann T, Stöbel-Richter Y, Zenger M, Brähler E. The quality of life questionnaire EQ-5D-5L: psychometric properties and normative values for the general German population. *Qual Life Res*. 2014 Mar 7;23:443–7.⁴³

the flattening in patients in the ICU without COVID-19, there is still improvement in physical functioning and health status between 3 and 6 months after ICU discharge. Moreover, the current study showed that men had a better physical functioning (on PROMIS-PF) than women. Based on the current data, there is no explanation for this.

Although not statistically significant, the difference in handgrip strength between all patients in the ICU with COVID-19 (mean = 74.2% of normative values) and patients in the ICU without COVID-19 (mean = 84.2% of normative values) is

considered clinically relevant as the MCID for HGS is smaller than the difference between these groups.⁴⁹ This indicates that muscle strength training might deserve additional attention during treatment in COVID-19 ICU patients.

Our post hoc analyses showed that patients in the COVID-19 ICU group who had an ICU admission in the first wave had a significant lower handgrip strength than patients in the non-COVID-19 ICU group. This difference disappeared in the second and third wave. In these waves, the handgrip strength of the patients with COVID-19 and the patients

Table 3. Associations Between Dependent Variables Handgrip Strength, Physical Functioning, and Health Status and Independent Variables in the COVID-19 ICU Group (n = 92)^a

Independent Variable	Handgrip Strength (n = 90) ^b			Physical Functioning ^c			Health Status ^d		
	B	95% CI	P ^e	B	95% CI	P ^e	B	95% CI	P ^e
Age	-0.41	-1.18 to 0.36	.293	0.14	-0.12 to 0.39	.30	0.04	-0.003 to 0.01	.22
Sex	3.936	-11.99 to 19.86	.615	5.779	1.54 to 10.02	.008	0.04	-0.09 to 0.17	.50
Body mass index	-0.07	-0.97 to 0.96	.989	-0.23	-0.56 to 0.11	.18	-0.007	-0.02 to 0.002	.13
Premorbid functioning ^f	-3.85	-10.83 to 3.13	.265	0.84	-1.17 to 2.85	.83	-0.03	-0.03 to 0.10	.30
Comorbidities ^g	2.31	-3.38 to 8.01	.394	-1.37	-2.64 to -0.11	.03	-0.03	-0.07 to 0.01	.20

^aICU = intensive care unit. ^bDetermined with a JAMAR handheld dynamometer and reported as percentages of reference values. ^cReported as a Patient-Reported Outcomes Measurement Information System physical function *t* value. ^dReported as a EuroQol 5 Dimensions 5 Levels index value. ^eValues in bold type were significant at *P* < .05. ^fDetermined with the Identification of Seniors at Risk–Hospitalized Patients instrument. ^gDetermined with the Charlson Comorbidity Index.

Table 4. Post Hoc Associations Between Dependent Variables Handgrip Strength, Physical Functioning, and Health Status and Independent Variables in the COVID-19 ICU Group (n = 92)^a

Independent Variable	Handgrip Strength (n = 90) ^b			Physical Functioning ^c			Health Status ^d		
	B	95% CI #	P ^e	B	95% CI	P ^e	B	95% CI	P ^e
Age	-0.36	-1.08 to 0.41	.38	-0.18	-0.79 to 0.43	.18	0.005	-0.002 to 0.01	.19
Sex	4.72	-11.17 to 20.61	.62	6.46	2.22 to 10.70	.003	-0.04	-0.09 to 0.18	.53
Body mass index	-0.18	-1.15 to 0.79	.71	-0.32	-0.65 to 0.02	.07	-0.008	-0.02 to 0.0002	.11
Premorbid functioning ^f	-3.04	-10.19 to 4.11	.39	1.18	-1.04 to 3.40	.29	0.03	-0.03 to 0.10	.31
Comorbidities ^g	2.63	-2.97 to 8.22	.33	-1.43	-2.67 to -0.20	.03	-0.02	-0.07 to 0.02	.24
ICU length of stay, d	-0.53	-1.01 to -0.04	.04	-0.19	-0.34 to -0.04	.01	-0.002	-0.007 to 0.002	.30
Duration of mechanical ventilation, d	-6.13	-24.20 to 11.94	.51	4.94	-1.57 to 11.45	.14	-0.043	-0.23 to 0.14	.65
Use of ECLS/ECMO	11.18	-27.39 to 49.75	.57	2.70	-9.92 to 15.33	.68	0.09	-0.24 to 0.42	.59

^aECLS = extracorporeal life support; ECMO = extracorporeal membrane oxygenation; ICU = intensive care unit. ^bDetermined with a JAMAR handheld dynamometer and reported as percentages of reference values. ^cReported as a Patient-Reported Outcomes Measurement Information System physical function *t* value. ^dReported as a EuroQol 5 Dimensions 5 Levels index value. ^eValues in bold type were significant at *P* < .05. ^fDetermined with the Identification of Seniors at Risk–Hospitalized Patients instrument. ^gDetermined with the Charlson Comorbidity Index.

Table 5. Comparison of Handgrip Strength, Self-Reported Physical Functioning, and Health Status Between the COVID-19 ICU Group and the Non-COVID-19 ICU Group by Wave^a

Parameter and Wave	COVID-19 ICU Group ^b		Non-COVID-19 ICU Group ^c		B	95% CI	P
	No.	Mean (SEM)	No.	Mean (SEM)			
Handgrip strength ^d							
First wave ^e	43	70.9 (3.30)	89	84.2 (4.44)	-13.37	-24.90 to -1.841	.02 ^f
Second wave ^g	39	78.7 (5.49)	89	84.2 (4.44)	-5.57	-22.55 to 11.40	.49
Third wave ^h	8	70.5 (9.39)	89	84.2 (4.44)	-13.77	-39.23 to 11.69	.28
Physical functioning ⁱ							
First wave	44	40.4 (1.46)	90	39.8 (0.99)	0.66	-2.76 to 4.09	.70
Second wave	39	37.8 (1.47)	90	39.8 (0.99)	-1.97	-5.55 to 1.62	.28
Third wave	8	36.4 (2.90)	90	39.8 (0.99)	-3.38	-10.28 to 3.53	.34
Health status ^j							
First wave	44	0.8 (0.03)	90	0.7 (0.03)	0.05	-0.04 to 0.15	.28
Second wave	39	0.7 (0.05)	90	0.7 (0.03)	-0.03	-0.14 to 0.08	.55
Third wave	8	0.7 (0.10)	90	0.7 (0.03)	-0.03	-0.30 to 0.23	.80

^aICU = intensive care unit; SEM = standard error of the mean. ^bPatients in the specific wave were included. ^cAll patients in the non-COVID-19 ICU group were included. ^dDetermined with a JAMAR handheld dynamometer and reported as percentages of reference values. ^eFirst wave: March 2020–June 2020. ^fSignificant at *P* < .05. ^gSecond wave: July 2020–January 2021. ^hThird wave: Since February 2021. ⁱReported as a Patient-Reported Outcomes Measurement Information System physical function *T* value. ^jReported as a EuroQol 5 Dimensions 5 Levels index value.

without COVID-19 was similar. This might be explained by the duration of mechanical ventilation, which was higher in the first wave compared with the second and third waves.⁵⁰ A prolonged mechanical ventilation increases the risk of muscle weakness.⁴⁸ Another explanation can be the ICU and hospital

length of stay, which decreased for patients with COVID-19 in the Netherlands since the second wave.⁵¹ In patients in the ICU without COVID-19, the association between muscle weakness and ICU length of stay was already known.⁵² Muscle weakness is also associated with hospital length of

stay.⁵³ The current post hoc analyses confirmed that handgrip strength was negatively associated with ICU stay in the patients with COVID-19.

No differences in handgrip strength, physical functioning, and health status were shown between the patients in the COVID-19 ICU group and the non-COVID-19 ICU group. In the [Figure](#), the results of handgrip strength, physical functioning, and health status of both patient groups are compared with healthy people or normative values, based on literature. Both patient groups score lower than healthy people in handgrip strength, physical functioning, and health status.^{20,21,43} Health status, measured with EQ-5D-5L, is highly correlated with performance-based outcomes, except for handgrip strength.⁵⁴ This is in line with the results on handgrip strength in our study.

Our study has several strengths. To our knowledge, it is the first published study to examine the differences in handgrip strength, physical functioning, and health status between patients with COVID-19 in the ICU and patients without COVID-19 in the ICU after ICU discharge, and all data were obtained from regular care. Furthermore, all the primary outcomes were standardized to reference values before analyses, which facilitated comparison with other studies on physical outcomes in these 2 patient populations. However, there are also some limitations. First, the sample size of this study is quite small for this kind of observational study. An increase in the number of inclusions may have increased the reliability and validity of the results; however, the minimum sample size was well reached. In the post hoc univariate linear regression analyses ([Tab. 5](#)), none of the subgroups reached the minimum group size as calculated in the power calculation. Therefore, these results should be interpreted accordingly. Second, the primary outcomes were measured only once. These outcome data collected on different time points would have increased the insight as to whether patients with COVID-19 in the ICU and patients without COVID-19 in the ICU have a different course in their recovery. Third, there might be selection bias due to multiple causes. It is possible that the patients without COVID-19 in this study differed from patients without COVID-19 before the pandemic because, due to the COVID-19 pandemic, surgeries requiring ICU admission were postponed. Furthermore, patients in the ICU were invited to receive outpatient ICU aftercare except for patients ≥ 80 years of age, with an ICU length of stay < 48 hours, a neurological disease as ICU admission diagnosis, or ICU discharge with home-based mechanical ventilation. Selection bias may have occurred because not all of the patients visited the ICU outpatient aftercare service for multiple reasons. Practice shows that some patients give no importance to visiting ICU outpatient aftercare, some patients seem to be anxious, and some patients are followed up in another hospital or in primary care. Although there are no data to objectify this, aftercare attendance is subjectively rated as good by clinicians. It is likely that nonattendance is likely to be similar between groups and, therefore, selection bias is unlikely to differ between groups. Finally, data were collected in only one hospital. UMC Utrecht is an academic hospital and treats patients with higher complexity than those treated in community hospitals. For example, at UMC Utrecht, heart and lung transplants are performed more often, which could explain the more frequent use of ECLS/ECMO in the non-COVID-19 ICU group. It is, therefore, possible that the outcomes seen in the current study may differ from those of patients who have been admitted in a community hospital.

Our study shows that both the patients in the COVID-19 ICU group and the patients in the non-COVID-19 ICU group have a lower physical functioning and health status than healthy people around 3 months after discharge. Our study confirms the importance of tailored rehabilitation after ICU discharge in the patients who had COVID-19, as previously highlighted in the literature.^{13,55,56} As with patients without COVID-19 in the ICU, a functional (re)assessment is recommended 3 months after hospital discharge, with referral to the appropriate rehabilitation or specialist services if the patient appears to be recovering at a slower rate than anticipated or if the patient has developed unanticipated physical (or non-physical) morbidity that was not previously identified. In both patient groups, it is recommended to initiate rehabilitation on the physical domain of PICS during all phases of recovery. Furthermore, it is important to give support if the patient is not recovering as quickly as they anticipated.¹²

There are also recommendations for future research. First, to provide more tailor-made care to patients who were in the ICU with COVID-19, more insight must be gained into the course of the recovery of physical status during the first period following hospital discharge. Therefore, we recommend observational studies with physical, mental, and cognitive outcomes measured on multiple time points early after ICU discharge. Second, the current study focused on differences in physical status outcomes between patients with and without COVID-19 in the ICU. It is possible that both groups differ on other outcomes that might affect recovery. Future studies should, therefore, focus on participation outcomes and modifiable factors; for example, cognitive, emotional, and psychic functioning and societal participation.

Conclusion

Based on our observational study, we conclude that, at 3 months ICU postdischarge, handgrip strength, physical functioning, and health status are comparable both for patients who were in the ICU with COVID-19 and for patients who were in the ICU without COVID-19.

Both patient groups have a lower physical status and health status than healthy people, thus requiring personalized physical rehabilitation. We recommend aftercare in the physical domain of PICS after ICU discharge in patients who were in the ICU with or without COVID-19 with an ICU length of stay > 48 hours, in primary and/or secondary care. Finally, we recommend performing a functional assessment for patients in both the COVID-19 and non-COVID-19 groups 3 months after hospital discharge.

Author Contributions

Concept/idea/research design: B. Cijs, K. Valkenet, J.M.A. Visser-Meily, M. van der Schaaf

Writing: B. Cijs, K. Valkenet, M. van der Schaaf

Data collection: J.M.A. Visser-Meily, G. Heijnen

Data analysis: B. Cijs, K. Valkenet, M. van der Schaaf

Consultation (including review of manuscript before submitting):

G. Heijnen, J.M.A. Visser-Meily

Ethics Approval

Ethical approval for the usage of daily care data was obtained from the Medical Ethical Research Committee of University Medical Center Utrecht.

Funding

There are no funders to report for this study.

Disclosures

The authors completed the ICMJE Form for Disclosure of Potential Conflicts of Interest and reported no conflicts of interest.

References

- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497–506.
- National Institute for Public Health and the Environment. *In derde golf sterke stijging ziekenhuisopnames jongere leeftijdsgroepen*. Accessed May 31, 2023. <https://www.rivm.nl/nieuws/in-derde-golf-sterke-stijging-ziekenhuisopnames-jongere-leeftijdsgroepen>.
- Abate SM, Ali SA, Mantfardo B, Basu B. Rate of intensive care unit admission and outcomes among patients with coronavirus: a systematic review and meta-analysis. *PLoS One*. 2020;15:1–19. <https://doi.org/10.1371/journal.pone.0235653>.
- Moonen HPFX, Strookappe B, Zanten ARH. Physical recovery of COVID-19 pneumosepsis intensive care survivors compared with non-COVID pneumosepsis intensive care survivors during post-intensive care hospitalization: the RECOVID retrospective cohort study. *J Parenter Enter Nutr*. 2022;46:798–804.
- Neville TH, Hays RD, Tseng CH, et al. Survival after severe COVID-19: long-term outcomes of patients admitted to an intensive care unit. *J Intensive Care Med*. 2022;1019:1–10. <https://doi.org/http://journals.sagepub.com/doi/10.1177/08850666221092687>, 1028.
- Thiolliere F, Falandry C, Allaouchiche B, et al. Intensive care-related loss of quality of life and autonomy at 6 months post-discharge: does COVID-19 really make things worse? *Crit Care*. 2022;26:94. <https://doi.org/10.1186/s13054-022-03958-6>.
- Needham DM, Davidson J, Cohen H, et al. Improving long-term outcomes after discharge from intensive care unit: report from a stakeholders' conference. *Crit Care Med*. 2012;40:502–509. <https://doi.org/10.1097/CCM.0b013e318232da75>.
- Ohtake PJ, Lee AC, Scott JC, et al. Physical impairments associated with post-intensive care syndrome: systematic review based on the world health organization's international classification of functioning, disability and health framework. *Phys Ther*. 2018;98:631–645.
- Hodgson CL, Denehy L. Measuring physical function after ICU: one step at a time. *Intensive Care Med*. 2017;43:1901–1903.
- Ámundadóttir ÓR, Jónasdóttir RJ, Sigvaldason K, et al. Predictive variables for poor long-term physical recovery after intensive care unit stay: an exploratory study. *Acta Anaesthesiol Scand*. 2020;64:1477–1490.
- Connolly B. Describing and measuring recovery and rehabilitation after critical illness. *Curr Opin Crit Care*. 2015;21:445–452.
- National Institute for Health and Care Excellence LU. *NICE: Rehabilitation After Critical Illness*. 2009. Accessed October 20, 2020. <http://www.nice.org.uk/guidance/cg83>. Manchester, UK: National Institute for Health and Care Excellence (NICE).
- Connolly B, Salisbury L, O'Neill B, et al. Exercise rehabilitation following intensive care unit discharge for recovery from critical illness: executive summary of a Cochrane collaboration systematic review. *J Cachexia Sarcopenia Muscle*. 2016;7:520–526.
- Phua J, Weng L, Ling L, et al. Intensive care management of coronavirus disease 2019 (COVID-19): challenges and recommendations. *Lancet Respir Med*. 2020;8:506–517.
- Brinkman S, De Jonge E, Abu-Hanna A, et al. Mortality after hospital discharge in ICU patients. *Crit Care Med*. 2013;41:1229–1236.
- Grasselli G, Zangrillo A, Zanella A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region, Italy. *JAMA*. 2020;323:1574–1581.
- Stuurgroep Onderzoek Corona. *Advies Basisset Klinimetric Nazorgpoli*. Accessed October 20, 2020. https://revalidatiegeneeskunde.nl/sites/default/files/attachments/Beleid/COVID-19/advies_basisset_klinimetric_ic_nazorg_poli_-_30_april_2020.pdf.
- Fess EE. A method for checking Jamar dynamometer calibration. *J Hand Ther*. 1987;1:28–32.
- Tipping CJ, Young PJ, Romero L, Saxena MK, Dulhunty J, Hodgson CL. A systematic review of measurements of physical function in critically ill adults. *Crit Care Resusc*. 2012;14:302–311.
- Steiber N. Strong or weak handgrip? Normative reference values for the German population across the life course stratified by sex, age, and body height. *PLoS One*. 2016;11:e0163917. <https://doi.org/10.1371/journal.pone.0163917>.
- Terwee CB, Roorda LD, de HCW, et al. Dutch-Flemish translation of 17 item banks from the patient-reported outcomes measurement information system (PROMIS). *Qual Life Res*. 2014;23:1733–1741.
- Crins MHP, van der Wees PJ, Klausch T, van Dulmen SA, Roorda LD, Terwee CB. Psychometric properties of the PROMIS physical function item bank in patients receiving physical therapy. *PLoS One*. 2018;13:e0192187. <https://doi.org/10.1371/journal.pone.0192187>.
- Schalet BD, Hays RD, Jensen SE, Beaumont JL, Fries JF, Cella D. Validity of PROMIS physical function measured in diverse clinical samples. *J Clin Epidemiol*. 2016;73:112–118.
- EuroQol. EuroQol—a new facility for the measurement of health-related quality of life. *Health Policy*. 1990;3:199–208.
- Janssen MF, Pickard AS, Golicki D, et al. Measurement properties of the EQ-5D-5L compared to the EQ-5D-3L across eight patient groups: a multi-country study. *Qual Life Res*. 2013;22:1717–1727.
- Nolan CM, Longworth L, Lord J, et al. The EQ-5D-5L health status questionnaire in COPD: validity, responsiveness and minimum important difference. *Thorax*. 2016;71:493–500.
- Scalone L, Ciampichini R, Fagioli S, et al. Comparing the performance of the standard EQ-5D 3L with the new version EQ-5D 5L in patients with chronic hepatic diseases. *Qual Life Res*. 2013;22:1707–1716.
- Golicki D, Niewada M, Buczek J, et al. Validity of EQ-5D-5L in stroke. *Qual Life Res*. 2015;24:845–850.
- Hernandez G, Garin O, Dima AL, et al. EuroQol (EQ-5D-5L) validity in assessing the quality of life in adults with asthma: cross-sectional study. *J Med Internet Res*. 2019;21:1–12.
- Milton A, Schandl A, Soliman I, et al. ICU discharge screening for prediction of new-onset physical disability—a multinational cohort study. *Acta Anaesthesiol Scand*. 2020;64:789–797.
- Földi M, Farkas N, Kiss S, et al. Obesity is a risk factor for developing critical condition in COVID-19 patients: a systematic review and meta-analysis. *Obes Rev*. 2020;21:e13095. <https://doi.org/10.1111/obr.13095>.
- Garrow JS, Webster J. Quetelet's index (W/H²) as a measure of fatness. *Int J Obes*. 1985;9:147–153.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40:373–383.
- Hoogerduijn JG, Grobbee DE, Schuurmans MJ. Prevention of functional decline in older hospitalized patients: nurses should play a key role in safe and adequate care. *Int J Nurs Pract*. 2014;20:106–113.
- Li TS, Gomersall CD, Joynt GM, Chan DPS, Leung P, Hui DSC. Long-term outcome of acute respiratory distress syndrome caused by severe acute respiratory syndrome (SARS): an observational study. *Crit Care Resusc*. 2006;8:302–308.
- De Groot V, Beckerman H, Lankhorst GJ, Bouter LM. How to measure comorbidity: a critical review of available methods. *J Clin Epidemiol*. 2003;56:221–229.

37. Umegaki T, Ikai H, Imanaka Y. The impact of acute organ dysfunction on patients' mortality with severe sepsis. *J Anaesthesiol Clin Pharmacol*. 2011;27:180–184.
38. Póvoa P, Almeida E, Moreira P, et al. C-reactive protein as an indicator of sepsis. *Intensive Care Med*. 1998;24:1052–1056.
39. Nakamura M, Oda S, Sadahiro T, et al. Correlation between high blood IL-6 level, hyperglycemia, and glucose control in septic patients. *Crit Care*. 2012;16:R58. <https://doi.org/10.1186/cc11301>.
40. *SPSS Statistics 27*. Armonk, NY: IBM; 2020.
41. Morris SB. Estimating effect sizes from pretest-posttest-control group designs. *Organ Res Methods*. 2008;11:364–386.
42. Sterne JAC, White IR, Carlin JB, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ*. 2009;338:b2393. <https://doi.org/10.1136/bmj.b2393>.
43. Hinz A, Kohlmann T, Stöbel-Richter Y, Zenger M, Brähler E. The quality of life questionnaire EQ-5D-5L: psychometric properties and normative values for the general German population. *Qual Life Res*. 2014;23:443–447.
44. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. *Statistical Power Analysis for the Behavioral Sciences*. New York, NY: Taylor and Francis; 2013: 400.
45. Austin PC, Steyerberg EW. The number of subjects per variable required in linear regression analyses. *J Clin Epidemiol*. 2015;68: 627–636.
46. Eggmann S, Luder G, Verra ML, Irincheeva I, Bastiaenen CHG, Jakob SM. Functional ability and quality of life in critical illness survivors with intensive care unit acquired weakness: a secondary analysis of a randomised controlled trial. *PLoS One*. 2020;15:e0229725. <https://doi.org/10.1371/journal.pone.0229725>.
47. Herridge MS, Chu LM, Matte A, et al. The RECOVER program: disability risk groups and 1-year outcome after 7 or more days of mechanical ventilation. *Am J Respir Crit Care Med*. 2016;194: 831–844.
48. Fan E, Cheek F, Chlan L, et al. An official American thoracic society clinical practice guideline: the diagnosis of intensive care unit-acquired weakness in adults. *Am J Respir Crit Care Med*. 2014;190:1437–1446.
49. Bohannon RW. Minimal clinically important difference for grip strength: a systematic review. *J Phys Ther Sci*. 2019;31: 75–78.
50. Karagiannidis C, Hentschker C, Westhoff M, et al. Changes in utilization and outcomes of mechanical ventilation of COVID-19 during the course of the pandemic in Germany in 2020: an observational study of 7,490 patients. *medRxiv*. 2021;17: 1–12.
51. National Institute for Public Health and the Environment. *Het nieuwe coronavirus in Nederland. Wat is het verschil tussen de eerste golf en de tweede golf*. 2020: 2.
52. Hermans G, van den Berghe G. Clinical review: intensive care unit acquired weakness. *Crit Care*. 2015;19:274–283.
53. Vanhorebeek I, Latronico N, Van den Berghe G. ICU-acquired weakness. *Intensive Care Med*. 2020;46:637–653.
54. Paton M, Lane R, Paul E, Linke N, Shehabi Y, Hodgson CL. Correlation of patient-reported outcome measures to performance-based function in critical care survivors: PREDICTABLE. *Austral Crit Care*. 2022. <https://doi.org/10.1016/j.aucc.2022.05.006>.
55. Major ME, Kwakman R, Kho ME, et al. Surviving critical illness: what is next? An expert consensus statement on physical rehabilitation after hospital discharge. *Crit Care*. 2016;20:354. <https://doi.org/10.1186/s13054-016-1508-x>.
56. Major ME, Dettling-Ihnenfeldt D, Ramaekers SPJ, Engelbert RHH, van der Schaaf M. Feasibility of a home-based interdisciplinary rehabilitation program for patients with post-intensive care syndrome: the REACH study. *Crit Care*. 2021;25:1–15. <https://doi.org/10.1186/s13054-021-03709-z>.