

ORIGINAL RESEARCH

Prevalence and Determinants of Pain in Spinal Cord Injury During Initial Inpatient Rehabilitation: Data From the Dutch Spinal Cord Injury Database



Tim C. Crul, MD,^{a,b} Marcel W.M. Post, PhD,^{a,b,c} Johanna M.A. Visser-Meily, MD, PhD,^{a,b,d} Janneke M. Stolwijk-Swüste, MD, PhD^{a,b,e}

From the ^aCentre of Excellence for Rehabilitation Medicine, UMC Utrecht Brain Centre, University Medical Centre Utrecht, Utrecht; ^bDe Hoogstraat Rehabilitation, Utrecht; ^cUniversity of Groningen, University Medical Centre Groningen, Centre for Rehabilitation, Groningen; ^dDepartment of Rehabilitation, Physical Therapy Science, and Sports, UMC Utrecht Brain Center, University Medical Center Utrecht, Utrecht; and ^eDepartment of Spinal Cord Injury and Orthopedics, De Hoogstraat Rehabilitation, Utrecht, the Netherlands.

Abstract

Objective: To describe the prevalence and characteristics of spinal cord injury (SCI)-related pain during initial inpatient rehabilitation and to investigate relationships with demographic and lesion characteristics.

Design: Cohort during inpatient rehabilitation.

Setting: Eight specialized SCI rehabilitation centers in the Netherlands.

Participants: Patients with newly acquired SCI admitted for inpatient rehabilitation between November 2013 and August 2019 (N=1432).

Interventions: Not applicable.

Main Outcome Measures: Presence of pain at admission and discharge. Logistic regression analyses were used to study the prevalence of pain related to sex, age, etiology, completeness, and level of injury.

Results: Data from 1432 patients were available. Of these patients 64.6% were male, mean age was 56.8 years, 59.9% had a nontraumatic SCI, 63.9% were classified as American Spinal Cord Injury Association Impairment Scale (AIS) D and 56.5% had paraplegia. Prevalence of pain was 61.2% at admission (40.6% nociceptive pain [NocP], 30.2% neuropathic pain [NeuP], 5.4% other pain) and 51.5% at discharge (26.0% NocP, 31.4% NeuP, 5.7% other pain). Having NocP at admission was associated with traumatic SCI. AIS B had a lower risk of NocP than AIS D at admission. Having NocP at discharge was associated with female sex and traumatic SCI. AIS C had a lower risk of NocP at discharge than AIS D. Having NeuP at admission was associated with female sex. Having NeuP at discharge was associated with female sex, age younger than 65 years vs age older than 75 years and tetraplegia.

Conclusions: SCI-related pain is highly prevalent during inpatient rehabilitation. Prevalence of NocP decreased during inpatient rehabilitation, and prevalence of NeuP stayed the same. Different patient and lesion characteristics were related to the presence of SCI-related pain. Healthcare professionals should be aware of these differences in screening patients on presence and development of pain during inpatient rehabilitation.

Archives of Physical Medicine and Rehabilitation 2023;104:74–82

© 2022 by the American Congress of Rehabilitation Medicine. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

The development of the Dutch Spinal Cord Injury Database (DSCID) was supported by Rehabilitation Netherlands (Revalidatie Nederland), Innovation in Rehabilitation program, grant IPR2011-12. All spinal cord injury rehabilitation centers in the Netherlands contributed to the DSCID.

Disclosures: none

Spinal cord injury (SCI) results in varying degrees of function loss distal to the level of the injury.¹ Besides motor and sensory function loss, SCI is often accompanied by secondary conditions that affect quality of life.^{2,3} Pain is one of the secondary conditions that is frequently reported by patients with SCI. The prevalence of SCI-related pain is estimated at 61%, with a large amount of

0003-9993/\$36 - see front matter © 2022 by the American Congress of Rehabilitation Medicine. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

<https://doi.org/10.1016/j.apmr.2022.07.005>

heterogeneity in studies. Pain is one of the most challenging symptoms to manage.⁴ The prognosis of SCI-related pain is poor, with many patients reporting the pain to remain the same or even worsen over time.⁵ In addition, SCI-related pain is known to have a severe negative effect on quality of life.^{2,6}

SCI-related pain is classified by the International Association of the Study of Pain into 2 main types: nociceptive pain (NocP) and neuropathic pain (NeuP).⁵ Nociceptive pain is caused by tissue damage^{2,6} and is divided in 2 main categories: musculoskeletal pain and visceral pain.⁷ Examples of musculoskeletal pain include pain resulting from joint arthritis, spinal fractures, muscle injury, rotator cuff tendinopathy, and muscle spasms.⁷ Examples of causes of visceral pain are constipation, urinary tract infection, ureteral calculus, and fecal impaction.⁷ Neuropathic pain is caused by damage or dysfunction of the nervous system.^{8,9} In SCI, NeuP can be located above, at, or below injury level.² Pain above injury level is often not because of the SCI itself but caused by concomitant compression neuropathy or by regional pain syndromes.² Pain at injury level can be caused by damage or disease of the nerve root or spinal cord.^{2,7} Pain below injury level is caused directly by injury of the spinal cord.^{2,7}

The treatment of SCI-related pain remains a challenge.¹⁰⁻¹² Pharmacologic treatment strategy for NocP is based on the stepped care of paracetamol, nonsteroidal anti-inflammatory drugs, and opioids in accordance with the World Health Organization Cancer Pain Relief.^{13,14} For NeuP, pregabalin and gabapentin are the most evidence based treatment options and have shown a significant reduction in pain intensity in multiple studies but with limited evidence on long-term effects.^{11,12,15} Besides pharmacologic interventions, frequently used nonpharmacologic treatments, such as acupuncture, exercise programs, cranial electrotherapy stimulation, and psychological therapies such as cognitive behavioral therapy are used.^{15,16} Little evidence is found on the effectiveness of these treatments on pain relief.^{15,16}

It is important to gain a better understanding of SCI-related pain during inpatient rehabilitation to improve future treatment. Presence of SCI-related pain tends to change over time; therefore, studies on pain in community-dwelling patients with SCI might not relate to pain at inpatient rehabilitation.³ Large studies on the presence of pain during inpatient rehabilitation are scarce and do not report the type of pain¹⁷ or do not relate the patient and lesion characteristics to the type of pain.¹⁸ Finding patient and lesion characteristics associated with SCI-related pain and type of pain can identify which patients are at risk for developing pain and facilitate patient screening during inpatient rehabilitation. Addressing pain at an early stage could potentially benefit treatment in the long-term. The objective of this study is to describe the prevalence of SCI-related pain during inpatient rehabilitation, describe the characteristics of such pain, and investigate possible associations to different patient and lesion characteristics.

List of abbreviations:

AIS	American Spinal Cord Injury Association Impairment Scale
DSCID	Dutch Spinal Cord Injury Database
NeuP	neuropathic pain
NocP	nociceptive pain
NRS	numeric rating scale
SCI	spinal cord injury

Methods

Participants

This study uses data from the Dutch Spinal Cord Injury Database (DSCID).¹⁹ The DSCID is a joint effort of the 8 SCI specialized Dutch rehabilitation centers. This study used data of the DSCID collected between November 2013 and August 2019. Patients were included in the current study if data on presence of pain (yes or no) were available at both admission and discharge, patients did not object to the sharing of deidentified data, and time between onset of SCI and admission to the rehabilitation center was within 9 months (to exclude readmissions).

Procedure

This study was conducted according the principles of the Declaration of Helsinki. Informed consent varies between centers. Some centers ask for written informed consent in advance, whereas other centers inform patients with oral and written information about the use of anonymous data for research purposes. Patients can decline the use of these data, including the data of the DSCID. Clinical data of patients with SCI were collected at admission and discharge. A designated member of the clinical staff is responsible for data collection at each center. Information is either retrieved from the electronic medical record retrospectively or added prospectively at admission of the patient to the rehabilitation center and entered into the DSCID anonymously. The notation in the electronic medical record or the admission to the rehabilitation center is always supervised by a SCI rehabilitation physician. In this way, entering NocP or NeuP is similar in all rehabilitation centers.

Instruments

The DSCID contains data on age, sex, etiology of SCI, various secondary conditions of SCI, functional status, and quality of life. American Spinal Cord Injury Association Impairment Scale (AIS) classification and neurologic level of injury were assessed according to the International Standards for Neurological Classification of Spinal Cord Injury, revised version of 2011.²⁰ Data on pain were collected according to an amendment to the Spinal Cord Injury Pain Basic Data Set, which was a shorter version.²¹ Information on the presence of pain; the type of pain divided into NocP (which were subdivided into musculoskeletal and visceral pain), NeuP (which was divided into above level, at level and below level of injury, where at level pain extends to 3 levels below the level of injury), and other pain; intensity of pain on the 11-point numeric rating scale (NRS); treatment of pain; and type of treatment (oral medication, physiotherapy, psychology, other) was available from the DSCID.

Analysis

The statistical analyses were performed using SPSS Version 26.^a Frequencies were reported on the sociodemographic characteristics of the included patients and the characteristics of pain. Independent samples *t* tests were used to compare means for continuous variables, and chi-square tests were used to compare categorical variables on baseline characteristics of patients included and excluded from the DSCID. An independent samples

t test was used to compare mean time since onset between patients with pain and without pain. Logistic regression analyses were performed to examine whether a prediction on the prevalence of pain could be made using the variables age, sex, etiology, AIS classification, and level of injury for which unadjusted and adjusted odds ratios were calculated. Little's Missing Completely at Random tests were used to evaluate whether missing data were distributed randomly.

Results

From a total of 2700 patients registered in the DSCID, 1432 were included in the analysis. A flow diagram of inclusion of patients is shown in [fig 1](#). Patient characteristics at admission of the included and excluded patients are shown in [table 1](#). The included group contained less patients with AIS A and more patients with AIS D than the excluded group. Exclusion of patients readmitted to the rehabilitation center (time since onset >9 months) resulted in a significant difference in time since onset between included and excluded patients. Among the included patients, most common causes of traumatic SCI were falls (54.5%), traffic collisions (22.3%), and sports-related accidents (14.3%). Most common causes of nontraumatic SCI were vascular (22.4%), benign or malignant tumor growth (21.8%), degeneration (20.8%), and inflammation (15.9%). Most patients had an incomplete SCI in which AIS D occurred most frequently.

Prevalence of pain is reported in [table 2](#). Pain was reported in 61.1% at admission and in 51.5% at discharge. NocP was reported in 40.4% of the patients at admission and in 26.1% at discharge. NeuP was reported in 30.5% of the patients at admission and in 31.7% at discharge. Change in pain is reported in [table 3](#). Characteristics of NocP and NeuP are shown in [tables 4 and 5](#), respectively. Above-level NeuP was described as other NeuP. This

occurred in 3 patients, of which 2 had a combination with at-level and below-level NeuP.

Time since onset of SCI was not significantly different in the group with pain and without pain in the total sample (mean, 38.1 days and 38.2 days; $P=.96$ at admission and mean, 130 and 126 days; $P=.39$ at discharge, respectively) nor in the subgroups with nociceptive pain (mean, 44.8 and 47.6 days; $P=.47$ at admission and mean, 126 and 129 days; $P=.56$ at discharge, respectively) and neuropathic pain (mean, 38.7 and 37.3 days; $P=.63$ at admission and mean, 132 and 136 days; $P=.25$ at discharge, respectively).

Patient characteristics on pain

The odds ratios of NocP and NeuP in relationship to patient characteristics is shown in [tables 6 and 7](#), respectively. At admission, patients with AIS B had a lower risk of having NocP than those with AIS D. Patients with traumatic SCI had a higher risk of having NocP. At discharge female patients and patients with traumatic SCI had a higher risk of having NocP, and patients with AIS C had a lower risk of having NocP than those with AIS D. Sex, age, etiology, and level of injury showed significant relations to the prevalence of NeuP. At admission, female patients had a higher risk of having NeuP. At discharge female patients, patients younger than 65 years compared with those older than 75 years, and patients with tetraplegia had a higher risk of having NeuP.

Missing data

Little's Missing Completely at Random tests showed that missing data were distributed randomly and did not occur more frequently in specific subgroups.

Discussion

In our population of 1432 patients with newly acquired SCI, pain is highly prevalent. The prevalence of NocP decreased during inpatient rehabilitation, and prevalence of NeuP stayed the same. Female patients and patients having traumatic SCI had a higher risk of having NocP. Female patients, younger patients, and patients with tetraplegia had a higher risk of having NeuP.

The prevalence of pain reported in this study is lower than reported in the current literature. One study performed in Switzerland on pain during inpatient rehabilitation reported a pain prevalence of 87% at admission and 74% at discharge.¹⁷ This study did not make a distinction between different types of pain. Another study performed in Italy on pain at inpatient rehabilitation showed a prevalence of 72% during inpatient stay, with a prevalence of NocP and NeuP of 52% and 48%, respectively.¹⁸ Most studies report the presence of pain in community-dwelling patients with SCI, which is different from pain prevalence at inpatient rehabilitation. In a recent meta-analysis, with large variation in time since onset, the pooled prevalence of musculoskeletal pain was 56%, and the pooled prevalence of visceral pain was 20%.²² The differences in prevalence between the studies can also be explained by the different patient characteristics of the studies. Musculoskeletal pain could develop in long-term wheelchair users (range, 5-56 years) because of overuse of the upper extremity, spasticity, and muscle contractures.²³ During inpatient rehabilitation there seems to be a decrease in NocP. [Table 3](#) shows that 296 patients ceased to have NocP during inpatient rehabilitation, while 104 developed




Fig 1 Flow diagram.

Table 1 Patient characteristics at admission

Characteristic	Included Participants=1432	Excluded Participants=1268	P Value
Sex, n (%)			.474
Female	507 (35.4)	465 (36.7)	
Male	925 (64.6)	801 (63.3)	
Missing	0	2	
Age (y)			.334
Mean \pm SD	56.8 \pm 16.5	57.3 \pm 16.8	
Median (IQR)	60 (22)	61 (23)	
Missing	10	6	
Time since onset at admission (d)			<.001*
Mean \pm SD	38.2 \pm 45.9	56.4 \pm 23.01	
Median (IQR)	22 (26)	30 (56)	
Missing	0	340	
Length of stay (d)			.595
Mean \pm SD	92.4 \pm 78.7	90.38 \pm 65	
Median (IQR)	70 (84)	72 (74.5)	
Missing	43	547	
AIS classification, n (%)			<.001*
AIS A	159 (12.3)	160 (18.4)	
AIS B	125 (9.7)	92 (10.6)	
AIS C	183 (14.2)	127 (14.6)	
AIS D	825 (63.9)	492 (56.5)	
Missing	140	397	
Etiology of SCI, n (%)			.060
Traumatic	566 (40.1)	417 (36.5)	
Nontraumatic	846 (59.9)	727 (63.5)	
Missing	20	124	
Tetraplegia/paraplegia			.070
Tetraplegia	557 (43.5)	412 (47.5)	
Paraplegia	722 (56.5)	455 (52.5)	
Missing	153	401	

Abbreviations: IQR, interquartile range.

* Significant difference between groups.

NocP. This could be explained by pain resulting from the trauma or surgery disappearing over time. It is also possible that intensive physical treatment from a physical therapist during inpatient rehabilitation has a positive effect on managing NocP symptoms as

has been described for SCI-related shoulder pain in those using wheelchairs for over 1 year.²⁴ Increased use of physical therapy as a treatment for NocP during inpatient rehabilitation is also described in [table 4](#), where 28.2% of patients with NocP use

Table 2 Prevalence of pain at admission and discharge

Total Participants=1432 Pain, n (%)	Admission	Discharge
Total pain		
Yes	877 (61.2)	737 (51.5)
No	555 (38.8)	695 (48.5)
Nociceptive pain		
Yes	556 (40.6)	352 (26.0)
No	814 (59.4)	1002 (74.0)
Missing	62	78
Neuropathic pain		
Yes	411 (30.2)	436 (31.4)
No	951 (69.8)	952 (68.6)
Missing	70	44
Other pain		
Yes	74 (5.4)	78 (5.7)
No	1296 (94.6)	1293 (94.3)
Missing	62	61

Table 3 Change in pain during inpatient rehabilitation

Pain, n (%)	At Discharge	
	Yes, n (%)	No, n (%)
Total pain at admission		
Yes	580 (66.1)	297 (33.9)
No	157 (28.3)	398 (71.7)
Nociceptive pain at admission		
Yes	230 (43.7)	296 (56.3)
No	104 (13.4)	673 (86.6)
Missing*	129	
Neuropathic pain at admission		
Yes	240 (59.4)	164 (40.6)
No	176 (19.0)	748 (81.0)
Missing*	104	
Other pain at admission		
Yes	9 (12.9)	61 (87.1)
No	66 (5.3)	1180 (94.7)
Missing*	106	

* Missing data on type of pain at admission and/or discharge.

Table 4 Characteristics of nociceptive pain

Pain	Patients With Nociceptive Pain at Admission (n=556)	Patients With Nociceptive Pain at Discharge (n=352)
Pain intensity		
NRS, mean \pm SD (min-max)	4.37 \pm 2.15 (0-10)	4.19 \pm 2.12 (0-10)
NRS, median (IQR)	4 (3)	4 (3)
Missing	244	102
Type of pain, n (%)		
Musculoskeletal pain	448 (90.7)	250 (92.3)
Visceral pain	26 (5.3)	12 (4.4)
Other	20 (4.0)	9 (3.3)
Missing	62	81
Treatment, n (%)		
Yes	486 (93.6)	291 (85.6)
No	33 (6.4)	49 (14.4)
Missing	37	12
Type of treatment, n (%)*		
Oral medication	475 (91.5)	270 (79.4)
Physiotherapy	77 (14.8)	96 (28.2)
Psychotherapy	1 (0.2)	4 (1.2)
Other	13 (2.5)	16 (4.7)

Abbreviations: IQR, interquartile range.

* % of included patients with nociceptive pain and with data on treatment. Multiple treatments or no treatment can be used in 1 patient. Therefore, total percentages can exceed or be lower than 100%.

Table 5 Characteristics of neuropathic pain

Characteristic	Patients With Neuropathic Pain at Admission (n=411)	Patients With Neuropathic Pain at Discharge (n=436)
Pain intensity		
NRS, mean \pm SD (min-max)	4.69 \pm 2.15 (0-10)	4.38 \pm 2.29 (0-10)
NRS, median (IQR)	5 (4)	4 (3)
Missing	156	84
Level of pain, n (%)		
At level	82 (22.3)	85 (21.9)
Below level	214 (58.2)	223 (57.3)
At and below level	65 (17.7)	70 (18.0)
Other	7 (1.9)	11 (2.8)
Missing	43	47
Treatment, n (%)		
Yes	291 (72.6)	352 (82.7)
No	110 (27.4)	74 (17.3)
Missing	10	8
Type of treatment, n (%)*		
Oral medication	287 (71.6)	346 (80.8)
Physiotherapy	17 (4.2)	41 (9.6)
Psychotherapy	1 (0.2)	5 (1.2)
Other	5 (1.2)	14 (3.3)

* % of included patients with nociceptive pain and with data on treatment. Multiple treatments or no treatment can be used in 1 patient. Therefore total percentages can exceed or be lower than 100%.

physical therapy as a treatment compared with 14.8% at admission. Inactivity during hospitalization could also explain the higher pain levels of NocP at admission and decrease at discharge because training with a physical therapist increases during inpatient rehabilitation. The fact that inpatients start using wheelchairs, however, could explain the portion of patients that develop NocP during inpatient rehabilitation because rotator cuff pathology occurs more frequently in manual wheelchair users with SCI.²⁵

The prevalence of NeuP reported in this study is lower than other studies report. A wide range of percentages on the prevalence of NeuP exist in current literature, varying between 32% and 92%.²⁶⁻³³ Meta-analyses published in 2017 and 2021 estimated the pooled prevalence of NeuP to be 53% and 58%, respectively.^{8,22} This study collected information during inpatient rehabilitation and therefore did not examine the prevalence of NeuP over a longer period of time. As shown in table 3, the number of patients that develop NeuP during inpatient rehabilitation is similar to the number of patients that cease to have NeuP (176 vs 164, respectively). NeuP mostly develops in a few years' time, and prevalence increases after 1 year.^{8,26,28} Most patients reported NeuP to be present below level of lesion. Other studies also report NeuP to be mostly present below neurologic level of injury.^{28,30-32,34,35} In the meta-analysis of Burke et al, pooled prevalence of at-level NeuP in patients with SCI was 19%, whereas pooled prevalence of below-level NeuP was 27%.⁸ In this study, the prevalence of NeuP can only be generalized to patients with SCI during inpatient rehabilitation and not the entire population with SCI. Long-term follow-up at inpatient rehabilitation and after discharge would be useful to study the change in pain over time.

Demographic characteristics in this study were comparable with other studies on SCI. Mean age at admission was 57 years. The mean age in other studies on SCI range from 40-62 years.^{3,36-42} The majority of patients (64.6%) in this study are male, which is similar to other studies.^{4,37,43} Etiology of SCI is comparable with many other studies that include both traumatic and nontraumatic SCI.^{37,42,43} A minority of the patients in this study had a complete SCI (AIS A). This low number of complete SCI is in accordance to other SCI studies.^{3,36,37} There are differences on demographics in different parts of the world, for example, traumatic SCI is more prevalent in Northern America than Western Europe.⁴⁴

The mean pain intensity of NocP and NeuP reported in this study is comparable with other studies reporting pain during inpatient rehabilitation.^{17,45} It is lower than in other studies reporting on pain in the 5-8 years post injury, where pain ranged from 5.7-5.8 on the NRS in musculoskeletal pain, 6.6 in visceral pain, and 5.7-6.4 in NeuP.^{3,35} This difference could be explained by pain worsening over time, as described before.⁵ It is likely that assessment in other studies focus solely on pain, which could have led to a higher score on the NRS because of selection bias. Patients included in the current study were assessed on multiple questions regarding their SCI. This could have led to the large amount of missing data on NRS because the priority might not have been on pain during their inpatient rehabilitation.

A meta-analysis performed on determinants of pain in SCI did not find a relationship between pain and patient or lesion characteristics.⁴ In this study women and patients having a traumatic SCI had a higher risk of having NocP, and female patients, patients younger than 65 years, and patients with tetraplegia had a higher risk of having NeuP. Other studies also found sex influenced SCI-related pain, where pain is related to female sex.^{26,46,47} This could also be explained by the different presentation of pain in sex, unrelated to SCI.⁴⁸ In accordance with results on this study, Siddall et

Table 6 Prevalence of nociceptive pain in relationship to patient and lesion characteristics

Variable	Admission			Discharge		
	Patients With Pain (n=556), n (%) [*]	Patients With Pain vs Without Pain		Patients With Pain (n=352), n (%) [*]	Patients With Pain vs Without Pain	
		Unadjusted OR (95% CI)	Adjusted OR [†] (95% CI)		Unadjusted OR (95% CI)	Adjusted OR [†] (95% CI)
Sex						
Female	199 (41.3)	1.05 (0.84-1.31)	1.10 (0.86-1.41)	138 (29.2)	1.28 (1.00-1.65)	1.37 (1.04-1.82) [‡]
Male	357 (40.2)	1.00	1.00	214 (24.3)	1.00	1.00
Missing	0			0		
Age (y)						
≤44	135 (44.7)	1.17 (0.80-1.71)	1.04 (0.68-1.59)	78 (25.7)	1.07 (0.69-1.66)	1.05 (0.64-1.73)
45-54	99 (44.4)	1.16 (0.78-1.73)	1.07 (0.69-1.66)	60 (27.0)	1.15 (0.72-1.82)	1.05 (0.63-1.76)
55-64	106 (35.2)	0.79 (0.54-1.16)	0.78 (0.51-1.18)	81 (27.6)	1.18 (0.76-1.82)	1.20 (0.74-1.96)
65-74	142 (39.4)	0.95 (0.65-1.37)	0.96 (0.65-1.44)	89 (24.9)	1.03 (0.67-1.58)	1.05 (0.66-1.68)
≥75	71 (40.8)	1.00	1.00	41 (24.4)	1.00	1.00
Missing	3			3		
AIS classification at admission						
AIS A	74 (47.7)	1.32 (0.94-1.87)	1.02 (0.70-1.50)	45 (29.2)	1.19 (0.81-1.74)	1.01 (0.65-1.58)
AIS B	40 (33.3)	0.72 (0.48-1.08)	0.64 (0.42-0.98) [‡]	23 (19.3)	0.69 (0.43-1.12)	0.69 (0.42-1.14)
AIS C	65 (36.9)	0.85 (0.60-1.19)	0.71 (0.50-1.01)	34 (20.2)	0.73 (0.49-1.10)	0.60 (0.38-0.93) [‡]
AIS D	323 (40.9)	1.00	1.00	202 (25.8)	1.00	1.00
Missing	54			48		
Etiology of the SCI						
Traumatic	262 (48.2)	1.68 (1.35-2.10) [‡]	1.98 (1.50-2.62) [‡]	157 (29.5)	1.35 (1.05-1.71) [‡]	1.53 (1.12-2.09) [‡]
Nontraumatic	287 (35.6)	1.00	1.00	191 (23.8)	1.00	1.00
Missing	7			4		
Level of injury						
Tetraplegia	220 (40.7)	1.00 (0.80-1.26)	0.83 (0.64-1.07)	129 (24.6)	0.97 (0.75-1.26)	0.95 (0.68-1.24)
Paraplegia	279 (40.7)	1.00	1.00	173 (25.1)	1.00	1.00
Missing	57			50		

Abbreviations: CI, confidence interval; OR, odds ratio.
^{*} Percentage of pain within subgroup.
[†] Adjusted for sex, age, time since onset, AIS classification, etiology of SCI, and level of injury.
[‡] Statistically significant (CI≠1.00).

al described patients with tetraplegia to have a positive correlation to having NeuP.³ A study in the Swiss population found that chronic pain, with no distinction between types of pain, was more prevalent in the oldest age group (61 years or older) than in the youngest age group (16-30 years).⁴⁶ Another study at inpatient rehabilitation also found that having pain was related to older age.¹⁸ These studies did not differentiate between types of pain. The relationship between age and specifically having NeuP has not been described in previous research.

This study found a lower risk for patients with AIS B and AIS C. There was no relationship between the presence of NeuP and injury severity. Until now, only few studies have investigated the influence of injury severity on pain, mostly on the presence of NeuP. Different studies show conflicting results on the relationship between injury severity and pain. Norrbrink Budh et al reported that patients with AIS D had a significantly higher prevalence of SCI-related pain than patients with other AIS classifications.⁴⁷ In contrast, Werhagen et al showed patients with complete SCI (AIS A) had more below-level NeuP than patients with incomplete SCI.³¹ Another study of Werhagen showed no difference between injury severity and the prevalence of NeuP in non-traumatic SCI.³² Siddall et al reported that allodynia was more present in patients with incomplete lesions.⁴⁹ A better understanding of the relationship between pain and injury severity can

possibly be achieved if the pathophysiology of pain is further explained or a more in depth assessment of pain characteristics is performed.

Study limitations

Several limitations of this study deserve to be mentioned. First, this study only reports on pain during initial inpatient rehabilitation. Therefore the prevalence of pain can only be generalized to patients during inpatient rehabilitation, with a short time since onset of SCI. Second, this study describes presence on pain, type of pain, type of treatment, and NRS but does not report on further information such as type of medication or burden caused by pain. Third, the DSCID has a high percentage of missing data. Because data collection is either done prospectively at admission or by extracting information from the electronic medical record, the amount of missing data could have been caused by a lack of information in the electronic medical record. Also, in most cases there is a delay in entering data in the DSCID, making it difficult to retrieve missing data from the patient. Future funding could possibly create time in which data are collected routinely prospectively in the different centers. This could enhance data completeness and accuracy as has been described before.¹⁹ Last, there is a significant difference in AIS classifications in the

Table 7 Prevalence of neuropathic pain in relationship to patient and lesion characteristics

	Admission			Discharge		
	Patients With Pain (n=411) n (%) [*]	Patients With Pain vs Without Pain		Patients With Pain (n=436), n (%) [*]	Patients With Pain vs Without Pain	
		Unadjusted OR (95% CI)	Adjusted OR [†] (95% CI)		Unadjusted OR (95% CI)	Adjusted OR [†] (95% CI)
Sex						
Female	158 (32.6)	1.20 (0.94-1.52)	1.35 (1.04-1.76) [‡]	183 (37.2)	1.51 (1.19-1.90) [‡]	1.69 (1.30-2.20) [‡]
Male	253 (28.8)	1.00	1.00	253 (28.2)	1.00	1.00
Missing	0			0		
Age (y)						
≤44	103 (34.6)	1.59 (1.04-2.41) [‡]	1.56 (0.98-2.48)	98 (31.7)	1.50 (0.98-2.29)	1.89 (1.16-3.07) [‡]
45-54	82 (35.7)	1.66 (1.07-2.58) [‡]	1.54 (0.96-2.47)	83 (36.6)	1.86 (1.19-2.89) [‡]	2.23 (1.35-3.67) [‡]
55-64	84 (28.3)	1.18 (0.77-1.81)	1.07 (0.67-1.70)	102 (33.8)	1.64 (1.08-2.51) [‡]	2.08 (1.29-3.37) [‡]
65-74	96 (27.0)	1.11 (0.73-1.68)	0.98 (0.62-1.53)	108 (29.4)	1.34 (0.89-2.04)	1.55 (0.97-2.48)
≥75	43 (25.0)	1.00	1.00	41 (23.7)	1.00	1.00
Missing	3			4		
AIS classification at admission						
AIS A	47 (30.1)	0.98 (0.68-1.43)	0.82 (0.54-1.24)	54 (34.0)	1.18 (0.82-1.70)	1.08 (0.71-1.65)
AIS B	31 (25.8)	0.79 (0.51-1.23)	0.74 (0.47-1.17)	38 (30.9)	1.03 (0.68-1.55)	1.02 (0.66-1.58)
AIS C	50 (28.4)	0.91 (0.63-1.30)	0.85 (0.58-1.24)	53 (30.5)	1.01 (0.71-1.44)	0.86 (0.58-1.27)
AIS D	239 (30.5)	1.00	1.00	243 (30.3)	1.00	1.00
Missing	44			48		
Etiology of the SCI						
Traumatic	176 (32.7)	1.22 (0.96-1.55)	1.31 (0.98-1.77)	194 (35.3)	1.34 (1.06-1.69) [‡]	1.31 (0.98-1.74)
Nontraumatic	229 (28.5)	1.00	1.00	238 (29.0)	1.00	1.00
Missing	6			4		
Level of injury						
Tetraplegia	168 (31.1)	1.13 (0.89-1.45)	1.09 (0.83-1.44)	185 (34.1)	1.30 (1.02-1.65) [‡]	1.32 (1.00-1.74) [‡]
Paraplegia	194 (28.5)	1.00	1.00	201 (28.6)	1.00	1.00
Missing	49			50		

Abbreviations: CI, confidence interval; OR, odds ratio,

* Percentage of pain within subgroup.

[†] Adjusted for sex, age, time since onset, AIS classification, etiology of SCI, and level of injury.

[‡] Statistically significant (CI≠1.00).

included and excluded group. Patients with complete SCI were less likely to have data on presence of pain at both admission and discharge. This could be explained by the fact that patients with complete SCI are admitted longer to the rehabilitation center and the information at discharge is not yet collected. In addition, there are more patients with AIS D in the analysis, which could have led to an overestimation of the prevalence of NocP because participants with AIS D were more likely to have NocP than those with AIS B and C.

Conclusions

This study showed that SCI-related pain is highly prevalent during initial inpatient rehabilitation. Prevalence of NocP decreased during inpatient rehabilitation, and prevalence of NeuP remained the same. Pain intensity was similar for different types of SCI-related pain, with a score on the 11-point NRS with means ranging from 4.19 (NocP at discharge) to 4.69 (NeuP at admission). Female patients and patients with traumatic SCI have a higher risk of having NocP, and female patients, patients younger than 65 years, and patients with tetraplegia have a higher risk of having NeuP. Patients with AIS B and C have a lower risk of having NocP than those with AIS D. These differences in pain prevalence between

different groups of patients with SCI are important in screening patients on having pain and developing pain during inpatient rehabilitation.

Suppliers

a. SPSS Version 26; IBM, Armonk NY.

Keywords

Neuralgia; Nociceptive pain; Pain; Rehabilitation; Spinal cord injuries

Corresponding author

J. Stolwijk, Rembrandtkade, MD, PhD, 10, 3583 TM, Utrecht, the Netherlands. *E-mail address:* j.stolwijk@dehoogstraat.nl.

Acknowledgments

We thank these rehabilitation centers and the people involved in data collection.

References

- Nas K, Yazmalar L, Şah V, Aydın A, Öneş K. Rehabilitation of spinal cord injuries. *World J Orthop* 2015;6:8–16.
- Hagen EM, Rekand T. Management of neuropathic pain associated with spinal cord injury. *Pain Ther* 2015;4:51–65.
- Siddall PJ, McClelland JM, Rutkowski SB, Cousins MJ. A longitudinal study of the prevalence and characteristics of pain in the first 5 years following spinal cord injury. *Pain* 2003;103:249–57.
- Van Gorp S, Kessels AG, Joosten EA, Van Kleef M, Patijn J. Pain prevalence and its determinants after spinal cord injury: a systematic review. *Eur J Pain* 2015;19:5–14.
- Siddall PJ. Management of neuropathic pain following spinal cord injury: now and in the future. *Spinal Cord* 2009;47:352–9.
- Nicholson B. Differential diagnosis: nociceptive and neuropathic pain. *Am J Manag Care* 2006;12(9 Suppl):S256–62.
- Bryce TN, Biering-Sørensen F, Finnerup NB, et al. International spinal cord injury pain classification: part I. Background and description. *Spinal Cord* 2012;50:413–7.
- Burke D, Fullen BM, Stokes D, Lennon O. Neuropathic pain prevalence following spinal cord injury: a systematic review and meta-analysis. *Eur J Pain* 2017;21:29–44.
- Finnerup NB, Haroutounian S, Kamerman P, et al. Neuropathic pain: an updated grading system for research and clinical practice. *Pain* 2016;157:1599–606.
- Baron R, Binder A, Wasner G. Neuropathic pain: diagnosis, pathophysiological mechanisms, and treatment. *Lancet Neurol* 2010;9:807–19.
- Teasell RW, Mehta S, Aubut J-AL, et al. A systematic review of pharmacologic treatments of pain after spinal cord injury. *Arch Phys Med Rehabil* 2010;91:816–31.
- Mehta S, McIntyre A, Janzen S, et al. Systematic Review of Pharmacologic Treatments of Pain After Spinal Cord Injury: An Update. *Arch Phys Med Rehabil* 2016;97: 1381-91.e1.
- Max M. World Health Organization. World Health Organization cancer pain relief program: network news. *J Pain Symptom Manage* 1986;1:53–7.
- Franz S, Schulz B, Wang H, et al. Management of pain in individuals with spinal cord injury: guideline of the German-speaking Medical Society for Spinal Cord Injury. *Ger Med Sci* 2019;17:1–26.
- Guy SD, Mehta S, Casalino A, et al. The CanPain SCI Clinical Practice Guidelines for Rehabilitation Management of Neuropathic Pain after Spinal Cord: recommendations for treatment. *Spinal Cord* 2016;54(Suppl 1):S14–23.
- Boldt I, Eriks-Hoogland I, Brinkhof MWG, de Bie R, Joggi D, von Elm E. Non-pharmacological interventions for chronic pain in people with spinal cord injury. *Cochrane Database Syst Rev* 2014;11: CD009177.
- Zanca JM, Dijkers MP, Hammond FM, Horn SD. Pain and its impact on inpatient rehabilitation for acute traumatic spinal cord injury: analysis of observational data collected in the SCIREhab Study. *Arch Phys Med Rehabil* 2013;94(4 Suppl):S137–44.
- Stampacchia G, Gerini A, Morganti R, et al. Pain characteristics in Italian people with spinal cord injury: a multicentre study. *Spinal Cord* 2022;60:604–11.
- Post MWM, Nachttegaal J, Van Langeveld SA, et al. Progress of the Dutch Spinal Cord Injury Database: completeness of database and profile of patients admitted for inpatient rehabilitation in 2015. *Top Spinal Cord Inj Rehabil* 2018;24:141–50.
- Kirschblum SC, Waring W, Biering-Sørensen F, et al. Reference for the 2011 revision of the International Standards for Neurological Classification of Spinal Cord Injury. *J Spinal Cord Med* 2011;34:547–54.
- Widerström-Noga E, Biering-Sørensen F, Bryce TN, et al. The International Spinal Cord Injury Pain Basic Data Set (version 2.0). *Spinal Cord* 2014;52:282–6.
- Hunt C, Moman R, Peterson A, et al. Prevalence of chronic pain after spinal cord injury: a systematic review and meta-analysis. *Reg Anesth Pain Med* 2021;46:328–36.
- Finnerup NB, Bastrup C. Spinal cord injury pain: mechanisms and management. *Curr Pain Headache Rep* 2012;16:207–16.
- Wellisch M, Lovett K, Harrold M, et al. Treatment of shoulder pain in people with spinal cord injury who use manual wheelchairs: a systematic review and meta-analysis. *Spinal Cord* 2022;60:107–14.
- Jahani O, Van Straaten MG, Barlow JD, Murthy NS, Morrow MMB. Progression of rotator cuff tendon pathology in manual wheelchair users with spinal cord injury: a 1-year longitudinal study. *J Spinal Cord Med* 2022 Apr 14. [Epub ahead of print].
- Adriaansen JJE, Post MWM, de Groot S, et al. Secondary health conditions in persons with spinal cord injury: a longitudinal study from one to five years post-discharge. *J Rehabil Med* 2013;45:1016–22.
- Aito S, D'Andrea M, Werhagen L, et al. Neurological and functional outcome in traumatic central cord syndrome. *Spinal Cord* 2007;45:292–7.
- Finnerup NB, Norrbrink C, Trok K, et al. Phenotypes and predictors of pain following traumatic spinal cord injury: a prospective study. *J Pain* 2014;15:40–8.
- New PW, Lim TC, Hill ST, Brown DJ. A survey of pain during rehabilitation after acute spinal cord injury. *Spinal Cord* 1997;35:658–63.
- Wen H, Reinhardt JD, Gosney JE, Baumberger M, Zhang X, Li J. Spinal cord injury-related chronic pain in victims of the 2008 Sichuan earthquake: a prospective cohort study. *Spinal Cord* 2013;51:857–62.
- Werhagen L, Budh CN, Hultling C, Molander C. Neuropathic pain after traumatic spinal cord injury—relations to gender, spinal level, completeness, and age at the time of injury. *Spinal Cord* 2004;42:665–73.
- Werhagen L, Hultling C, Molander C. The prevalence of neuropathic pain after non-traumatic spinal cord lesion. *Spinal Cord* 2007;45:609–15.
- Werhagen L, Aito S, Tucci L, Strayer J, Hultling C. 25 years or more after spinal cord injury: clinical conditions of individuals in the Florence and Stockholm areas. *Spinal Cord* 2012;50:243–6.
- Zeilig G, Enosh S, Rubin-Asher D, Lehr B, Defrin R. The nature and course of sensory changes following spinal cord injury: predictive properties and implications on the mechanism of central pain. *Brain* 2012;135:418–30.
- Barrett H, McClelland JM, Rutkowski SB, Siddall PJ. Pain characteristics in patients admitted to hospital with complications after spinal cord injury. *Arch Phys Med Rehabil* 2003;84:789–95.
- Vervoordeldonk JJ, Post MWM, New P, Clin Epi M, Van Asbeck FWA. Rehabilitation of patients with nontraumatic spinal cord injury in the Netherlands: etiology, length of stay, and functional outcome. *Top Spinal Cord Inj Rehabil* 2013;19:195–201.
- Osterthun R, Post MWM, van Asbeck FWA. Characteristics, length of stay and functional outcome of patients with spinal cord injury in Dutch and Flemish rehabilitation centres. *Spinal Cord* 2009;47:339–44.
- Ones K, Yilmaz E, Beydogan A, Gultekin O, Caglar N. Comparison of functional results in non-traumatic and traumatic spinal cord injury. *Disabil Rehabil* 2007;29:1185–91.
- McKinley WO, Seel RT, Hardman JT. Nontraumatic spinal cord injury: incidence, epidemiology, and functional outcome. *Arch Phys Med Rehabil* 1999;80:619–23.
- Schönherr MC, Groothoff JW, Mulder GA, Eisma WH. Rehabilitation of patients with spinal cord lesions in the Netherlands: an epidemiological study. *Spinal Cord* 1996;34:679–83.
- van den Berg MEL, Castellote JM, Mahillo-Fernandez I, de Pedro-Cuesta J. Incidence of nontraumatic spinal cord injury: a Spanish cohort study (1972-2008). *Arch Phys Med Rehabil* 2012;93:325–31.
- New PW, Simmonds F, Stevermuer T. A population-based study comparing traumatic spinal cord injury and non-traumatic spinal cord injury using a national rehabilitation database. *Spinal Cord* 2011;49:397–403.
- Ge L, Arul K, Ikpeze T, Baldwin A, Nickels JL, Mesfin A. Traumatic and nontraumatic spinal cord injuries. *World Neurosurg* 2018;111: e142–8.

44. Lee BB, Cripps RA, Fitzharris M, Wing PC. The global map for traumatic spinal cord injury epidemiology: update 2011, global incidence rate. *Spinal Cord* 2014;52:110–6.
45. Kim HY, Lee HJ, Kim T, et al. Prevalence and characteristics of neuropathic pain in patients with spinal cord injury referred to a rehabilitation center. *Ann Rehabil Med* 2020;44:438–49.
46. Müller R, Brinkhof MWG, Arnet U, et al. Prevalence and associated factors of pain in the Swiss spinal cord injury population. *Spinal Cord* 2017;55:346–54.
47. Norrbrink Budh C, Lund I, Ertzgaard P, et al. Pain in a Swedish spinal cord injury population. *Clin Rehabil* 2003;17:685–90.
48. Casale R, Atzeni F, Bazzichi L, et al. Pain in women: a perspective review on a relevant clinical issue that deserves prioritization. *Pain Ther* 2021;10:287–314.
49. Siddall PJ, Taylor DA, McClelland JM, Rutkowski SB, Cousins MJ. Pain report and the relationship of pain to physical factors in the first 6 months following spinal cord injury. *Pain* 1999;81:187–97.