

ORGANIC COMPOUNDS

Acta Cryst. (1994). C50, 410–411

**1-*tert*-Butyl-2-(*tert*-butylamino-methylidene)-3-pyrrolidinone,
C₁₃H₂₄N₂O**

ANTHONY L. SPEK

Bijvoet Center for Biomolecular Research,
Department of Crystal and Structural Chemistry,
Utrecht University, Padualaan 8, 3584 CH Utrecht,
The Netherlands

MAURITS D. JANSSEN, ELMO WISSING AND G. VAN KOTEN

Debye Institute, Department of Metal-Mediated
Synthesis, Utrecht University, Padualaan 8,
3584 CH Utrecht, The Netherlands


(Received 30 July 1993; accepted 27 September 1993)

Abstract

The eneamine moiety on the pyrrolidinone ring is coplanar to the carbonyl function, with an *E* configuration stabilized by an intramolecular N—H···O bond.

Comment

The molecular structure of the title compound (I) contains a pyrrolidinone ring with the eneamine fragment in the *E* configuration stabilized by an intramolecular N—H···O bond [N—H 0.92 (2), H···O 2.02 (2), N···O 2.754 (3) Å; N—H···O 135 (2)°]. The eneamine system is fully conjugated with the carbonyl function, as indicated by the short '*t*Bu(H)N—C single bond [1.334 (3) Å] and the relatively long olefinic C=C bond [1.360 (4) Å]. The H—N—C=C—C=O moiety is planar. This confirms the NMR data as well as the assignment of the structure in solution of the isomer of the parent compound, 1-*tert*-butyl-5-(*tert*-butylimino)-2-pyrrolidinone.



The pyrrolidinone ring is slightly puckered. The Cremer & Pople (1975) puckering parameters [$Q(2) = 0.223 (3)$ Å, $\varphi(2) = 338.1 (8)$ °] indicate a half-chair conformation. The twist about an approximate twofold axis running through C=O is illustrated most clearly by

the low $C_2(C7)$ asymmetry parameter [1.3 (3)°] (Duax, Weeks & Rohrer, 1976).

The geometry of both *tert*-butyl moieties deviates significantly from trigonal symmetry, due to intramolecular crowding [e.g. the *trans* bond angle N1—C1—C13, 105.8 (2)°, is significantly smaller than the two *gauche* bond angles N1—C1—C11, 109.9 (2)°, and N1—C1—C12, 110.2 (2)°]. Similar effects have been reported by de Vos, Kroon, Brouwer & Bos (1981). The structure contains no residual solvent-accessible areas.




Fig. 1. Structure of 1-*tert*-butyl-2-(*tert*-butylaminomethylidene)-3-pyrrolidinone, with displacement ellipsoids at 50% probability.

Experimental

The preparation of the title compound was performed according to the procedure described by Wissing, Kleijn, Boersma & Van Koten (1993). The crystal was mounted on top of a glass fibre.

Crystal data

C ₁₃ H ₂₄ N ₂ O	Mo $K\alpha$ radiation
$M_r = 224.35$	$\lambda = 0.71073$ Å
Monoclinic	Cell parameters from 25 reflections
$P2_1/c$	$\theta = 9\text{--}20^\circ$
$a = 19.646 (2)$ Å	$\mu = 0.064$ mm ⁻¹
$b = 6.1490 (10)$ Å	$T = 295$ K
$c = 11.466 (2)$ Å	Plate shaped
$\beta = 95.250 (10)$ °	0.50 × 0.35 × 0.35 mm
$V = 1379.3 (4)$ Å ³	Yellowish
$Z = 4$	
$D_x = 1.080$ Mg m ⁻³	

Data collection

Enraf-Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.08$
$\omega/2\theta$ scans [$\Delta\omega = (1.00 + 0.35\tan\theta)$ °]	$\theta_{\text{max}} = 27.5^\circ$
Absorption correction: none	$h = -25 \rightarrow 25$
5152 measured reflections	$k = -7 \rightarrow 0$
3170 independent reflections	$l = 0 \rightarrow 14$
1665 observed reflections [$I > 2.5\sigma(I)$]	3 standard reflections frequency: 30 min intensity variation: 5%

Refinement

Refinement on F	$w = 1/\sigma^2(F)$
$R = 0.0555$	$(\Delta/\sigma)_{\text{max}} = 0.33$
$wR = 0.0478$	$\Delta\rho_{\text{max}} = 0.18 \text{ e } \text{\AA}^{-3}$
$S = 0.62$	$\Delta\rho_{\text{min}} = -0.17 \text{ e } \text{\AA}^{-3}$
1665 reflections	Atomic scattering factors
221 parameters	from Cromer & Mann
Only coordinates of H atoms	(1968)
refined	

(SON) with financial aid from the Netherlands Organization of Scientific Research (NWO)

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the British Library Document Supply Centre as Supplementary Publication No. SUP 71668 (26 pp.). Copies may be obtained through The Technical Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England. [CIF reference: AB1111]

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (\AA^2)

$$U_{\text{eq}} = (1/3) \sum_i \sum_j U_{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j.$$

	x	y	z	U_{eq}
O1	0.30252 (11)	0.1314 (3)	0.43349 (18)	0.0695 (8)
N1	0.33334 (11)	0.5126 (4)	0.55298 (19)	0.0448 (7)
N2	0.18186 (10)	0.5417 (3)	0.33521 (15)	0.0383 (7)
C1	0.38595 (12)	0.6454 (4)	0.6193 (2)	0.0440 (9)
C2	0.28109 (13)	0.5892 (5)	0.4823 (2)	0.0385 (8)
C3	0.23989 (12)	0.4619 (4)	0.40900 (19)	0.0351 (8)
C4	0.11669 (12)	0.5560 (4)	0.3914 (2)	0.0409 (8)
C5	0.17865 (16)	0.3890 (5)	0.2327 (2)	0.0510 (10)
C6	0.21343 (17)	0.1794 (5)	0.2721 (3)	0.0575 (11)
C7	0.25757 (14)	0.2448 (5)	0.3818 (2)	0.0462 (9)
C11	0.43035 (17)	0.7569 (7)	0.5357 (3)	0.0662 (13)
C12	0.35274 (19)	0.8132 (6)	0.6923 (3)	0.0617 (11)
C13	0.42781 (19)	0.4880 (6)	0.6989 (3)	0.0670 (14)
C41	0.06300 (16)	0.6632 (6)	0.3055 (3)	0.0571 (11)
C42	0.12800 (18)	0.7038 (6)	0.4984 (3)	0.0605 (11)
C43	0.09049 (17)	0.3363 (5)	0.4311 (3)	0.0592 (11)

Table 2. Selected geometric parameters (\AA , $^\circ$)

O1—C7	1.234 (3)	C1—C13	1.519 (4)
N1—C1	1.473 (3)	C2—C3	1.360 (4)
N1—C2	1.334 (3)	C3—C7	1.421 (4)
N2—C3	1.442 (3)	C4—C41	1.525 (4)
N2—C4	1.488 (3)	C4—C42	1.527 (4)
N2—C5	1.501 (3)	C4—C43	1.530 (4)
C1—C11	1.517 (4)	C5—C6	1.508 (4)
C1—C12	1.514 (4)	C6—C7	1.515 (4)
C1—N1—C2	125.6 (2)	C2—C3—C7	122.2 (2)
C3—N2—C4	115.56 (17)	N2—C4—C41	108.5 (2)
C3—N2—C5	102.76 (19)	N2—C4—C42	108.4 (2)
C4—N2—C5	113.57 (19)	N2—C4—C43	113.8 (2)
N1—C1—C11	109.9 (2)	C41—C4—C42	107.6 (2)
N1—C1—C12	110.2 (2)	C41—C4—C43	110.0 (2)
N1—C1—C13	105.8 (2)	C42—C4—C43	108.3 (2)
C11—C1—C12	110.1 (3)	N2—C5—C6	108.3 (2)
C11—C1—C13	110.9 (2)	C5—C6—C7	103.4 (2)
C12—C1—C13	109.8 (2)	O1—C7—C3	127.2 (2)
N1—C2—C3	123.6 (3)	O1—C7—C6	125.4 (3)
N2—C3—C2	124.1 (2)	C3—C7—C6	107.4 (2)
N2—C3—C7	112.8 (2)		

Positional parameters of the H atoms were refined with four common isotropic thermal parameters. Data collection: locally modified CAD-4 Software (Enraf-Nonius, 1989). Cell refinement: SET4 (de Boer & Duisenberg, 1984). Data reduction: HELENA (Spek, 1993). Program(s) used to solve structure: SHELXS86 (Sheldrick, 1986). Program(s) used to refine structure: SHELX76 (Sheldrick, 1976). Molecular graphics: PLATON (Spek, 1990). Software used to prepare material for publication: PLATON.

X-ray data were collected by A. J. M. Duisenberg. The investigations were supported in part (EW and ALS) by the Netherlands Foundation for Chemical Research

References

- Boer, J. L. de & Duisenberg, A. J. M. (1984). *Acta Cryst. A* **40**, C-410.
 Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
 Cromer, D. T. & Mann, J. B. (1968). *Acta Cryst. A* **24**, 321–324.
 Duax, W. L., Weeks, C. M. & Rohrer, D. C. (1976). *Topics in Stereochemistry*, Vol. 9, edited by N. L. Allenger & E. L. Eliel, pp. 271–383. New York: John Wiley.
 Enraf-Nonius (1989). CAD-4 Software. Version 5.0. Enraf-Nonius, Delft, The Netherlands.
 Sheldrick, G. M. (1976). SHELX76. Program for Crystal Structure Determination. Univ. of Cambridge, England.
 Sheldrick, G. M. (1986). SHELXS86. Program for the Solution of Crystal Structures. Univ. of Göttingen, Germany.
 Spek, A. L. (1990). *Acta Cryst. A* **46**, C-34.
 Spek, A. L. (1993). HELENA. Program for Data Reduction. Laboratorium voor Kristal- en Structuurchemie. Univ. of Utrecht, The Netherlands.
 Vos, A. M. de, Kroon, J., Brouwer, A. C. & Bos, H. J. T. (1981). *Acta Cryst. B* **37**, 161–164.
 Wissing, E., Kleijn, H., Boersma, J. & Van Koten, G. (1993). *Recl Trav. Chim. Pays-Bas.* **112**, 618–626.

Acta Cryst. (1994). **C50**, 411–413

2,6-Difluorophenylxenon(II)-tetrafluoroborat

THEO GILLES, ROBERT GNANN, DIETER NAUMANN
UND KARL-FRIEDRICH TEBBE

Institut für Anorganische Chemie, Universität zu Köln,
Greinstraße 6, D-50939 Köln, Germany

(Eingegangen am 23. Februar 1993; angenommen am 17. September 1993)

Abstract

The crystal structure of the compound 2,6-difluorophenylxenon(II) tetrafluoroborate is described. The Xe atom is bonded to a C atom and weakly coordinated by one F atom of the tetrafluoroborate anion.

Kommentar

Seit kurzem kennt man Verbindungen mit Xenon-Kohlenstoff Bindungen (Naumann, Butler, Gnann & Tyrra, 1993 und dort genannte Literatur), von denen eine bereits durch