

## References

- [1] J. P. Buhler, B. H. Gross, D. B. Zagier, On the conjecture of Birch and Swinnerton-Dyer for an elliptic curve of rank 3, *Math. Comp.* **44** (1985), 473–481.
- [2] A. Brumer, K. Kramer, The rank of elliptic curves, *Duke Math. J.* **44** (1977), 715–743.
- [3] Z. Djabri, E. F. Schaefer, N. P. Smart, Computing the  $p$ -Selmer group of an elliptic curve, to appear in *Trans. Amer. Math. Soc.*
- [4] S. Fermigier, An elliptic curve over  $\mathbf{Q}$  of rank  $\geq 22$ , [www.fermigier.com](http://www.fermigier.com).
- [5] L. J. Mordell, On the rational solutions of the indeterminate equations of the 3rd and 4th degrees, *Proc. Camb. Phil. Soc.* **21** (1922), 179–192.
- [6] L. J. Mordell, *Diophantine equations*, Academic Press, London and New York, 1969.
- [7] A. Néron, Modèles minimaux des variétés abéliennes sur les corps locaux et globaux, *IHES Publ. Math.*, **21** (1964), 361–482.
- [8] B. Poonen, E. F. Schaefer, Explicit descent for Jacobians of cyclic covers of the projective line, *J. reine und angew. Math.* **488** (1997), 141–188.
- [9] E. F. Schaefer, Class groups and Selmer groups, *J. Number theory* **56** (1996), 79–114.
- [10] E. F. Schaefer, 2-Descent on the Jacobians of hyperelliptic curves, *J. Number Th.* **51** (1995), 219–232.
- [11] E. F. Schaefer, Computing a Selmer group of a Jacobian using functions on the curve, *Math. Ann.* **310** (1998), 447–471.
- [12] J-P. Serre, Propriétés galoisiennes des points d’ordre fini des courbes elliptiques, *Inv. Math.* **15** (1972), 259–331.
- [13] J. H. Silverman, *The arithmetic of elliptic curves*, Graduate Texts in Math. **106**, Springer-Verlag, New York, 1986.
- [14] J. Top, Descent by 3-isogeny and the 3-rank of quadratic fields, In: *Advances in number theory* (ed. by F. Q. Gouvea and N. Yui), Clarendon Press, Oxford 1993, 303–317.